



Small Mass Limit of a Langevin Equation on a Manifold

Jeremiah Birrell, Scott Hottovy, Giovanni Volpe and Jan Wehr

Abstract. We study damped geodesic motion of a particle of mass m on a Riemannian manifold, in the presence of an external force and noise. Lifting the resulting stochastic differential equation to the orthogonal frame bundle, we prove that, as $m \rightarrow 0$, its solutions converge to solutions of a limiting equation which includes a *noise-induced drift* term. A very special case of the main result presents Brownian motion on the manifold as a limit of inertial systems.

1. Introduction

Brownian motion (BM) plays a central role in many phenomena of scientific and technological significance. It lies at the foundation of stochastic calculus [1], which is applied to model a variety of phenomena, ranging from non-equilibrium statistical mechanics to stock market fluctuations to population dynamics. In particular, Brownian motion occurs naturally in systems where microscopic and nanoscopic particles are present, as a consequence of thermal agitation [2]. Brownian motion of micro- and nanoparticles occurring in complex environments can often be represented as motion on two-dimensional or one-dimensional manifolds embedded within a three-dimensional space. For example, the motion of proteins on cellular membranes occurs effectively on two-dimensional manifolds and is currently at the center of an intense experimental activity [3]. The single file diffusion of particles in porous nanomaterials occurs in an effectively one-dimensional environment and plays a crucial role in many phenomena such as drug delivery, chemical catalysis and oil recovery [4]. Several interesting phenomena can emerge in these conditions, such as anomalous diffusion [5] and inhomogeneous diffusion [6]. Similar phenomena also occur when considering active matter systems, such as living matter, which are characterized by being in a far-from-equilibrium state [7]; such systems often interact with complex environments that can be effectively modeled by

low-dimensional manifolds embedded in a three-dimensional space. To gain a deeper understanding of these phenomena it is necessary to explore the properties of Brownian motion on manifolds.

The original motivation for this paper is to present Brownian motion on a manifold as the zero-mass limit of an inertial system. Our main result is significantly more general and contains a rigorous version of the above statement as a special case. In this section, we will first outline this motivating problem and then discuss some earlier work on similar questions. For the sake of clarity, we do not spell out all technical assumptions and the arguments presented in the introduction are heuristic.

Brownian motion on an n -dimensional Riemannian manifold (M, g) can be introduced as a mathematical object—a Markov process x_t on M with the generator expressed in terms of the Riemannian metric g , which uniquely determines its law. In this form, it has been a subject of an immense amount of study, both for its own sake and beauty, and for applications to analysis and geometry. The reader is referred to [8, 9] and references therein. In local coordinates, the components of BM satisfy the stochastic differential equation (SDE):

$$dx_t^i = -\frac{1}{2}g^{jk}\Gamma_{jk}^i dt + \sum_{\alpha=1}^n \sigma_\alpha^i dW_t^\alpha, \quad (1.1)$$

where σ is the positive-definite square root of the inverse metric tensor g^{-1} , in the sense that $\sum_{\alpha=1}^n \sigma_\alpha^i \sigma_\alpha^k = g^{ik}$, see p. 87 of [8]. From the applied point of view, BM on a manifold is an idealized probabilistic description of diffusive motion performed by a particle constrained to M . This can be justified at various levels, depending on what one is willing to assume. Let us mention in particular the work of van Kampen [10] which studies the conditions on the constraints, restricting motion of a diffusing particle to a manifold, under which its effective motion becomes Brownian.

Here, as our point of departure, we take equations describing inertial motion of a particle of mass m , in the presence of two forces: damping and noise. The equations of motion in local coordinates are

$$\begin{aligned} dx_t^i &= v_t^i dt, & (1.2) \\ m dv_t^i &= -m\Gamma_{jk}^i v_t^j v_t^k dt - \gamma_j^i v_t^j dt + \sum_{\alpha=1}^n \sigma_\alpha^i dW_t^\alpha, & (1.3) \end{aligned}$$

where Γ_{jk}^i are the Christoffel's symbols of the metric g , γ denotes the damping tensor and the vector fields σ_α , $\alpha = 1, \dots, n$, couple the particle to n standard Wiener processes, acting as noise sources. The summation convention is used here and throughout the paper. The reason the sum over α here is written explicitly is that it does not play the role of a covariant index.

For the purposes of this motivating discussion, we assume that the damping and noise satisfy a fluctuation–dissipation relation known from non-equilibrium statistical mechanics [11]; this assumption will not be needed in the more general theorem that will be presented later. Note that the covariance of

the noise is equal to $\sum_{\alpha} \sigma_{\alpha}^i \sigma_{\alpha}^k$ and is thus a tensor of type $\binom{2}{0}$ (a contravariant tensor of rank two). We want to relate it to a quantity of the same type. Since the damping tensor γ_j^i has type $\binom{1}{1}$, we raise its lower index using the metric and state the fluctuation–dissipation relation as:

$$\sum_{\alpha} \sigma_{\alpha}^i \sigma_{\alpha}^j = 2\beta^{-1} g^{jk} \gamma_k^i, \tag{1.4}$$

where $\beta^{-1} = k_B T$, k_B is the Boltzmann constant and T denotes the temperature. In particular, if damping is isotropic, $\gamma_k^i = \gamma \delta_k^i$, we get

$$\sum_{\alpha} \sigma_{\alpha}^i \sigma_{\alpha}^j = 2\beta^{-1} \gamma g^{ij} \tag{1.5}$$

and Eq. (1.3) becomes

$$m dv_t^i = -m \Gamma_{jk}^i v_t^j v_t^k dt - \gamma v_t^i dt + \sum_{\alpha} \sigma_{\alpha}^i dW_t^{\alpha}, \tag{1.6}$$

with σ_{α}^i satisfying the relation Eq. (1.5). The problem motivating this work can now be stated as follows. Consider the solutions of Eqs. (1.2) and (1.6) with the initial conditions $x^{(m)}(0) = x_0, v^{(m)}(0) = v_0$. We want to show that, as $m \rightarrow 0, x^{(m)}$ converges to the solution of the SDE

$$dx_t^i = -\frac{1}{\beta \gamma^2} g^{jk} \Gamma_{jk}^i dt + \frac{1}{\gamma} \sum_{\alpha} \sigma_{\alpha}^i dW_t^{\alpha} \tag{1.7}$$

with the same initial condition. Since this equation describes (a rescaled) BM on the manifold M in local coordinates, this will realize our original goal. Related results in the physics content are reported in [12].

We now present a sketch of the argument. The remarks that follow it explain its relation to the actual proof in later sections. Our guiding principle is that the kinetic energy of the particle is of order 1, so that the components of the velocity (in a fixed coordinate chart) behave as $\frac{1}{\sqrt{m}}$ in the limit $m \rightarrow 0$.

Solving for $v_t^i dt$ in Eq. (1.6), we obtain

$$dx_t^i = v_t^i dt = -\frac{m}{\gamma} dv_t^i - \frac{m}{\gamma} \Gamma_{jk}^i v_t^j v_t^k dt + \frac{1}{\gamma} \sum_{\alpha} \sigma_{\alpha}^i dW_t^{\alpha}. \tag{1.8}$$

In the limit $m \rightarrow 0$, we expect no contribution from the first term, since γ is constant and thus $\frac{m}{\gamma} dv_t^i$ is the differential of the expression $\frac{1}{\gamma} m v_t^i$ which vanishes in the limit. We *do* expect a nonzero limit from the quadratic term. This is again based on the analogy with [13], where such a term appears as a result of integration by parts. In the case discussed here, it is present in the equation from the start, reflecting the manifold geometry. In the limit, we expect the fast velocity variable to average, giving rise to an x -dependent drift term. To calculate this term, we use the method of [13], together with the heuristics that v^i should be of order $m^{-\frac{1}{2}}$. Consider the differential (vanishing in the $m \rightarrow 0$ limit):

$$\begin{aligned}
 d(mv_t^j mv_t^k) &= d(mv_t^j)mv_t^k + mv_t^j d(mv_t^k) + d(mv_t^j) d(mv_t^k) \\
 &= \left(-m\Gamma_{li}^j v_t^l v_t^i dt - \gamma v_t^j dt + \sum_{\alpha} \sigma_{\alpha}^j dW_t^{\alpha} \right) mv_t^k \\
 &\quad + mv_t^j \left(-m\Gamma_{li}^k v_t^l v_t^i dt - \gamma v_t^k dt + \sum_{\alpha} \sigma_{\alpha}^k dW_t^{\alpha} \right) + \sum_{\alpha} \sigma_{\alpha}^j \sigma_{\alpha}^k dt.
 \end{aligned}
 \tag{1.9}$$

Based on our assumption about the order of magnitude of v^i , in the limit we get

$$-2m\gamma v_t^j v_t^k dt + \sum_{\alpha} \sigma_{\alpha}^j \sigma_{\alpha}^k dt = 0.
 \tag{1.10}$$

Substituting this into Eq. (1.8) and leaving out the terms which vanish in the limit, we obtain

$$dx_t^i = -\frac{1}{\beta\gamma^2} g^{jk} \Gamma_{jk}^i dt + \frac{1}{\gamma} \sum_{\alpha} \sigma_{\alpha}^i dW_t^{\alpha},
 \tag{1.11}$$

which describes a rescaled BM on the manifold.

The limiting process x_t satisfies an equation driven by the same Wiener processes, W^{α} , that drove the equations for the original processes $x^{(m)}$, and is thus defined on the same probability space. The processes $x^{(m)}$ will be proven to converge to x in the sense that the L^p -norm (in the ω variable) of the uniform distance on $[0, T]$ between the realizations of x and $x^{(m)}$ goes to zero for every T . This is much stronger than convergence in law on compact time intervals.

Zero-mass limits of diffusive systems have been studied in numerous works, starting from [14]. See [2] for a masterly review of the early history. The analysis of such models have been extended in many directions. For example [15] studies the limit of a particle system driven by a fluid model that is coupled to noise. The extension of diffusion processes to the relativistic setting has been studied in, for example [16, 17]. Other related works study random perturbations of the geodesic flow on a Riemannian manifold. In [18], convergence of the transition semigroups of a family of transport processes on a manifold to that of Brownian motion was shown. See also the paper [19] where homogenization of the velocity variable for equations on manifolds is studied. Families of Ornstein–Uhlenbeck processes on manifolds were studied in [20, 21]. Two interesting recent papers are [22] and [23]. They prove convergence in law to Brownian motion in appropriate limits. We remark that in the special case of constant damping (as considered in this introduction), the generator of the process defined by Eqs. (1.2)–(1.3) is the *hypoelliptic Laplacian*, introduced in [24]—an important analytical object, encoding geometric properties of the manifold. In this case [24] proves a convergence-in-law result. This has been extended to cover the convergence of kernels and their derivatives [25].

More recently, several authors address the case when a general position-dependent forcing is included and the damping and/or noise coefficients depend on the state of the system. In particular, they study the associated phenomenon

of the *noise-induced drift* that arises in the limit. See references in the recent paper [13], where a formula for the noise-induced drift has been established for a large class of systems in Euclidean space of an arbitrary dimension. See also [26], where some of the assumptions made in [13] are relaxed.

The analysis applied here is most similar to that of [13], with two important differences. First, we prove the fundamental momentum (or kinetic energy) bound in a different way, which would also lead to an alternative proof of the main result of [13]. Second, in this paper we pose the problem on a compact Riemannian manifold. This leads to complications of a geometric nature that are absent in Euclidean space: to control the quadratic terms in the geodesic equation more efficiently, we lift the equations to the orthogonal frame bundle. Another, equally important, consequence of lifting to the orthogonal frame bundle is that the equations of motion, including the noise term, can be formulated geometrically, without reference to local coordinate charts. While this increases the number of variables and makes the equations more complicated, it simplifies the analytical aspects of the problem. Frame bundle techniques similar to this have been used by many other authors, see for example [8, 17, 20–22].

As our main result, we derive the equation satisfied by an inertial system in the limit $m \rightarrow 0$. In the case of constant damping and noise (and thus satisfying the fluctuation–dissipation relation) with zero forcing, we obtain Brownian motion, as suggested by the informal derivation leading to Eq. (1.11). Various formulations that cover this classical case have been studied by previous authors [21, 24]. Forcing terms were also considered in [21]. The theorem presented here is general enough to include the classical case, as well as cover position-dependent forcing, damping and noise, including a derivation of the formula for the noise-induced drift. In addition, the damping and the noise coefficients are not required to satisfy a fluctuation–dissipation relation (unlike in the motivating discussion above), which leads to a fully general formula for the resulting noise-induced drift in the small mass limit. Physically, this result is particularly relevant when considering active matter and systems far from thermodynamic equilibrium [27].

1.1. Summary of the Main Result

In this section, we summarize the main result of this paper, including the required assumptions. Motivation for the equations under consideration, further detail on the notation, and a proof of the result will follow in subsequent sections.

Let (M, g) be a compact, connected n -dimensional Riemannian manifold without boundary, $F_O(M)$ be its frame bundle with canonical projection π (see Sect. 3), and $N \equiv F_O(M) \times \mathbb{R}^n$. For each $m > 0$ (representing the particle mass), we will consider the following SDE on N for fixed, non-random initial condition $(u_0, v_0) \in N$:

$$u_t^m = u_0 + \int_{t_0}^t H_{v_s^m}(u_s^m) ds, \tag{1.12}$$

$$v_t^m = v_0 + \frac{1}{m} \int_{t_0}^t [F(u_s^m) - \gamma(u_s^m)v_s^m] ds + \frac{1}{m} \int_{t_0}^t \sigma(u_s^m) dW_s. \quad (1.13)$$

Here, $H_\alpha(u)$ are the canonical horizontal vector fields on $F_O(M)$ (see Lemma 3.2), $F(x)$ is a smooth vector field on M (the forcing), $\gamma(x)$ is a smooth $\binom{1}{1}$ tensor field on M (the damping), the noise coefficients are given by a $\mathbb{R}^{n \times k}$ -valued function, $\sigma(u)$, on $F_O(M)$, W_t is a \mathbb{R}^k -valued Wiener process, and we define $F(u) = u^{-1}F(\pi(u))$ and $\gamma(u) = u^{-1}\gamma(\pi(u))u$.

Let (u_t^m, v_t^m) be a family of solutions to Eqs. (1.12)–(1.13), corresponding to mass values $m > 0$. To study the $m \rightarrow 0$ limit of u_t^m , we assume that the symmetric part of $\gamma, \gamma^s = \frac{1}{2}(\gamma + \gamma^T)$, has eigenvalues bounded below by a positive constant. This coercivity assumption is crucial to our results, as it provides the damping necessary to ensure that the momentum degrees of freedom, $p_t^m = mv_t^m$, become negligible in the limit (see Sect. 5).

Under the assumptions stated above, the main result (Theorem 7.1) gives the limiting behavior of u_t^m as $m \rightarrow 0$. More specifically, we prove the following:

Fix $T > 0$ and a Riemannian metric tensor field on $F_O(M)$. Let d be the associated metric on the connected component of $F_O(M)$ that contains u_0 . Then for any $q > 0$ and any $0 < \kappa < q/2$, we have

$$E \left[\sup_{t \in [0, T]} d(u_t^m, u_t)^q \right] = O(m^\kappa) \text{ as } m \rightarrow 0, \quad (1.14)$$

where u_t solve the following SDE on $F_O(M)$ with initial condition u_0 :

$$du_t = H_{(\gamma^{-1}F)(u_t)}(u_t)dt + S(u_t)dt + H_{(\gamma^{-1}\sigma)(u_t)}(u_t) \circ dW_t. \quad (1.15)$$

The additional drift vector field, $S(u)$, on $F_O(M)$ that arises in the limit, given by Eqs. (6.32)–(6.33), will be called the *noise-induced drift*.

2. Forced Geodesic Motion on the Tangent Bundle

We now begin the task of making the results outlined in the introduction and summary precise. Let (M, g) be an n -dimensional smooth connected Riemannian manifold with tangent bundle (TM, π) , where π is the natural projection. Let $V : TM \rightarrow TM$ be smooth and $\pi \circ V = \pi$, i.e., V maps each fiber into itself. The deterministic dynamical system that we eventually want to couple to noise is defined by the equation

$$\nabla_{\dot{x}} \dot{x} = V(\dot{x}), \quad (2.1)$$

where ∇ is the Levi–Civita connection. In this section and the next, we focus on the non-random system. The coupling to noise will be discussed in Sect. 4. We refer to the system Eq. (2.1) as the geodesic equation with (velocity-dependent) forcing V . Note that \dot{x} is an element of TM , and so it contains both position and velocity information. In particular, Eq. (2.1) contains the special case where V is independent of the velocity degrees of freedom, i.e., V is a vector field on M .

Equation (2.1) is more general than the system outlined in the introduction. In particular, it contains a deterministic forcing, other than the drag. The process that we will eventually find in the small mass limit will, therefore, be more general than Brownian motion on M , but will include Brownian motion as a special case.

We now interpret Eq. (2.1) as an ordinary differential equation (ODE) on the tangent bundle TM . With $V \equiv 0$ it is the standard geodesic equation. Below we give some facts, starting from this case in points 1–4 and then, in points 5 and 6, we add forcing. These facts will not be used in our subsequent analysis, but they give one an idea of how forced geodesic motion on manifold, Eq. (2.1), can be reformulated as a flow on a larger space. We will build on this idea in the next section.

1. For $v \in TM$, let x_v be the geodesic with velocity v at $t = 0$. Define the geodesic vector field $G : TM \rightarrow T(TM)$ by $G(v) = \frac{d}{dt}(\dot{x}_v)_{t=0}$, i.e., the tangent vector to the curve $\dot{x}_v : I \rightarrow TM$ at $t = 0$. G is a smooth vector field on TM and $x : I \rightarrow M$ is a geodesic iff \dot{x} (interpreted as a curve in TM) is an integral curve of G .
2. If η is an integral curve of G then $x \equiv \pi \circ \eta$ is a geodesic on M and $\dot{x} = \eta$.
3. The flow of G is $(t, v) \rightarrow \dot{x}_v(t)$.
4. In a chart x^i for M and induced coordinates (x^i, v^i) on TM , G takes the form

$$G(x, v) = v^i \partial_{x^i}|_{(x,v)} - \Gamma_{jk}^i(x) v^j v^k \partial_{v^i}|_{(x,v)}, \tag{2.2}$$

where Γ_{jk}^i are the Christoffel symbols of the Levi–Civita connection in the coordinate system x^i .

5. Using V we can define a vector field $V : TM \rightarrow T(TM)$ given in an induced chart on TM by $V = V^i \partial_{v^i}$ (we will let context dictate whether we consider V as mapping into TM or $T(TM)$). This produces a well-defined smooth vector field on TM that is independent of the choice of charts.
6. We let

$$Y = G + V. \tag{2.3}$$

If τ is an integral curve of Y , then $x = \pi \circ \tau$ satisfies Eq. (2.1) and $\tau = \dot{x}$. Conversely, if x satisfies Eq. (2.1) then \dot{x} is an integral curve of Y .

This last point implies that the equation of interest, Eq. (2.1), defines a smooth dynamical system on the tangent bundle of M and the vector field of this dynamical system is Y .

3. Forced Geodesic Motion on the Frame Bundle

The metric tensor g on M defines a reduction of the structure group of TM to the orthogonal group, $O(\mathbb{R}^n)$ [28], with local trivializations induced by local orthonormal (o.n.) frames on M (i.e., collections of local vector fields that form an o.n. basis at each point of their domain). In turn, this lets one construct the orthogonal frame bundle, $(F_O(M), \pi)$ (we will let context distinguish between the various projections π). By reformulating Eq. (2.1) as a dynamical system

using the orthogonal frame bundle of M , in a similar manner to the procedure outlined in the previous section, we will arrive at equations that are more amenable to being coupled to noise.

Our expanded system will be defined via a vector field on the manifold $N \equiv F_O(M) \times \mathbb{R}^n$.

3.1. Coordinate-Independent Definition

Fix $(u, v) \in N$. We will define a vector $X_{(u,v)} \in T_{(u,v)}N$ as follows. Let $x(t)$ be the solution to

$$\nabla_{\dot{x}} \dot{x} = V(\dot{x}), \quad x(0) = \pi(u), \quad \dot{x}(0) = u(v), \tag{3.1}$$

i.e., the integral curve of Y , defined in Eq. (2.3), starting at $u(v) \in T_{\pi(u)}M$.

Let $U_\alpha(t)$ be the parallel translates of $u(e_\alpha)$ along $x(t)$ (e_α is the standard basis for \mathbb{R}^n), i.e.,

$$\nabla_{\dot{x}} U_\alpha = 0, \quad U_\alpha(0) = u(e_\alpha). \tag{3.2}$$

Parallel transport via the Levi-Civita connection preserves inner products, so $\tau(t)$ defined by $\tau(t)e_\alpha = U_\alpha(t)$ is a smooth section of $F_O(M)$ along $x(t)$. Define the smooth curve in \mathbb{R}^n , $v(t) = \tau(t)^{-1}\dot{x}(t)$.

With these definitions, $\eta(t) = (\tau(t), v(t))$ is a smooth curve in N starting at (u, v) . Define the vector field X by

$$X_{(u,v)} = \dot{\eta}(0). \tag{3.3}$$

3.2. Coordinate Expression

We now derive a formula for X in a coordinate system defined below and thereby prove it is a smooth vector field on N .

Let (U, ϕ) be a coordinate chart on M and E_α be an o.n. frame on U . We will let Roman indices denote quantities in the coordinate frame and Greek indices denote quantities in the local o.n. frame. The connection coefficients in the o.n. frame, $A_{\beta\eta}^\alpha$, are defined by $\nabla_{E_\beta} E_\eta = A_{\beta\eta}^\alpha E_\alpha$. The coordinate frame, ∂_i , and the o.n. frame, E_α , are related by an invertible matrix-valued smooth function Λ_i^α on U ,

$$\partial_i = \Lambda_i^\alpha E_\alpha. \tag{3.4}$$

Let ψ be the local section of $F_O(M)$ induced by E_α , i.e., $\psi(x)v = v^\alpha E_\alpha(x)$. We have the diffeomorphism $\Phi : \pi^{-1}(U) \rightarrow \phi(U) \times O(\mathbb{R}^n)$, $u \rightarrow (\phi(\pi(u)), h)$, where h is uniquely defined by $u = \psi(\pi(u))h$. In turn, this gives a diffeomorphism $\Phi \times id$ on $\pi^{-1}(U) \times \mathbb{R}^n \subset N$.

Lemma 3.1. *The pushforward of the vector field X to $\phi(U) \times O(\mathbb{R}^n) \times \mathbb{R}^n$ by the diffeomorphism $\Phi \times id$ is given by*

$$\begin{aligned} ((\Phi \times id)_* X)|_{(x,h,v)} &= (\Lambda^{-1})_\alpha^j(\pi(u))h_\beta^\alpha v^\beta \partial_j - h_\alpha^\eta A_{\delta\eta}^\beta(\pi(u))h_\xi^\delta v^\xi \partial_{e_\alpha^\beta} \\ &\quad + (h^{-1})_\beta^\alpha V^\beta(u(v))\partial_{v^\alpha} \end{aligned} \tag{3.5}$$

where $u = \Phi^{-1}(\phi^{-1}(x), h)$, V^β are the components of V in the o.n. frame E_β (not the coordinate frame ∂_i), v^α are the standard coordinates on \mathbb{R}^n , and e_α^β are the standard coordinates on $\mathbb{R}^{n \times n}$. In particular, X is a smooth vector field.

Proof. Using

$$\tau(t) = \psi(x(t))h(t) = u(t), \quad v(t) = \tau^{-1}(t)\dot{x}(t) \tag{3.6}$$

we obtain

$$\begin{aligned} v(t) &= h^{-1}(t)\psi^{-1}(x(t))\dot{x}^i(t)\partial_i = \dot{x}^i(t)\Lambda_i^\alpha(x(t))h^{-1}(t)\psi^{-1}(x(t))E_\alpha(x(t)) \\ &= \dot{x}^i(t)\Lambda_i^\alpha(x(t))h^{-1}(t)e_\alpha = \dot{x}^i(t)\Lambda_i^\alpha(x(t))(h^{-1})^\beta_\alpha(t)e_\beta. \end{aligned} \tag{3.7}$$

Solving for $\dot{x}(t)$ we find

$$\dot{x}^j(t) = (\Lambda^{-1})^j_\alpha(x(t))h^\alpha_\beta(t)v^\beta(t). \tag{3.8}$$

This proves that the first term of Eq. (3.5) is correct.

$\nabla_{\dot{x}}U_\alpha = 0$ implies

$$\begin{aligned} 0 &= \nabla_{\dot{x}}\psi(x(t))h(t)e_\alpha = \nabla_{\dot{x}}h^\beta_\alpha(t)E_\beta(x(t)) \\ &= \dot{h}^\beta_\alpha(t)E_\beta(x(t)) + h^\beta_\alpha(t)\dot{x}^k(t)\nabla_{\partial_k}E_\beta \\ &= \dot{h}^\beta_\alpha(t)E_\beta(x(t)) + h^\beta_\alpha(t)\dot{x}^k(t)\Lambda_k^\eta(x(t))\nabla_{E_\eta}E_\beta \\ &= \left(\dot{h}^\beta_\alpha(t) + h^\beta_\alpha(t)\dot{x}^k(t)\Lambda_k^\delta(x(t))A_{\delta\eta}^\beta(x(t)) \right) E_\beta(x(t)). \end{aligned} \tag{3.9}$$

Therefore,

$$\dot{h}^\beta_\alpha(t) = -h^\eta_\alpha(t)A_{\delta\eta}^\beta(x(t))\Lambda_k^\delta(x(t))\dot{x}^k(t). \tag{3.10}$$

Using Eq. (3.8), we get

$$\begin{aligned} \dot{h}^\beta_\alpha(t) &= -h^\eta_\alpha(t)A_{\delta\eta}^\beta(x(t))\Lambda_k^\delta(x(t))((\Lambda^{-1})^k_\alpha(x(t))h^\kappa_\xi(t)v^\xi(t)) \\ &= -h^\eta_\alpha(t)A_{\delta\eta}^\beta(x(t))h^\delta_\xi(t)v^\xi(t). \end{aligned} \tag{3.11}$$

This proves that the second term in Eq. (3.5) is correct.

Differentiating Eq. (3.7) (and dropping the time dependence in our notation) we find

$$\dot{v}^\alpha = -(h^{-1})^\alpha_\beta \dot{h}^\beta_\eta (h^{-1})^\eta_\xi \Lambda_i^\xi \dot{x}^i + (h^{-1})^\alpha_\beta \partial_t \Lambda_i^\beta \dot{x}^i \dot{x}^i + (h^{-1})^\alpha_\beta \Lambda_i^\beta \ddot{x}^i. \tag{3.12}$$

From $\nabla_{\dot{x}}\dot{x} = V$, we obtain $\ddot{x}^i + \Gamma^i_{jk}\dot{x}^j\dot{x}^k = V^i$ where V^i are the components of V in the coordinate frame ∂_i (not to be confused with V^α , the components in the o.n. frame E_α). We need to convert from Γ^i_{jk} to $A_{\beta\eta}^\alpha$,

$$\begin{aligned} \Gamma^i_{jk}\partial_i &= \nabla_{\partial_j}\partial_k = \nabla_{\partial_j}\Lambda_k^\alpha E_\alpha = \partial_j\Lambda_k^\alpha E_\alpha + \Lambda_k^\alpha \nabla_{\partial_j}E_\alpha \\ &= \partial_j\Lambda_k^\alpha E_\alpha + \Lambda_k^\alpha \Lambda_j^\beta \nabla_{E_\beta}E_\alpha \\ &= (\partial_j\Lambda_k^\alpha + \Lambda_k^\eta \Lambda_j^\beta A_{\beta\eta}^\alpha)E_\alpha = (\partial_j\Lambda_k^\alpha + \Lambda_k^\eta \Lambda_j^\beta A_{\beta\eta}^\alpha)(\Lambda^{-1})^i_\alpha \partial_i. \end{aligned} \tag{3.13}$$

Using this we obtain

$$\begin{aligned} \dot{v}^\alpha &= -(h^{-1})^\alpha_\beta \dot{h}^\beta_\eta (h^{-1})^\eta_\delta \Lambda_i^\delta \dot{x}^i + (h^{-1})^\alpha_\beta \partial_t \Lambda_i^\beta \dot{x}^i \dot{x}^i + (h^{-1})^\alpha_\beta \Lambda_i^\beta (V^i - \Gamma^i_{jm}\dot{x}^j\dot{x}^m) \\ &= -(h^{-1})^\alpha_\beta \dot{h}^\beta_\eta (h^{-1})^\eta_\delta \Lambda_i^\delta \dot{x}^i + (h^{-1})^\alpha_\beta \partial_t \Lambda_i^\beta \dot{x}^i \dot{x}^i \\ &\quad + (h^{-1})^\alpha_\beta \Lambda_i^\beta (V^i - (\partial_j\Lambda_m^\eta + \Lambda_m^\delta \Lambda_j^\xi A_{\xi\delta}^\eta)(\Lambda^{-1})^i_\eta \dot{x}^j \dot{x}^m) \\ &= (h^{-1})^\alpha_\beta \Lambda_i^\beta V^i - (h^{-1})^\alpha_\beta \dot{h}^\beta_\eta (h^{-1})^\eta_\delta \Lambda_i^\delta \dot{x}^i + (h^{-1})^\alpha_\beta \partial_t \Lambda_i^\beta \dot{x}^i \dot{x}^i \end{aligned}$$

$$\begin{aligned}
& - (h^{-1})_{\beta}^{\alpha} (\partial_j \Lambda_m^{\beta} + \Lambda_m^{\delta} \Lambda_j^{\xi} A_{\xi\delta}^{\beta}) \dot{x}^j \dot{x}^m \\
= & (h^{-1})_{\beta}^{\alpha} \Lambda_i^{\beta} V^i - (h^{-1})_{\beta}^{\alpha} h_{\eta}^{\beta} (h^{-1})_{\xi}^{\eta} \Lambda_i^{\xi} \dot{x}^i + (h^{-1})_{\beta}^{\alpha} \partial_l \Lambda_i^{\beta} \dot{x}^l \dot{x}^i \\
& - (h^{-1})_{\beta}^{\alpha} \partial_j \Lambda_m^{\beta} \dot{x}^j \dot{x}^m - (h^{-1})_{\beta}^{\alpha} \Lambda_m^{\eta} \Lambda_j^{\xi} A_{\xi\eta}^{\beta} \dot{x}^j \dot{x}^m.
\end{aligned} \tag{3.14}$$

The third and fourth terms cancel. Using Eq. (3.10), the second can be written

$$\begin{aligned}
- (h^{-1})_{\beta}^{\alpha} h_{\eta}^{\beta} (h^{-1})_{\xi}^{\eta} \Lambda_j^{\xi} \dot{x}^j &= (h^{-1})_{\beta}^{\alpha} (h_{\xi}^{\eta} A_{\delta\eta}^{\beta} \Lambda_i^{\delta} \dot{x}^i) (h^{-1})_{\epsilon}^{\xi} \Lambda_j^{\epsilon} \dot{x}^j \\
&= (h^{-1})_{\beta}^{\alpha} A_{\xi\eta}^{\beta} \Lambda_i^{\xi} \Lambda_j^{\eta} \dot{x}^i \dot{x}^j.
\end{aligned} \tag{3.15}$$

Therefore,

$$\begin{aligned}
\dot{v}^{\alpha}(t) &= (h^{-1})_{\beta}^{\alpha} \Lambda_i^{\beta} V^i + (h^{-1})_{\beta}^{\alpha} A_{\xi\eta}^{\beta} \Lambda_i^{\eta} \Lambda_j^{\xi} \dot{x}^i \dot{x}^j - (h^{-1})_{\beta}^{\alpha} A_{\xi\eta}^{\beta} \Lambda_j^{\eta} \Lambda_m^{\xi} \dot{x}^j \dot{x}^m \\
&= (h^{-1})_{\beta}^{\alpha} \Lambda_i^{\beta} V^i = (h^{-1})_{\beta}^{\alpha} V^{\beta}.
\end{aligned} \tag{3.16}$$

Note that we have converted from the components in the coordinate basis, V^i , to the coordinates in the o.n. basis, V^{β} , in the last line. This proves that the final term of Eq. (3.5) is correct.

Note that the equation for $v(t)$ can also be written as

$$\dot{v}(t) = \tau^{-1}(t) V(\tau(t) v(t)). \tag{3.17}$$

□

The cancelation of the Christoffel terms in the equation for \dot{v} is not unexpected. In the absence of forcing $V, x(t)$ is a geodesic and hence its tangent vector is parallel transported along itself. Therefore, v^i , the components of the tangent vector in the parallel-transported frame, U_{α} , must be constants when V vanishes. This is in contrast to the geodesic equation in an arbitrary coordinate system, in which the equation for \ddot{x}^i is non-trivial even in the absence of forcing. This fact simplifies the analysis when we study the small mass limit of the noisy system and is one of the advantages of the orthogonal frame bundle formulation.

Using the above lemma, we can write the equation for the integral curves of X , Eq. (3.3), in coordinates.

Corollary 3.1. *In a coordinate system defined as in Lemma 3.1, an integral curve of X , $(x^i(t), h_{\delta}^{\beta}(t), v^{\alpha}(t))$, satisfies*

$$\dot{x}^j(t) = (\Lambda^{-1})_{\alpha}^j(x(t)) h_{\beta}^{\alpha}(t) v^{\beta}(t), \tag{3.18}$$

$$\dot{h}_{\beta}^{\alpha}(t) = -h_{\beta}^{\eta}(t) A_{\xi\eta}^{\alpha}(x(t)) h_{\delta}^{\xi}(t) v^{\delta}(t), \tag{3.19}$$

$$\dot{v}^{\alpha}(t) = (h^{-1})_{\beta}^{\alpha}(t) V^{\beta}(\dot{x}(t)), \tag{3.20}$$

where from Eq. (3.6), we see that $\dot{x}(t) = v^{\xi}(t) h_{\xi}^{\eta}(t) E_{\eta}(x(t))$. Recall that V^{β} are the components of V in the o.n. basis E_{β} .

In the process of proving Lemma 3.1, we have also characterized the relation between integral curves of X , Eq. (3.3), and integral curves of Y , Eq. (2.3), as expressed by the following corollaries.

Corollary 3.2. *Let $x(t)$ be the solution to*

$$\nabla_{\dot{x}}\dot{x} = V(\dot{x}), \quad x(0) = \pi(u), \quad \dot{x}(0) = u(v), \tag{3.21}$$

i.e., the integral curve of Y , defined in Eq. (2.3), starting at $u(v) \in T_{\pi(u)}M$.

Let $U_\alpha(t)$ be the parallel translates of $u(e_\alpha)$ along $x(t)$ (e_α is the standard basis for \mathbb{R}^n), i.e.,

$$\nabla_{\dot{x}}U_\alpha = 0, \quad U_\alpha(0) = u(e_\alpha). \tag{3.22}$$

Parallel transport preserves inner products, so $\tau(t)$ defined by $\tau(t)e_\alpha = U_\alpha(t)$ is a smooth section of $F_O(M)$ along $x(t)$. Define the smooth curve in \mathbb{R}^n , $v(t) = \tau(t)^{-1}\dot{x}(t)$.

Define the smooth curve in N , $\eta(t) = (\tau(t), v(t))$. This is an integral curve of X starting at (u, v) .

Conversely, uniqueness of integral curves gives us the following.

Corollary 3.3. *Let $(\tau(t), v(t))$ be an integral curve of X starting at (u, v) . Define $x(t) = \pi(\tau(t))$ and $U_\alpha(t) = \tau(t)e_\alpha$. Then $x(t)$ is a solution to*

$$\nabla_{\dot{x}}\dot{x} = V(\dot{x}), \quad x(0) = \pi(u), \quad \dot{x}(0) = u(v), \tag{3.23}$$

the U_α are parallel along $x(t)$, and $v(t) = \tau^{-1}(t)\dot{x}(t)$.

3.3. A Second Coordinate-Independent Formulation

In this section, we introduce a natural set of horizontal vector fields on the orthogonal frame bundle and, using the coordinate expression for X , Eq. (3.5), we show that these vector fields can be used to characterize the dynamical system Eq. (2.1), yielding another coordinate-independent formulation. This will also show the relationship between the Eqs. (3.18)–(3.20) and the equations in [22]. The formulation we give in this section will be utilized for the remainder of the paper, as it has several advantages over our previous characterizations of the system Eq. (2.1). These advantages will be made clear as we progress.

Lemma 3.2. *On an n -dimensional Riemannian manifold, there exists a canonical linear map from \mathbb{R}^n to horizontal vector fields on $F_O(M)$ defined as follows (see [8, 22]).*

For each $v \in \mathbb{R}^n$ and $u \in F_O(M)$, define $H_v(u) \in T_uF_O(M)$ by $H_v(u) = (u(v))^h$, i.e., the horizontal lift of $u(v) \in T_{\pi(u)}M$ to $T_uF_O(M)$.

This is a smooth horizontal vector field on $F_O(M)$. Pushing forward to $U \times O(\mathbb{R}^n)$ via a local trivialization (U, Φ) of $F_O(M)$ about u with corresponding o.n. frame E_α , as in Sect. 3.1, they have the form

$$H_v(u) = v^\alpha h_\alpha^\beta E_\beta(\pi(u)) - v^\alpha h_\alpha^\beta h_\xi^\eta A_{\beta\eta}^\delta(\pi(u)) \partial_{e_\xi}^\delta, \tag{3.24}$$

where $\Phi(u) = (\pi(u), h)$, $\nabla_{E_\alpha}E_\beta = A_{\alpha\beta}^\eta E_\eta$, v^α are the components of v in the standard basis for \mathbb{R}^n , and e_α^β are the standard coordinates on $\mathbb{R}^{n \times n}$. Note that the second term defines a vector field on $\mathbb{R}^{n \times n}$, but it is in fact tangent to $O(\mathbb{R}^n)$. Our expression Eq. (3.24) differs slightly from the one found in [8], as we have written it in an o.n. frame rather than a coordinate frame.

If e_α is the standard basis for \mathbb{R}^n we will let $H_\alpha \equiv H_{e_\alpha}$. Therefore, $H_v = v^\alpha H_\alpha$ for any $v \in \mathbb{R}^n$, where we employ the summation convention.

Under right multiplication by $g \in O(\mathbb{R}^n)$, these vector fields satisfy

$$(R_g)_*(H_v(u)) = H_{g^{-1}v}(ug). \tag{3.25}$$

Remark 1. The implied summations in $v^\alpha h_\alpha^\beta, v^\alpha H_\alpha$, etc., are summations over components in the standard basis for \mathbb{R}^n . The α 's here are not tensor indices on M, TM , or $F_O(M)$ and do not transform under change of coordinates or frame. This is in contrast with the index β in h_α^β , which does transform under a change of the o.n. frame E_β . We will occasionally revisit this point going forward for emphasis.

The horizontal vector fields Eq. (3.24) can be used to relate geodesic motion and parallel transport on M to a flow on the frame bundle.

Lemma 3.3. *Let $u \in F_O(M)$ and $v \in \mathbb{R}^n$. Let τ be the integral curve of H_v starting at u . Then $x \equiv \pi \circ \tau$ is the geodesic starting at $\pi(u)$ with initial velocity $u(v)$ and for any $w \in \mathbb{R}^n, \tau(t)w$ is parallel transported along $x(t)$.*

Proof. τ is a horizontal curve in $F_O(M)$ iff $\tau(w)$ is horizontal in TM for any $w \in \mathbb{R}^n$. In a vector bundle, horizontal and parallel transported are synonymous. Hence, $\tau(t)w$ is parallel transported along $x = \pi \circ \tau$. Therefore, to prove $x(t)$ is the claimed geodesic it suffices to show $\dot{x} = \tau(v)$.

In a local trivialization $\Phi, \Phi(\tau(t)) = (x(t), h(t))$. Hence, using Eq. (3.24), we have

$$\dot{x}(t) = v^\alpha h_\alpha^\beta(t) E_\beta(\pi(\tau(t))) = \tau(t)v. \tag{3.26}$$

□

Uniqueness of geodesics, parallel transport, and integral curves then gives the following.

Lemma 3.4. *Let $x \in M, u$ be a frame at x , and $v \in \mathbb{R}^n$. Let $x(t)$ be the geodesic starting at x with initial velocity $u(v)$. Let e_α be the standard basis for \mathbb{R}^n and U_α be the parallel translates of $u(e_\alpha)$ along $x(t)$. Let $\tau(t)$ be the corresponding section of $F_O(M)$, i.e., $\tau(t)e_\alpha = U_\alpha(t)$. Then τ is the integral curve of H_v starting at u .*

We can also use the H 's to lift vector fields from M to the frame bundle.

Lemma 3.5. *Let b be a smooth vector field on M and b^h be the horizontal lift of b to $F_O(M)$. Recall that this is a smooth vector field on $F_O(M)$. We have*

$$b^h(u) = H_{u^{-1}b(\pi(u))}(u). \tag{3.27}$$

If R_g denotes right multiplication by $g \in O(\mathbb{R}^n)$ then $(R_g)_*b^h = b^h$.

Proof. To prove the first assertion, by the definition of H ,

$$H_{u^{-1}b(\pi(u))}(u) = (u(u^{-1}b(\pi(u))))^h = (b(\pi(u)))^h = b^h(u). \tag{3.28}$$

As for the second,

$$\pi_*(((R_g)_*b^h)(u)) = (\pi \circ R_g)_*b^h(ug^{-1}) = b(\pi(ug^{-1})) = b(\pi(u)). \tag{3.29}$$

$(R_g)_*$ preserves the horizontal subspaces, hence $(R_g)_*b^h$ is the horizontal lift of b . □

Lemma 3.6. *Let b be a smooth vector field on M . If τ is an integral curve of b^h starting at u then $x \equiv \pi \circ \tau$ is an integral curve of b starting at $\pi(u)$ and for any $v \in \mathbb{R}^n, \tau(t)v$ is the parallel translate of $u(v)$ along $x(t)$.*

Conversely, if $x(t)$ is an integral curve of b starting at $\pi(u)$ and $U_\alpha(t)$ are the parallel translates of $u(e_\alpha)$ along $x(t)$ then $\tau(t)$ defined by $\tau(t)e_\alpha = U_\alpha(t)$ is the integral curve of b^h starting at u .

Proof. Suppose τ is an integral curve of b^h starting at u . Then

$$\dot{x} = \pi_*\dot{\tau} = \pi_*b^h(\tau) = b(x). \tag{3.30}$$

So $x(t)$ is an integral curve of b . τ has horizontal tangent vector for all t , hence $\tau(t)v$ is parallel in TM .

Conversely, if $x(t)$ is an integral curve of b starting at $\pi(u)$ and $U_\alpha(t)$ are the parallel translates of $u(e_\alpha)$ then $\tau(t)$ defined by $\tau(t)e_\alpha = U_\alpha(t)$ is a smooth horizontal curve in $F_O(M)$ and $\tau(t_0) = u$. We have

$$\pi_*\dot{\tau} = \dot{x} = b(x). \tag{3.31}$$

$\dot{\tau}$ is horizontal, so

$$\dot{\tau} = (b(x))^h = b^h(\tau). \tag{3.32}$$

□

The prior lemmas show that geodesic motion, parallel transport, and flows on M can all be related to flows on $F_O(M)$. Therefore, it should not come as a surprise that the vector field X , Eq. (3.3), whose integral curves characterize the trajectories of our deterministic system, can be written in terms of the H_v 's and the forcing, V .

Proposition 3.1. *Let $N = F_O(M) \times \mathbb{R}^n$ and $(u, v) \in N$. Then $X_{(u,v)}$, defined by Eq. (3.3), is given by*

$$X_{(u,v)} = (H_v(u), u^{-1}V(u(v))) \tag{3.33}$$

where we have identified $T\mathbb{R}^n$ with \mathbb{R}^n .

Proof. In a local trivialization induced by an o.n. frame E_α , Eq. (3.5) implies that

$$X|_{(x,h,v)} = h^\alpha_\beta v^\beta E_\alpha(x) - h^\eta_\alpha A^\beta_{\delta\eta}(x) h^\delta_\xi v^\xi \partial_{e_\alpha} + (h^{-1})^\alpha_\beta V^\beta(u(v)) \partial_{v^\alpha}. \tag{3.34}$$

The proposition then follows from Eq. (3.24). □

The geometric significance of the H_v 's will make Eq. (3.33) simpler to work with than our initial definition of the vector field X , Eq. (3.3).

Proposition 3.1 implies that the deterministic dynamics of the system of interest, Eq. (2.1), lifted to $N = F_O(M) \times \mathbb{R}^n$, are given by

$$\dot{u} = H_v(u), \quad \dot{v} = u^{-1}V(u(v)), \quad (u(t_0), v(t_0)) = (u_0, v_0). \tag{3.35}$$

We want to emphasize that v is defined in terms of the dynamical frame u , and not in reference to any choice of coordinates on M or $F_O(M)$. In other

words, the components v^α of v in the standard basis for \mathbb{R}^n are the components of the particle's velocity *in its own parallel-transported frame*. They are not tied to a particular coordinate system on M or $F_O(M)$ and do not transform under coordinate changes on either space.

4. Randomly Perturbed Geodesic Flow With Forcing

In this section, we will show how we couple noise to the system Eq. (3.35) to obtain a stochastic differential equation on N .

4.1. Stochastic Differential Equations on Manifolds

First, we recall the definition and some basic properties of semimartingales and stochastic differential equations on manifolds. The definition and lemmas in this section are adapted from [8], but we repeat them here for completeness. The general theory outlined in this section does not require a Riemannian metric on M .

Definition 1. Let M be an n -dimensional smooth manifold, $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$ be a filtered probability space satisfying the usual conditions [1], and X_t be a continuous adapted M -valued process. X is called an M -valued continuous semimartingale if $f \circ X_t$ is an \mathbb{R} -valued semimartingale for all $f \in C^\infty(M)$. We will only deal with continuous semimartingales, so we drop the adjective continuous from now on.

Note that, by Itô's formula, if $M = \mathbb{R}^n$ then this agrees with the usual definition.

Definition 2. Let V be a k -dimensional vector space and Z_t be a V -valued semimartingale, called the driving process. Let M be a smooth manifold, X_t be an M -valued semimartingale, and σ be a smooth section of $TM \otimes V^*$. We say that X_t is a solution to the SDE

$$X_t = X_{t_0} + \int_{t_0}^t \sigma(X_s) \circ dZ_s \tag{4.1}$$

if

$$f(X_t) = f(X_{t_0}) + \int_{t_0}^t \sigma(X_s)[f] \circ dZ_s \tag{4.2}$$

P -a.s. for all $f \in C^\infty(M)$, where $\int \dots \circ dZ_s$ denotes the stochastic integral in the Stratonovich sense. We use the notation $Y[f]$ to denote the smooth function one obtains by operating with some vector field, Y , on a smooth function, f , and in the stochastic integral we contract over the V^* and V factors in $\sigma[f]$ and Z , respectively. We will equivalently write the SDE (4.1) in differential notation

$$dX_t = \sigma(X_s) \circ dZ_s. \tag{4.3}$$

Note that when M is a finite dimensional vector space, this definition agrees with the usual one (in the Stratonovich sense). Using a basis for V and the dual basis for V^* to write the contraction in Eq. (4.2) as a sum over components in these bases, we arrive at a formula analogous to the definition in [8] (page 21). However, we find it useful to use the above formulation in terms of a vector space and its dual to justify use of the summation convention over contracted indices.

The Stratonovich integral is used in Eq. (4.2) to make the definition diffeomorphism invariant, as captured by the following Stratonovich calculus variant of the Itô change-of-variables formula (see [8, pp. 20–21]).

Lemma 4.1. *Let X_t be an M -valued semimartingale that satisfies the SDE*

$$X_t = X_{t_0} + \int_{t_0}^t \sigma(X_s) \circ dZ_s, \tag{4.4}$$

N be another smooth manifold, and $\Phi : M \rightarrow N$ be a diffeomorphism. Then $\tilde{X} \equiv \Phi \circ X$ is an N -valued semimartingale and satisfies the SDE

$$\tilde{X}_t = \tilde{X}_{t_0} + \int_{t_0}^t (\Phi_* \sigma)(\tilde{X}_s) \circ dZ_s \tag{4.5}$$

where Φ_* denotes the pushforward.

Definition 4.2 can be restated in terms of the Itô integral as follows, similar to p.23 of [8].

Lemma 4.2. *X_t is a solution to the SDE*

$$X_t = X_{t_0} + \int_{t_0}^t \sigma(X_s) \circ dZ_s \tag{4.6}$$

iff

$$f(X_t) = f(X_{t_0}) + \int_{t_0}^t \sigma(X_s)[f]dZ_s + \frac{1}{2} \int_{t_0}^t \sigma_\alpha(X_s)[\sigma_\beta[f]]d[Z^\alpha, Z^\beta]_s \tag{4.7}$$

for all $f \in C^\infty(M)$ where the summation convention is employed and the sum is over the components in any pair of dual bases for V and V^ . This is another manifestation of the Itô formula for the stochastic differential of the composition of a smooth function with a semimartingale.*

4.2. Coupling to Noise

For the remainder of this paper, we will assume M is compact, connected, and without boundary. Note that this also implies $F_O(M)$ is compact and without boundary. In this section, we describe the coupling of the dynamical system Eq. (3.35) to noise, and hence we must also assume M is equipped with a Riemannian metric.

Let W be an \mathbb{R}^k -valued Wiener process and $\sigma : F_O(M) \rightarrow \mathbb{R}^{n \times k}$ be smooth. We are interested in the following SDE for $(u, v) \in N = F_O(M) \times \mathbb{R}^n$,

$$u_t = u_0 + \int_{t_0}^t H_{v_s}(u_s)ds, \tag{4.8}$$

$$v_t = v_0 + \frac{1}{m} \int_{t_0}^t u_s^{-1} V(u_s v_s) ds + \frac{1}{m} \int_{t_0}^t \sigma(u_s) \circ dW_s. \tag{4.9}$$

Note that we have replaced V in Eq. (3.35) with $\frac{1}{m}V$ (where now V is independent of m), making the dependence on particle mass, m , explicit.

To connect with Definition 2, one must view $H_v(u)$, $\frac{1}{m}u^{-1}V(uv)$, and $\frac{1}{m}\sigma(u)$ as sections of $TN \otimes (\mathbb{R}^{k+1})^*$ (identifying $T_{(u,v)}N$ with $T_uF_O(M) \oplus \mathbb{R}^n$), and use the driving \mathbb{R}^{k+1} -valued semimartingale $Z_t = (t, W_t)$. Alternatively, one could view the above objects as $k + 1$ vector fields on N and include sums over indices, as done in [8], but for economy of notation, we wish to avoid employing indices and explicit summations when possible.

Because the Wiener process only couples to the equation for v , which is a process with values in the second factor of the product space $N = F_O(M) \times \mathbb{R}^n$, a solution of the SDE (4.8)–(4.9) on the manifold N in the sense of Eq. (4.2) is equivalent to the existence of an N -valued semimartingale, (u, v) , such that the first component is pathwise C^1 and pathwise satisfies the ODE

$$\dot{u}_t = H_{v_t}(u_t), \quad u(t_0) = u_0 \tag{4.10}$$

and the second component satisfies the SDE on \mathbb{R}^n

$$v_t = v_0 + \frac{1}{m} \int_{t_0}^t u_s^{-1} V(u_s v_s) ds + \frac{1}{m} \int_{t_0}^t \sigma(u_s) dW_s. \tag{4.11}$$

Note that u has locally bounded variation, so the choice of stochastic integral in the second equation is not significant. We use the Itô notation here. We emphasize that while the machinery of Sect. 4.1 is not needed to formulate the above system, it will be required when we pass to the limit $m \rightarrow 0$.

For the remainder of the paper, we will make the following assumption.

Assumption 1. We will assume that the deterministic vector field V is the sum of a position-dependent force term and a position-dependent linear drag term

$$V(w) = F(x) - \gamma(x)w, \quad w \in T_xM, \quad x = \pi(w) \in M, \tag{4.12}$$

where F is a smooth vector field on M and γ is a smooth $\binom{1}{1}$ tensor field on M . We will *not* assume that the force field F comes from a potential.

As stated in Sect. 1.1, we will also assume that the symmetric part of γ , $\gamma^s = \frac{1}{2}(\gamma + \gamma^T)$, has eigenvalues bounded below by a constant $\gamma_1 > 0$ on all of M . We again emphasize that this coercivity assumption will be crucial for the momentum decay estimates of Sect. 5.

In the following it will be useful to denote $u^{-1}F(\pi(u))$ by $F(u)$ and $u^{-1}\gamma(\pi(u))u$ by $\gamma(u)$, letting the context distinguish between the different notations. These are smooth \mathbb{R}^n - and $\mathbb{R}^{n \times n}$ -valued functions on $F_O(M)$, respectively. With these definitions, the SDE (4.8)–(4.9) becomes

$$u_t = u_0 + \int_{t_0}^t H_{v_s}(u_s) ds, \tag{4.13}$$

$$v_t = v_0 + \frac{1}{m} \int_{t_0}^t [F(u_s) - \gamma(u_s)v_s] ds + \frac{1}{m} \int_{t_0}^t \sigma(u_s) dW_s. \tag{4.14}$$

Given k vector fields, $\sigma_\alpha(x)$, on M , these induce corresponding noise coefficients on the frame bundle, $\sigma(u)$, given by

$$\sigma(u)e_\alpha = u^{-1}\sigma_\alpha(\pi(u)). \tag{4.15}$$

Additionally, one is often interested in the case where $k = n$ and $\sigma(u)$ comes from a $\binom{1}{1}$ -tensor field on a M , denoted $\sigma(x)$, in the same manner as $\gamma(u)$, i.e.,

$$\sigma(u) = u^{-1}\sigma(\pi(u))u. \tag{4.16}$$

For most of this work, we keep the discussion general and deal only with $\sigma(u)$.

The following lemmas will be useful.

Lemma 4.3. *Let γ^s denote the symmetric part of γ . Then*

$$\gamma^s(u) = u^{-1}\gamma^s(\pi(u))u \tag{4.17}$$

Proof. We are done if we can show

$$\gamma^T(u) = u^{-1}\gamma^T(\pi(u))u. \tag{4.18}$$

Letting \cdot be the Euclidean inner product on \mathbb{R}^n , for $x, y \in \mathbb{R}^n$ we have

$$\begin{aligned} y \cdot \gamma^T(u)x &= (\gamma(u)y) \cdot x = (u^{-1}\gamma(\pi(u))uy) \cdot x = g(\gamma(\pi(u))uy, ux) \\ &= g(uy, \gamma^T(\pi(u))ux) = y \cdot (u^{-1}\gamma^T(\pi(u))ux). \end{aligned} \tag{4.19}$$

This holds for all x, y and so the proof is complete. □

Corollary 4.1. *The eigenvalues of $\gamma^s(u)$ and $\gamma^s(\pi(u))$ are the same. In particular, by Assumption 1, the eigenvalues of $\gamma^s(u)$ are also bounded below by $\gamma_1 > 0$ for all $u \in F_O(M)$.*

This also implies that the real parts of the eigenvalues of $\gamma(u)$ are bounded below by γ_1 for all $u \in F_O(M)$. In addition,

$$\|e^{-t\gamma(u)}\| \leq e^{-\gamma_1 t}, \quad \|e^{-t\gamma(u)^T}\| \leq e^{-\gamma_1 t}, \tag{4.20}$$

for any $u \in F_O(M)$ and any $t \geq 0$ (see, for example, p. 86 of [29]).

Lemma 4.4. *For each $(u_0, v_0) \in N$ there exists a unique globally defined solution $(u_t, v_t), t \in [0, \infty)$ to the SDE (4.13)–(4.14) that pathwise satisfies the initial conditions. It can be chosen so that pathwise, $t \mapsto u_t$ is C^1 and satisfies the ODE (4.10). We emphasize that the global-in-time existence relies on the compactness of M .*

Proof. The diffusion term for the SDE is independent of v and the drift is an affine function of v , so compactness of $F_O(M)$ implies that the drift and diffusion are linearly bounded in v , uniformly in u . Therefore, by embedding $F_O(M)$ compactly in some \mathbb{R}^l , one can use the results on global existence and uniqueness of solutions to a vector-valued SDE with linearly bounded coefficients (see for example [1, Theorem 5.2.9]) to prove existence of a unique globally defined solution to the SDE that pathwise satisfies the initial conditions. One can modify the result on a measure zero set to ensure that the u component is also a C^1 -function of t and satisfies the ODE (4.10) everywhere, not just a.s. □

Often one is only interested in the evolution of the position, $x_t = \pi(u_t)$, and velocity, \dot{x}_t , degrees of freedom. The SDE (4.13) implies that $\dot{x}_t = u_t v_t$ and, pathwise, $u(t)$ is horizontal. In particular, for any $w \in \mathbb{R}^n$, $u(t)w$ is parallel transported along $x(t)$, the same as for the deterministic system. The following lemma captures the dependence of the solution on the choice of an initial frame in the case where σ is given by Eq. (4.16).

Lemma 4.5. *Let $h \in O(\mathbb{R}^n)$ and (u_t, v_t) be the solution to Eqs. (4.13)–(4.14) corresponding to the initial condition (u_0, v_0) . Suppose $\sigma(u)$ is obtained from $\sigma(x)$ as in Eq. (4.16). Then*

$$(\tilde{u}_t, \tilde{v}_t) \equiv (u_t h, h^{-1} v_t) \tag{4.21}$$

is the solution to Eqs. (4.13)–(4.14) with the initial condition $(u_0 h, h^{-1} v_0)$ and the Wiener process W_t replaced by the Wiener process $\tilde{W}_t = h^{-1} W_t$.

Proof. $(\tilde{u}_t, \tilde{v}_t)$ is a semimartingale starting at $(u_0 h, h^{-1} v_0)$. The map

$$\Phi(u, v) = (u h, h^{-1} v) \tag{4.22}$$

is a diffeomorphism of N and, therefore, Lemma 4.1 implies

$$\begin{aligned} d\tilde{u}_t &= (R_h)_*(H_{v(t)}(u_t))dt, \\ d\tilde{v}_t &= \frac{1}{m}(L_{h^{-1}})_*(F(u_t) - \gamma(u_t)v_t)dt + \frac{1}{m}(L_{h^{-1}})_*\sigma(u_s)dW_s \end{aligned} \tag{4.23}$$

where R and L denote right and left multiplication, respectively. Using the definitions of $F(u)$, $\gamma(u)$ and $\sigma(u)$ this simplifies to

$$d\tilde{u}_t = H_{\tilde{v}_t}(\tilde{u}_t)dt, \quad d\tilde{v}_t = \frac{1}{m}(F(\tilde{u}_t) - \gamma(\tilde{u}_t)\tilde{v}_t)dt + \frac{1}{m}\sigma(\tilde{u}_t)d\tilde{W}_t. \tag{4.24}$$

□

5. Rate of Decay of the Momentum

We now begin our investigation of the properties of the solutions of the SDE (4.13)–(4.14) in the small mass limit by proving that the momentum process, $p_t = m v_t$, converges to zero in several senses as $m \rightarrow 0$. To this end, we will introduce a superscript to the solutions, (u_t^m, v_t^m) , of Eqs. (4.13)–(4.14) to denote the corresponding value of the mass. The non-random initial conditions, u_0, v_0 , will be fixed independently of m .

More specifically, the momentum process will be shown to converge to zero at a rate dependent on powers of m . This convergence is shown with respect to the uniform L^p -metric on continuous paths (Proposition 5.1), L^p metric (Proposition 5.2), and as a stochastic integral with respect to the momentum (Proposition 5.3). To prove these propositions, the equation for v_t^m , Eq. (4.14), is solved in terms of u_t^m . Estimates are made on the Lebesgue integrals much like in the ordinary differential equation case. The stochastic integral term is rewritten to mirror the ODE case as closely as possible and then broken into small intervals which can be controlled using the Burkholder–Davis–Gundy inequalities.

First we give some useful lemmas.

5.1. Some Lemmas

Lemma 5.1. *Let $X_t = X_0 + M_t + A_t$ be a continuous \mathbb{R}^k -valued semimartingale on $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$ with local martingale component M_t and locally bounded variation component A_t . Let $V \in L^1_{loc}(A) \cap L^2_{loc}(M)$ be $\mathbb{R}^{n \times k}$ -valued and let $B(t)$ be a continuous $\mathbb{R}^{n \times n}$ -valued adapted process. Let $\Phi(t)$ be the adapted $\mathbb{R}^{n \times n}$ -valued C^1 process that pathwise solves the initial value problem (IVP)*

$$\dot{\Phi}(t) = B(t)\Phi(t), \quad \Phi(0) = I. \tag{5.1}$$

Then we have the P -a.s. equalities

$$\Phi(t) \int_0^t \Phi^{-1}(s)V_s dX_s = \int_0^t V_s dX_s + \Phi(t) \int_0^t \Phi^{-1}(s)B(s) \left(\int_0^s V_r dX_r \right) ds \tag{5.2}$$

$$= \Phi(t) \int_0^t V_s dX_s - \Phi(t) \int_0^t \Phi^{-1}(s)B(s) \left(\int_s^t V_r dX_r \right) ds \tag{5.3}$$

for all t .

If the eigenvalues of the symmetric part of $B, B^s = \frac{1}{2}(B + B^T)$, are bounded above by $-\alpha$ for some $\alpha > 0$ then for every $T \geq \delta > 0$ we have the P -a.s. bound

$$\begin{aligned} \sup_{t \in [0, T]} \left\| \Phi(t) \int_0^t \Phi^{-1}(s)V_s dX_s \right\| &\leq \left(1 + \frac{4}{\alpha} \sup_{s \in [0, T]} \|B(s)\| \right) \\ &\times \left(e^{-\alpha\delta} \sup_{t \in [0, T]} \left\| \int_0^t V_r dX_r \right\| + \max_{k=0, \dots, N-1} \sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t V_r dX_r \right\| \right) \end{aligned} \tag{5.4}$$

where $N = \max\{k \in \mathbb{Z} : k\delta < T\}$. Here and in the following, we use the ℓ^2 norm on every \mathbb{R}^k .

Proof. Using integration by parts, together with the fact that Φ is a process of locally bounded variation and $\dot{\Phi}(t) = B(t)\Phi(t)$, we obtain the P -a.s. equality

$$\Phi(t) \int_0^t \Phi^{-1}(s)V_s dX_s = \int_0^t V_s dX_s + \int_0^t B(s)\Phi(s) \int_0^s \Phi^{-1}(r)V_r dX_r ds \tag{5.5}$$

for all t .

Fix an $\omega \in \Omega$ for which the above equality holds and consider the resulting continuous functions $r(t) = \int_0^t V_s dX_s$ and $y(t) = \Phi(t) \int_0^t \Phi^{-1}(s)V_s dX_s$. Eq. (5.5) implies that these satisfy the integral equation

$$y(t) = r(t) + \int_0^t B(s)y(s)ds, \quad y(0) = 0. \tag{5.6}$$

The unique solution to this equation is [30]

$$y(t) = r(t) + \Phi(t) \int_0^t \Phi^{-1}(s)B(s)r(s)ds. \tag{5.7}$$

This proves the first equality in Eq. (5.2). For the second, we compute

$$\begin{aligned}
 & \Phi(t) \int_0^t V_s dX_s - \Phi(t) \int_0^t \Phi^{-1}(s)B(s) \left(\int_s^t V_r dX_r \right) ds \\
 &= \Phi(t) \left(\int_0^t V_s dX_s - \int_0^t \Phi^{-1}(s)B(s) \left(\int_0^t V_r dX_r - \int_0^s V_r dX_r \right) ds \right) \\
 &= \Phi(t) \int_0^t \Phi^{-1}(s)B(s) \left(\int_0^s V_r dX_r \right) ds \\
 &\quad + \Phi(t) \left(I - \int_0^t \Phi^{-1}(s)B(s) ds \right) \int_0^t V_r dX_r \\
 &= \Phi(t) \int_0^t \Phi^{-1}(s)B(s) \left(\int_0^s V_r dX_r \right) ds \\
 &\quad + \Phi(t) \left(I + \int_0^t \frac{d}{ds} \Phi^{-1}(s) ds \right) \int_0^t V_r dX_r \\
 &= \Phi(t) \int_0^t \Phi^{-1}(s)B(s) \left(\int_0^s V_r dX_r \right) ds + \int_0^t V_r dX_r,
 \end{aligned} \tag{5.8}$$

where we have used the formula

$$\frac{d}{ds} \Phi^{-1}(s) = -\Phi^{-1}(s)\dot{\Phi}(s)\Phi^{-1}(s). \tag{5.9}$$

To obtain the bound Eq. (5.4) we start from Eq. (5.3) and take the norm to find

$$\begin{aligned}
 \left\| \Phi(t) \int_0^t \Phi^{-1}(s)V_s dX_s \right\| &\leq \|\Phi(t)\| \left\| \int_0^t V_s dX_s \right\| \\
 &\quad + \int_0^t \|\Phi(t)\Phi^{-1}(s)\| \|B(s)\| \left\| \int_s^t V_r dX_r \right\| ds.
 \end{aligned} \tag{5.10}$$

For $t \geq s$, the fundamental solution $\Phi(t)\Phi^{-1}(s)$ satisfies the bound

$$\|\Phi(t)\Phi^{-1}(s)\| \leq e^{\int_s^t \lambda_{\max}(r) dr} \tag{5.11}$$

where $\lambda_{\max}(r)$ is the largest eigenvalue of $B^s(r)$ (see, for example, p. 86 of [29]). Therefore, assuming $\lambda_{\max} \leq -\alpha < 0$ gives

$$\begin{aligned}
 \left\| \Phi(t) \int_0^t \Phi^{-1}(s)V_s dX_s \right\| &\leq e^{-\alpha t} \left\| \int_0^t V_s dX_s \right\| \\
 &\quad + \sup_{s \in [0,t]} \|B(s)\| \int_0^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds.
 \end{aligned} \tag{5.12}$$

For any $T \geq \delta > 0$ we have the P -a.s. bounds

$$\sup_{t \in [0,T]} e^{-\alpha t} \left\| \int_0^t V_s dX_s \right\| \leq \sup_{t \in [0,\delta]} \left\| \int_0^t V_s dX_s \right\| + e^{-\alpha \delta} \sup_{t \in [\delta,T]} \left\| \int_0^t V_s dX_s \right\| \tag{5.13}$$

and

$$\begin{aligned} & \sup_{t \in [0, T]} \left(\sup_{s \in [0, t]} \|B(s)\| \int_0^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \right) \\ & \leq \sup_{s \in [0, T]} \|B(s)\| \left(\sup_{t \in [0, \delta]} \int_0^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \right. \\ & \quad \left. + \sup_{t \in [\delta, T]} \int_0^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \right). \end{aligned} \tag{5.14}$$

The first term can be bounded as follows:

$$\begin{aligned} & \sup_{t \in [0, \delta]} \int_0^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \\ & = \sup_{t \in [0, \delta]} \int_0^t e^{-\alpha(t-s)} \left\| \int_0^t V_r dX_r - \int_0^s V_r dX_r \right\| ds \\ & \leq \sup_{t \in [0, \delta]} \int_0^t e^{-\alpha(t-s)} 2 \sup_{0 \leq r \leq \delta} \left\| \int_0^r V_r dX_r \right\| ds \leq \frac{2}{\alpha} \sup_{0 \leq t \leq \delta} \left\| \int_0^t V_r dX_r \right\|. \end{aligned} \tag{5.15}$$

In the second term, we can split the integral to obtain

$$\begin{aligned} & \sup_{t \in [\delta, T]} \int_0^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \\ & = \sup_{t \in [\delta, T]} \left(\int_0^{t-\delta} e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds + \int_{t-\delta}^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \right) \\ & \leq \frac{2}{\alpha} e^{-\alpha\delta} \sup_{t \in [0, T]} \left\| \int_0^t V_r dX_r \right\| + \sup_{t \in [\delta, T]} \int_{t-\delta}^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds. \end{aligned} \tag{5.16}$$

Let $N = \max\{k \in \mathbb{Z} : k\delta < T\}$. Then P -a.s.

$$\begin{aligned} & \sup_{t \in [\delta, T]} \int_{t-\delta}^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \\ & \leq \max_{k=0, \dots, N-1} \sup_{t \in [(k+1)\delta, (k+2)\delta]} \int_{k\delta}^t e^{-\alpha(t-s)} \left\| \int_s^t V_r dX_r \right\| ds \\ & = \max_{k=0, \dots, N-1} \sup_{t \in [(k+1)\delta, (k+2)\delta]} \int_{k\delta}^t e^{-\alpha(t-s)} \left\| \int_{k\delta}^t V_r dX_r - \int_{k\delta}^s V_r dX_r \right\| ds \\ & \leq \frac{2}{\alpha} \max_{k=0, \dots, N-1} \sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t V_r dX_r \right\|. \end{aligned} \tag{5.17}$$

Combining Eqs. (5.15) and (5.17) and using the inequality

$$\sup_{t \in [0, \delta]} \left\| \int_0^t V_s dX_s \right\| \leq \max_{k=0, \dots, N-1} \sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t V_r dX_r \right\| \tag{5.18}$$

gives the P -a.s. bound

$$\begin{aligned}
 & \sup_{t \in [0, T]} \left\| \Phi(t) \int_0^t \Phi^{-1}(s) V_s dX_s \right\| \\
 & \leq \sup_{t \in [0, \delta]} \left\| \int_0^t V_s dX_s \right\| + e^{-\alpha \delta} \sup_{t \in [\delta, T]} \left\| \int_0^t V_s dX_s \right\| \\
 & \quad + \frac{2}{\alpha} \sup_{s \in [0, T]} \|B(s)\| \left(\sup_{t \in [0, \delta]} \left\| \int_0^t V_r dX_r \right\| + e^{-\alpha \delta} \sup_{t \in [0, T]} \left\| \int_0^t V_r dX_r \right\| \right) \\
 & \quad + \max_{k=0, \dots, N-1} \sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t V_r dX_r \right\| \\
 & \leq \left(1 + \frac{4}{\alpha} \sup_{s \in [0, T]} \|B(s)\| \right) \left(e^{-\alpha \delta} \sup_{t \in [0, T]} \left\| \int_0^t V_r dX_r \right\| \right. \\
 & \quad \left. + \max_{k=0, \dots, N-1} \sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t V_r dX_r \right\| \right) \tag{5.19}
 \end{aligned}$$

as claimed. □

It will also be useful to recall the following (see [1]).

Lemma 5.2. *If $M \in \mathcal{M}^{c,loc}$, $V \in L^2_{loc}(M)$, and $E[\int_{t_0}^t V^2 d[M]_s] < \infty$ for all t then $\int_{t_0}^t V_s dM_s$ is a martingale.*

5.2. Limit of the Momentum Process

In this section, we show three propositions about convergence of the momentum process $p_t^m = mv_t^m$ to zero as $m \rightarrow 0$.

Proposition 5.1. *For any $p > 0, T > 0$, and $0 < \beta < p/2$ we have*

$$E \left[\sup_{t \in [0, T]} \|p_t^m\|^p \right] = O(m^\beta) \quad \text{as } m \rightarrow 0. \tag{5.20}$$

Proof. The strategy here is to first rewrite the equation for p_t^m so that the stochastic integral term has the same form as the left-hand side of Eq. (5.2). Using the bound Eq. (5.4), we will then be able to show that both terms decay as $m \rightarrow 0$. The first term will decay exponentially, and the second term will decay because the stochastic integrals will be taken over “small” time intervals.

The momentum solves the SDE

$$dp_t^m = \left(F(u_t^m) - \frac{1}{m} \gamma(u_t^m) p_t^m \right) dt + \sigma(u_t^m) dW_t. \tag{5.21}$$

This is a linear SDE on \mathbb{R}^n where $F(u_t^m)$, $-\frac{1}{m} \gamma(u_t^m)$, and $\sigma(u_t^m)$ are pathwise continuous adapted vector- or matrix-valued processes, and so its unique solution can be written in terms of u_s^m

$$p_t^m = \Phi(t) \left(p_0^m + \int_0^t \Phi^{-1}(s) F(u_s^m) ds + \int_0^t \Phi^{-1}(s) \sigma(u_s^m) dW_s \right) \tag{5.22}$$

where $\Phi(t)$ is the adapted C^1 process that pathwise solves the IVP

$$\dot{\Phi}(t) = -\frac{1}{m}\gamma(u_t^m)\Phi(t), \quad \Phi(0) = I. \tag{5.23}$$

This technique of utilizing the explicit solution of a linear SDE to obtain estimates is used in [31], where the case of constant, scalar drag on flat Euclidean space is studied.

By Assumption 1, the symmetric part of $-\frac{1}{m}\gamma(u)$ has eigenvalues bounded above by $-\gamma_1/m < 0$ with the bound uniform in u . Therefore, using Eq. (5.11), for $s \leq t$,

$$\|\Phi(t)\Phi^{-1}(s)\| \leq e^{-\gamma_1(t-s)/m} \tag{5.24}$$

and hence for every $T > 0, p \geq 1$,

$$\begin{aligned} \sup_{t \in [0, T]} \|p_t^m\|^p &\leq \sup_{t \in [0, T]} 3^{p-1} \left(e^{-\gamma_1 pt/m} m^p \|v_0\|^p \right. \\ &\quad \left. + \left(\int_0^t e^{-\gamma_1(t-s)/m} \|F(u_s^m)\| ds \right)^p \right. \\ &\quad \left. + \left\| \Phi(t) \int_0^t \Phi^{-1}(s) \sigma(u_s^m) dW_s \right\|^p \right) \\ &\leq 3^{p-1} \left(m^p \|v_0\|^p + \frac{m^p}{\gamma_1^p} \|F\|_\infty^p \right. \\ &\quad \left. + \sup_{t \in [0, T]} \left\| \Phi(t) \int_0^t \Phi^{-1}(s) \sigma(u_s^m) dW_s \right\|^p \right), \end{aligned} \tag{5.25}$$

where $\|F\|_\infty$ denotes the supremum of $\|F(u)\|$ over u and we have employed the inequality

$$\left(\sum_{i=1}^N a_i \right)^p \leq N^{p-1} \sum_{i=1}^N a_i^p \tag{5.26}$$

for every $p \geq 1, N \in \mathbb{N}$.

Taking the p th power of Eq. (5.4), for any δ with $0 < \delta < T$ we have the P -a.s. bound

$$\begin{aligned} &\sup_{t \in [0, T]} \left\| \Phi(t) \int_0^t \Phi^{-1}(s) \sigma(u_s^m) dW_s \right\|^p \\ &\leq 2^{p-1} \left(1 + \frac{4}{\gamma_1} \sup_{s \in [0, T]} \|\gamma(u_s^m)\| \right)^p \left(e^{-p\delta\gamma_1/m} \sup_{t \in [0, T]} \left\| \int_0^t \sigma(u_r^m) dW_r \right\|^p \right. \\ &\quad \left. + \max_{k=0, \dots, N-1} \sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t \sigma(u_r^m) dW_r \right\|^p \right) \end{aligned} \tag{5.27}$$

where $N = \max\{k \in \mathbb{Z} : k\delta < T\}$.

We now return to bounding the momentum using Eq. (5.25). As was done in Eq. (5.25), the supremum of a quantity $\|A(u)\|$ will be denoted by $\|A\|_\infty$

for an arbitrary matrix- or vector-valued function A (rather than by the more precise but less readable $\| \| A \| \|_\infty$).

$$\begin{aligned} \sup_{t \in [0, T]} \| p_t^m \|^p &\leq 3^{p-1} \left[m^p \| v_0 \|^p + \frac{m^p}{\gamma_1^p} \| F \|^p_\infty + 2^{p-1} \left(1 + \frac{4}{\gamma_1} \| \gamma \|_\infty \right)^p \right. \\ &\quad \times \left(e^{-p\delta\gamma_1/m} \sup_{t \in [0, T]} \left\| \int_0^t \sigma(u_r^m) dW_r \right\|^p \right. \\ &\quad \left. \left. + \left(\sum_{k=0}^{N-1} \sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t \sigma(u_r^m) dW_r \right\|^{pq} \right)^{1/q} \right) \right] \end{aligned} \tag{5.28}$$

where we used Eq. (5.27) and the fact that the supremum norm on \mathbb{R}^N is bounded by the ℓ^q norm for any $q \geq 1$. We will take $q > 1$.

Taking the expected value and then using Hölder’s inequality on the expectations, we get

$$\begin{aligned} E \left[\sup_{t \in [0, T]} \| p_t^m \|^p \right] &\leq 3^{p-1} \left[m^p \| v_0 \|^p + \frac{m^p}{\gamma_1^p} \| F \|^p_\infty \right. \\ &\quad + 2^{p-1} \left(1 + \frac{4}{\gamma_1} \| \gamma \|_\infty \right)^p \left(e^{-p\delta\gamma_1/m} E \left[\sup_{t \in [0, T]} \left\| \int_0^t \sigma(u_r^m) dW_r \right\|^{pq} \right]^{1/q} \right. \\ &\quad \left. \left. + \left(\sum_{k=0}^{N-1} E \left[\sup_{t \in [k\delta, (k+2)\delta]} \left\| \int_{k\delta}^t \sigma(u_r^m) dW_r \right\|^{pq} \right] \right)^{1/q} \right) \right]. \end{aligned} \tag{5.29}$$

The Burkholder–Davis–Gundy inequalities (see for example Theorem 3.28 in [1]), for $d > 1$ imply the existence of a constant $C_{d,n} > 0$ such that

$$E \left[\sup_{0 \leq s \leq T} \left\| \int_0^s \sigma(u_r^m) dW_r \right\|^d \right] \leq C_{d,n} E \left[\left(\int_0^T \| \sigma(u_r^m) \|_F^2 dr \right)^{d/2} \right] \tag{5.30}$$

where $\| \cdot \|_F$ denotes the Frobenius (or Hilbert–Schmidt) norm.

Therefore, letting $\delta = m^{1-\kappa}$ for $0 < \kappa < 1$, we find

$$\begin{aligned} E \left[\sup_{t \in [0, T]} \| p_t^m \|^p \right] &\leq 3^{p-1} \left[m^p \| v_0 \|^p + \frac{m^p}{\gamma_1^p} \| F \|^p_\infty + 2^{p-1} C_{pq,n}^{1/q} \left(1 + \frac{4}{\gamma_1} \| \gamma \|_\infty \right)^p \right. \\ &\quad \times \left(e^{-p\gamma_1/m^\kappa} E \left[\left(\int_0^T \| \sigma(u_r^m) \|_F^2 dr \right)^{pq/2} \right]^{1/q} \right. \\ &\quad \left. \left. + \left(\sum_{k=0}^{N-1} E \left[\left(\int_{k\delta}^{(k+2)\delta} \| \sigma(u_r^m) \|_F^2 dr \right)^{pq/2} \right] \right)^{1/q} \right) \right] \end{aligned}$$

$$\begin{aligned} &\leq 3^{p-1} \left[m^p \|v_0\|^p + \frac{m^p}{\gamma_1^p} \|F\|_\infty^p + 2^{p-1} C_{pq,n}^{1/q} \|\sigma\|_{F,\infty}^p \right. \\ &\quad \left. \times \left(1 + \frac{4}{\gamma_1} \|\gamma\|_\infty \right)^p \left(e^{-p\gamma_1/m^\kappa} T^{p/2} + 2^{p/2} \left(N\delta^{pq/2} \right)^{1/q} \right) \right], \end{aligned} \tag{5.31}$$

where we define $\|\sigma\|_{F,\infty} = \sup_u \|\sigma(u)\|_F$.
 $N\delta < T$, hence

$$N\delta^{pq/2} < T\delta^{pq/2-1} = Tm^{(1-\kappa)(pq/2-1)}. \tag{5.32}$$

Therefore,

$$E \left[\sup_{t \in [0,T]} \|p_t^m\|^p \right] = O(m^{(1-\kappa)(p/2-1/q)}). \tag{5.33}$$

For any $0 < \beta < p/2$ we can choose $0 < \kappa < 1$ and $q > 1$ so that $(1 - \kappa)(p/2 - 1/q) = \beta$, thereby proving the claim for $p \geq 1$.

For any $0 < p < 1$ and $0 < \beta < p/2$, take $q \geq 1$. Then $\beta q/p < q/2$ so, using Hölder’s inequality, we find

$$E \left[\sup_{t \in [0,T]} \|p_t^m\|^p \right] \leq E \left[\left(\sup_{t \in [0,T]} \|p_t^m\|^p \right)^{q/p} \right]^{p/q} = O(m^\beta). \tag{5.34}$$

□

If we do not take the supremum over t inside the expectation, we can prove a stronger decay result.

Proposition 5.2. *For any $q > 0$ and any $m_0 > 0$ there exists a $C > 0$ such that*

$$\sup_{t \in [0,\infty)} E[\|p_t^m\|^q] \leq Cm^{q/2} \tag{5.35}$$

for all $0 < m \leq m_0$.

Proof. Let $\gamma_1 > \alpha > 0$ and define the process $z_t^m = e^{\alpha t/m} p_t^m$. By Itô’s formula

$$\begin{aligned} z_t^m &= p_0^m + \int_0^t \frac{\alpha}{m} z_s^m ds + \int_0^t e^{\alpha s/m} \left(F(u_s^m) - \frac{1}{m} \gamma(u_s^m) p_s^m \right) ds \\ &\quad + \int_0^t e^{\alpha s/m} \sigma(u_s^m) dW_s \end{aligned} \tag{5.36}$$

$$\begin{aligned} &= p_0^m + \int_0^t \left[-\frac{1}{m} (\gamma(u_s^m) - \alpha) z_s^m + e^{\alpha s/m} F(u_s^m) \right] ds \\ &\quad + \int_0^t e^{\alpha s/m} \sigma(u_s^m) dW_s. \end{aligned} \tag{5.37}$$

This holds for any α , but we will need $0 < \alpha < \gamma_1$ later.

Now, applying the Itô formula again to $\|z_t^m\|^2 = (z_t^m)^T z_t^m$ we find

$$\begin{aligned} \|z_t^m\|^2 &= \|p_0^m\|^2 + 2 \left(\int_0^t (z_s^m)^T \left(-\frac{1}{m}(\gamma(u_s^m) - \alpha)z_s^m + e^{\alpha s/m} F(u_s^m) \right) ds \right. \\ &\quad \left. + \int_0^t e^{\alpha s/m} (z_s^m)^T \sigma(u_s^m) dW_s \right) \\ &\quad + \sum_i \left[\sum_j \int_0^t e^{\alpha s/m} \sigma_j^i(u_s^m) dW_s^j, \sum_j \int_0^t e^{\alpha s/m} \sigma_j^i(u_s^m) dW_s^j \right]_t. \end{aligned} \quad (5.38)$$

The quadratic variation term is

$$\begin{aligned} &\sum_i \left[\sum_j \int_0^t e^{\alpha s/m} \sigma_j^i(u_s^m) dW_s^j, \sum_j \int_0^t e^{\alpha s/m} \sigma_j^i(u_s^m) dW_s^j \right]_t \\ &= \int_0^t e^{2\alpha s/m} \|\sigma(u_s^m)\|_F^2 ds. \end{aligned} \quad (5.39)$$

Therefore,

$$\begin{aligned} \|z_t^m\|^2 &= \|p_0^m\|^2 - \frac{2}{m} \int_0^t (z_s^m)^T (\gamma(u_s^m) - \alpha) z_s^m ds + 2 \int_0^t e^{\alpha s/m} (z_s^m)^T F(u_s^m) ds \\ &\quad + \int_0^t e^{2\alpha s/m} \|\sigma(u_s^m)\|_F^2 ds + 2 \int_0^t e^{\alpha s/m} (z_s^m)^T \sigma(u_s^m) dW_s. \end{aligned} \quad (5.40)$$

First we will show the result for $q = 2p$ with p a positive integer. Using Itô's formula one more time, we obtain

$$\begin{aligned} \|z_t^m\|^{2p} &= \|p_0^m\|^{2p} - \frac{2}{m} \int_0^t p \|z_s^m\|^{2(p-1)} (z_s^m)^T (\gamma(u_s^m) - \alpha) z_s^m ds \\ &\quad + 2 \int_0^t p \|z_s^m\|^{2(p-1)} e^{\alpha s/m} (z_s^m)^T F(u_s^m) ds \\ &\quad + \int_0^t p \|z_s^m\|^{2(p-1)} e^{2\alpha s/m} \|\sigma(u_s^m)\|_F^2 ds \\ &\quad + \frac{p(p-1)}{2} \int_0^t \|z_s^m\|^{2(p-2)} 4e^{2\alpha s/m} \|\sigma^T(u_s^m) z_s^m\|^2 ds \\ &\quad + 2 \int_0^t p \|z_s^m\|^{2(p-1)} e^{\alpha s/m} (z_s^m)^T \sigma(u_s^m) dW_s. \end{aligned} \quad (5.41)$$

By Assumption 1, we have $y^T \gamma(u) y \geq \gamma_1 \|y\|^2$ for all $u \in F_O(M)$, $y \in \mathbb{R}^n$. Hence, defining $\epsilon = \gamma_1 - \alpha > 0$, we deduce from here the following upper bound on the norm of p_t^m .

$$\begin{aligned} \|p_t^m\|^{2p} &\leq e^{-2p\alpha t/m} \|p_0^m\|^{2p} - \frac{2p\epsilon}{m} \int_0^t e^{-2p\alpha(t-s)/m} \|p_s^m\|^{2p} ds \\ &\quad + 2p \int_0^t e^{-2p\alpha(t-s)/m} \|p_s^m\|^{2(p-1)} (p_s^m)^T F(u_s^m) ds \end{aligned}$$

$$\begin{aligned}
 &+ p \int_0^t e^{-2p\alpha(t-s)/m} \|p_s^m\|^{2(p-1)} \|\sigma(u_s^m)\|_F^2 ds \\
 &+ 2p(p-1) \int_0^t e^{-2p\alpha(t-s)/m} \|p_s^m\|^{2(p-2)} \|\sigma^T(u_s^m)p_s^m\|^2 ds \\
 &+ 2p \int_0^t e^{-2p\alpha(t-s)/m} \|p_s^m\|^{2(p-1)} (p_s^m)^T \sigma(u_s^m) dW_s. \tag{5.42}
 \end{aligned}$$

Using Lemma 5.2, the following computation shows that

$$M_t \equiv \int_0^t e^{2p\alpha s/m} \|p_s^m\|^{2(p-1)} (p_s^m)^T \sigma(u_s^m) dW_s \tag{5.43}$$

is a martingale.

$$\begin{aligned}
 &E \left[\int_0^t \|e^{2p\alpha s/m} \|p_s^m\|^{2(p-1)} (p_s^m)^T \sigma(u_s^m)\|^2 ds \right] \\
 &\leq e^{4p\alpha t/m} \|\sigma\|_\infty^2 t E \left[\sup_{s \in [0,t]} \|p_s^m\|^{4p-2} \right] < \infty, \tag{5.44}
 \end{aligned}$$

where in the last step, the expectation is finite by Proposition 5.1. Using this and taking the expected value of Eq. (5.42) we find, for any $a > 0$,

$$\begin{aligned}
 E[\|p_t^m\|^{2p}] &\leq e^{-2p\alpha t/m} \|p_0^m\|^{2p} - \frac{2p\epsilon}{m} \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2p}] ds \\
 &+ 2p \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)} (p_s^m)^T F(u_s^m)] ds \\
 &+ p \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)} \|\sigma(u_s^m)\|_F^2] ds \\
 &+ 2p(p-1) \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-2)} \|\sigma^T(u_s^m)p_s^m\|^2] ds \tag{5.45}
 \end{aligned}$$

$$\begin{aligned}
 &\leq e^{-2p\alpha t/m} \|p_0^m\|^{2p} - \frac{2p\epsilon}{m} \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2p}] ds \\
 &+ p \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)} (a^2 \|p_s^m\|^2 + \|F\|_\infty^2/a^2)] ds \\
 &+ (p\|\sigma\|_{F,\infty}^2 + 2p(p-1)\|\sigma\|_\infty^2) \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)}] ds \\
 &= e^{-2p\alpha t/m} \|p_0^m\|^{2p} - p \left(\frac{2\epsilon}{m} - a^2 \right) \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2p}] ds \tag{5.46}
 \end{aligned}$$

$$\begin{aligned}
 &+ p\|F\|_\infty^2/a^2 \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)}] ds \\
 &+ (p\|\sigma\|_{F,\infty}^2 + 2p(p-1)\|\sigma\|_\infty^2) \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)}] ds. \tag{5.47}
 \end{aligned}$$

Letting $a^2 = 2\epsilon/m$ yields

$$E[\|p_t^m\|^{2p}] \leq e^{-2p\alpha t/m} \|p_0^m\|^{2p} + \left(\frac{pm}{2\epsilon} \|F\|_\infty^2 + p\|\sigma\|_{F,\infty}^2 + 2p(p-1)\|\sigma\|_\infty^2 \right) \times \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)}] ds. \tag{5.48}$$

For $p = 1$, this becomes

$$E[\|p_t^m\|^2] \leq m^2 e^{-2\alpha t/m} \|v_0\|^2 + \frac{m^2}{4\epsilon\alpha} \|F\|_\infty^2 + \frac{m}{2\alpha} \|\sigma\|_{F,\infty}^2. \tag{5.49}$$

Hence,

$$\sup_{t \in [0, \infty)} E[\|p_t^m\|^2] \leq m^2 \|v_0\|^2 + \frac{m^2}{4\epsilon\alpha} \|F\|_\infty^2 + \frac{m}{2\alpha} \|\sigma\|_{F,\infty}^2 \leq Cm \tag{5.50}$$

for all $0 < m \leq m_0$. So the claim holds for $p = 1$.

Now suppose it holds for $p - 1$. Then

$$\sup_{t \in [0, \infty)} E[\|p_t^m\|^{2p}] \leq m^{2p} \|v_0\|^{2p} + \left(\frac{pm}{2\epsilon} \|F\|_\infty^2 + p\|\sigma\|_{F,\infty}^2 + 2p(p-1)\|\sigma\|_\infty^2 \right) \times \sup_{t \in [0, \infty)} \int_0^t e^{-2p\alpha(t-s)/m} E[\|p_s^m\|^{2(p-1)}] ds \tag{5.51}$$

$$\leq m^{2p} \|v_0\|^{2p} + \left(\frac{pm}{2\epsilon} \|F\|_\infty^2 + p\|\sigma\|_{F,\infty}^2 + 2p(p-1)\|\sigma\|_\infty^2 \right) \times \sup_{t \in [0, \infty)} \int_0^t e^{-2p\alpha(t-s)/m} ds \sup_{t \in [0, \infty)} E[\|p_t^m\|^{2(p-1)}] \tag{5.52}$$

$$\leq m^{2p} \|v_0\|^{2p} + Cm^{p-1} \frac{m}{2p\alpha} \times \left(\frac{pm}{2\epsilon} \|F\|_\infty^2 + p\|\sigma\|_{F,\infty}^2 + 2p(p-1)\|\sigma\|_\infty^2 \right) \leq \tilde{C}m^p \tag{5.53}$$

for all $0 < m \leq m_0$. This proves the claim for all positive integers p by induction.

Finally, let $q > 0$ be arbitrary. Take $p \in \mathbb{N}$ with $2p > q$. Using Hölder's inequality we find

$$\sup_{t \in [0, \infty)} E[\|p_t^m\|^q] \leq \left(\sup_{t \in [0, \infty)} E[\|p_t^m\|^{2p}] \right)^{q/(2p)} \leq (Cm^p)^{q/(2p)} = \tilde{C}m^{q/2} \tag{5.54}$$

for all $0 < m \leq m_0$. Therefore, the result holds for any $q > 0$. □

We can use the above decay results to prove that certain integrals with respect to the momentum process also vanish in the limit.

Proposition 5.3. *For any smooth function f on $F_O(M)$, any $T > 0, p > 0$, and any $\alpha, \beta = 1, \dots, n$ we have*

$$E \left[\sup_{0 \leq t \leq T} \left| \int_0^t f(u_s^m) d((p_s^m)^\alpha (p_s^m)^\beta) \right|^p \right] = O(m^{p/2}) \text{ as } m \rightarrow 0, \tag{5.55}$$

where α, β refer to the components of the momentum process in the standard basis for \mathbb{R}^n .

Proof. First assume $p > 1$. Integrating by parts, and using the fact that u^m is pathwise C^1 , we get

$$\begin{aligned} \int_0^t f(u_s^m) d((p_s^m)^\alpha (p_s^m)^\beta) &= f(u_t^m) (p_t^m)^\alpha (p_t^m)^\beta - f(u_0^m) (p_0^m)^\alpha (p_0^m)^\beta \\ &\quad - \int_0^t (p_s^m)^\alpha (p_s^m)^\beta \frac{d}{ds} f(u_s^m) ds. \end{aligned} \tag{5.56}$$

From the original SDE, Eq. (4.13), u^m satisfies the ODE $\dot{u} = H_\nu(u)$, so

$$\frac{d}{ds} f(u_s^m) = \frac{1}{m} H_{p_s^m}(u_s^m)[f]. \tag{5.57}$$

$F_O(M)$ is compact, hence f is bounded. Therefore, decomposing the vector p_s^m in the standard basis e_ν of \mathbb{R}^n and using the notation H_ν introduced in Lemma 3.2, we have

$$\begin{aligned} &E \left[\sup_{0 \leq t \leq T} \left| \int_0^t f(u_s^m) d((p_s^m)^\alpha (p_s^m)^\beta) \right|^p \right] \\ &\leq 3^{p-1} \left(E \left[\sup_{0 \leq t \leq T} |f(u_t^m) (p_t^m)^\alpha (p_t^m)^\beta|^p \right] + E \left[|f(u_0^m) (p_0^m)^\alpha (p_0^m)^\beta|^p \right] \right. \\ &\quad \left. + E \left[\sup_{0 \leq t \leq T} \left| \int_0^t (p_s^m)^\alpha (p_s^m)^\beta \frac{1}{m} H_{p_s^m}(u_s^m)[f] ds \right|^p \right] \right) \\ &\leq 3^{p-1} \left(\|f\|_\infty^p E \left[\sup_{0 \leq t \leq T} \|p_t^m\|^{2p} \right] + m^{2p} \|f\|_\infty^p \|v_0\|^{2p} \right) \\ &\quad + \frac{3^{p-1}}{m^p} E \left[\sup_{0 \leq t \leq T} \left| \int_0^t (p_s^m)^\alpha (p_s^m)^\beta (p_s^m)^\nu H_\nu(u_s^m)[f] ds \right|^p \right]. \end{aligned} \tag{5.58}$$

We have assumed $p > 1$, so we can use Hölder’s inequality with exponents p and $p/(p-1)$ to estimate the last term. Using boundedness of $H_\nu[f]$ (implied by compactness of $F_O(M)$), this gives

$$\begin{aligned} &E \left[\sup_{0 \leq t \leq T} \left| \int_0^t (p_s^m)^\alpha (p_s^m)^\beta (p_s^m)^\nu H_\nu(u_s^m)[f] ds \right|^p \right] \\ &\leq E \left[\left(\int_0^T |(p_s^m)^\alpha (p_s^m)^\beta (p_s^m)^\nu H_\nu(u_s^m)[f]| ds \right)^p \right] \\ &\leq T^{p-1} E \left[\int_0^T |(p_s^m)^\alpha (p_s^m)^\beta (p_s^m)^\nu H_\nu(u_s^m)[f]|^p ds \right] \end{aligned}$$

$$\begin{aligned} &\leq CT^{p-1} E \left[\int_0^T \|p_s^m\|^{3p} ds \right] = CT^{p-1} \int_0^T E [\|p_s^m\|^{3p}] ds \\ &\leq CT^p \sup_{0 \leq s \leq T} E[\|p_s^m\|^{3p}] \end{aligned} \tag{5.59}$$

for some $C > 0$.

By Proposition 5.1, for any $0 < \kappa < p$ we obtain

$$\begin{aligned} &E \left[\sup_{0 \leq t \leq T} \left| \int_0^t f(u_s^m) d((p_s^m)^\alpha (p_s^m)^\beta) \right|^p \right] \\ &\leq O(m^\kappa) + \frac{3^{p-1}CT^p}{m^p} \sup_{0 \leq s \leq T} E [\|p_s^m\|^{3p}]. \end{aligned} \tag{5.60}$$

Applying Proposition 5.2 to the second term we get

$$\begin{aligned} &E \left[\sup_{0 \leq t \leq T} \left| \int_0^t f(u_s^m) d((p_s^m)^\alpha (p_s^m)^\beta) \right|^p \right] \\ &\leq O(m^\kappa) + \frac{3^{p-1}CT^p}{m^p} \tilde{C}m^{3p/2} \\ &\leq O(m^\kappa) + O(m^{p/2}). \end{aligned} \tag{5.61}$$

Taking $\kappa = p/2$ gives the result for all $p > 1$. The result for arbitrary $p > 0$ then follows by an application of Hölder’s inequality, as in Propositions 5.1 and 5.2. \square

6. Calculation of the Limiting SDE

We now manipulate the SDE, Eqs. (4.13)–(4.14), for (u_t^m, v_t^m) to extract the terms that survive when $m \rightarrow 0$ and thereby a candidate for the limiting SDE. The actual convergence proof will be given in Sect. 7.

To express the equations in terms of Lebesgue and Itô integrals on some \mathbb{R}^l , we consider the composition of a function $f \in C^\infty(F_O(M))$ with u_t^m . The following equations are then satisfied on \mathbb{R} and \mathbb{R}^n , respectively,

$$df(u_t^m) = H_{v_t^m}(u_t^m)[f]dt, \tag{6.1}$$

$$dv_t^m = \frac{1}{m}(F(u_t^m) - \gamma(u_t^m)v_t^m)dt + \frac{1}{m}\sigma(u_t^m)dW_t. \tag{6.2}$$

We know that the momentum, p_t^m , converges to zero (in various senses) and our objective will be to separate out such terms. We begin by solving for $v_t^m dt$ in the equation for dv_t^m . Note that, by assumption, $\gamma^s(u)$ is positive definite for all u , and hence $\gamma(u)$ is invertible for every u . Therefore,

$$v_t^m dt = -\gamma^{-1}(u_t^m)dp_t^m + \gamma^{-1}(u_t^m)F(u_t^m)dt + \gamma^{-1}(u_t^m)\sigma(u_t^m)dW_t. \tag{6.3}$$

Using the linearity of H_v in v , the first equation can be written

$$df(u_t^m) = (v_t^m)^\nu H_\nu(u_t^m)[f]dt, \tag{6.4}$$

where we are again using the notation H_ν introduced in Lemma 3.2 and components of the \mathbb{R}^n -valued process v_t^m will always refer to the standard basis. For any $n \times l$ matrix A , we will let $H_A(u)$ denote the element of $(\mathbb{R}^l)^*$ with action $w \mapsto H_{Aw}(u)$. For any \mathbb{R}^l -valued semimartingale, X , we will also write $H_A(u)dX$ as a shorthand for the contraction $H_{Ae_\eta}(u)dX^\eta$.

With these notations, after substituting Eq. (6.3) into Eq. (6.4) we obtain

$$\begin{aligned} df(u_t^m) &= -H_\nu(u_t^m)[f](\gamma^{-1})^\nu_\mu(u_t^m)d(p_t^m)^\mu \\ &\quad + H_\nu(u_t^m)[f](\gamma^{-1})^\nu_\mu(u_t^m)F^\mu(u_t^m)dt \\ &\quad + H_\nu(u_t^m)[f](\gamma^{-1})^\nu_\mu(u_t^m)\sigma_\eta^\mu(u_t^m)dW_t^\eta \\ &= -H_{\gamma^{-1}(u_t^m)}(u_t^m)[f]d(p_t^m) + H_{(\gamma^{-1}F)(u_t^m)}(u_t^m)[f]dt \\ &\quad + H_{(\gamma^{-1}\sigma)(u_t^m)}(u_t^m)[f]dW_t. \end{aligned} \tag{6.5}$$

Remark 2. Here and in the following, components of matrix- or \mathbb{R}^n -valued functions on the frame bundle, such as $\gamma(u)$ or $F(u)$, as well as the implied sums over repeated indices, will always refer to the standard basis. We emphasize that these components are unrelated to local coordinates on M or $F_O(M)$ and do not in any way imply that the statements have a local character. Rather, by lifting a tensor or vector field from M to a matrix- or vector-valued function on $F_O(M)$, we are able to speak about its components in each frame in a globally defined manner.

The second and third terms in Eq. (6.5) are independent of the momentum, so we focus on the first term. Integrating by parts and using the fact that u_t^m is pathwise C^1 and satisfies the ODE (4.13) we get

$$\begin{aligned} &H_{\gamma^{-1}(u_t^m)}(u_t^m)[f]d(p_t^m) \\ &= d(H_{\gamma^{-1}(u_t^m)}p_t^m(u_t^m)[f]) - (p_t^m)^\nu d(H_{\gamma^{-1}(u_t^m)}e_\nu(u_t^m)[f]) \\ &= d(H_{\gamma^{-1}(u_t^m)}p_t^m(u_t^m)[f]) - (p_t^m)^\nu \frac{d}{dt}(H_{\gamma^{-1}(u_t^m)}e_\nu(u_t^m)[f])dt \\ &= d(H_{\gamma^{-1}(u_t^m)}p_t^m(u_t^m)[f]) - (p_t^m)^\nu H_{v_t^m}(u_t^m)[(\gamma^{-1})^\mu_\nu H_\mu[f]]dt \\ &= d(H_{\gamma^{-1}(u_t^m)}p_t^m(u_t^m)[f]) - m(v_t^m)^\nu (v_t^m)^\xi K_{\nu\xi}^f(u_t^m)dt \end{aligned} \tag{6.6}$$

by Eq. (6.4), where we define

$$K_{\nu\xi}^f(u) = H_\xi(u)[(\gamma^{-1})^\mu_\nu]H_\mu(u)[f] + (\gamma^{-1})^\mu_\nu(u)H_\xi(u)[H_\mu[f]]. \tag{6.7}$$

The equation for $f(u_t^m)$ then becomes

$$\begin{aligned} df(u_t^m) &= -d(H_{\gamma^{-1}(u_t^m)}p_t^m(u_t^m)[f]) + m(v_t^m)^\nu (v_t^m)^\mu K_{\nu\mu}^f(u_t^m)dt \\ &\quad + H_{(\gamma^{-1}F)(u_t^m)}(u_t^m)[f]dt + H_{(\gamma^{-1}\sigma)(u_t^m)}(u_t^m)[f]dW_t. \end{aligned} \tag{6.8}$$

To simplify $m(v_t^m)^\nu (v_t^m)^\mu dt$, we follow [13] and compute

$$\begin{aligned} d(m(v_t^m)^\nu m(v_t^m)^\mu) &= m(v_t^m)^\nu d(m(v_t^m)^\mu) + m(v_t^m)^\mu d(m(v_t^m)^\nu) \\ &\quad + d([m(v_t^m)^\nu, m(v_t^m)^\mu]) \end{aligned}$$

$$\begin{aligned}
 &= m(v_t^m)^\nu d(m(v_t^m)^\mu) + m(v_t^m)^\mu d(m(v_t^m)^\nu) \\
 &\quad + \sum_{\delta} \sigma_{\delta}^{\nu}(u_t^m) \sigma_{\delta}^{\mu}(u_t^m) dt, \tag{6.9}
 \end{aligned}$$

where we have used the SDE for $(v_t^m)^\nu$, Eq. (6.2). Defining

$$\Sigma^{\nu\mu}(u) = \sum_{\delta} \sigma_{\delta}^{\nu}(u) \sigma_{\delta}^{\mu}(u) \tag{6.10}$$

and using the SDE for $(v_t^m)^\nu$ again, we get

$$\begin{aligned}
 d(m(v_t^m)^\nu m(v_t^m)^\mu) &= m(v_t^m)^\nu (F^\mu(u_t^m) - \gamma_{\xi}^{\mu}(u_t^m)(v_t^m)^{\xi}) dt \\
 &\quad + m(v_t^m)^\nu \sigma_{\eta}^{\mu}(u_t^m) dW_t^{\eta} \\
 &\quad + m(v_t^m)^\mu (F^\nu(u_t^m) - \gamma_{\xi}^{\nu}(u_t^m)(v_t^m)^{\xi}) dt \\
 &\quad + m(v_t^m)^\mu \sigma_{\eta}^{\nu}(u_t^m) dW_t^{\eta} + \Sigma^{\nu\mu}(u_t^m) dt. \tag{6.11}
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 &m(v_t^m)^\nu \gamma_{\xi}^{\mu}(u_t^m)(v_t^m)^{\xi} dt + m(v_t^m)^\mu \gamma_{\xi}^{\nu}(u_t^m)(v_t^m)^{\xi} dt \\
 &= -d(m(v_t^m)^\nu m(v_t^m)^\mu) + ((p_t^m)^\nu F^\mu(u_t^m) + (p_t^m)^\mu F^\nu(u_t^m)) dt \\
 &\quad + ((p_t^m)^\nu \sigma_{\eta}^{\mu}(u_t^m) + (p_t^m)^\mu \sigma_{\eta}^{\nu}(u_t^m)) dW_t^{\eta} + \Sigma^{\nu\mu}(u_t^m) dt. \tag{6.12}
 \end{aligned}$$

For scalar γ , one can immediately solve for $m(v_t^m)^\nu (v_t^m)^\mu dt$. To handle the general case, we utilize the technique developed in [13]. First we rewrite Eq. (6.12) in integral notation:

$$\begin{aligned}
 &\int_0^t \left(m(v_s^m)^\nu \gamma_{\xi}^{\mu}(u_s^m)(v_s^m)^{\xi} + m(v_s^m)^\mu \gamma_{\xi}^{\nu}(u_s^m)(v_s^m)^{\xi} \right) ds \\
 &= -m(v_t^m)^\nu m(v_t^m)^\mu + m(v_0)^\nu m(v_0)^\mu \\
 &\quad + \int_0^t ((p_s^m)^\nu F^\mu(u_s^m) + (p_s^m)^\mu F^\nu(u_s^m)) ds \\
 &\quad + \int_0^t ((p_s^m)^\nu \sigma_{\eta}^{\mu}(u_s^m) + (p_s^m)^\mu \sigma_{\eta}^{\nu}(u_s^m)) dW_s^{\eta} + \int_0^t \Sigma^{\nu\mu}(u_s^m) ds. \tag{6.13}
 \end{aligned}$$

The formula for the left-hand side implies that the right-hand side, which we denote by $C_t^{\mu\nu}$, is C^1 P -a.s. and so we can differentiate both sides with respect to t to obtain

$$m(v_t^m)^\nu \gamma_{\xi}^{\mu}(u_t^m)(v_t^m)^{\xi} + m(v_t^m)^\mu \gamma_{\xi}^{\nu}(u_t^m)(v_t^m)^{\xi} = \dot{C}_t^{\mu\nu}. \tag{6.14}$$

Define the matrix $V_t^{\nu\mu} = m(v_t^m)^\nu (v_t^m)^\mu$. The above equation can be written in matrix form

$$\gamma V + V \gamma^T = \dot{C}. \tag{6.15}$$

This is a Lyapunov equation, where $-\gamma$ has eigenvalues with real parts bounded above by $-\gamma_1$. Hence, we can solve uniquely for V ,

$$V = \int_0^\infty e^{-y\gamma} \dot{C} e^{-y\gamma^T} dy, \tag{6.16}$$

see for example Theorem 6.4.2 in [32]. Integrating with respect to t , we get

$$\int_0^t m(v_s^m)^\nu (v_s^m)^\mu ds = \int_0^t \int_0^\infty (e^{-y\gamma(u_s^m)})_\eta^\nu (e^{-y\gamma(u_s^m)})_\xi^\mu dy (\dot{C}_s)^{\eta\xi} ds. \tag{6.17}$$

Define the functions $G_{\eta\xi}^{\nu\mu}(u) = \int_0^\infty (e^{-y\gamma(u)})_\eta^\nu (e^{-y\gamma(u)})_\xi^\mu dy$. Using the dominated convergence theorem, along with the formula for the derivative of the matrix exponential found in [33], one can prove that G is a smooth function of u . Therefore, the $G_{\eta\xi}^{\nu\mu}(u_t^m)$ are semimartingales and

$$\begin{aligned} \int_0^t m(v_s^m)^\beta (v_s^m)^\alpha ds &= \int_0^t G_{\mu\nu}^{\beta\alpha}(u_s^m) dC_s^{\mu\nu} = - \int_0^t G_{\mu\nu}^{\beta\alpha}(u_s^m) d((p_s^m)^\nu (p_s^m)^\mu) \\ &+ \int_0^t G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\nu F^\mu(u_s^m) + (p_s^m)^\mu F^\nu(u_s^m)) ds \\ &+ \int_0^t G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\mu \sigma_\eta^\nu(u_s^m) + (p_s^m)^\nu \sigma_\eta^\mu(u_s^m)) dW_s^\eta \\ &+ \int_0^t G_{\mu\nu}^{\beta\alpha}(u_s^m) \Sigma^{\mu\nu}(u_s^m) ds. \end{aligned} \tag{6.18}$$

Note that if we define

$$J^{\beta\alpha}(u) = G_{\mu\nu}^{\beta\alpha}(u) \Sigma^{\mu\nu}(u) \tag{6.19}$$

then J is symmetric and is the unique solution to the Lyapunov equation

$$\gamma J + J \gamma^T = \Sigma. \tag{6.20}$$

Using Eq. (6.18) we find that $f(u_t^m)$ satisfies

$$\begin{aligned} df(u_t^m) &= H_{(\gamma^{-1}F)(u_t^m)}(u_t^m)[f]dt + J^{\beta\alpha}(u_t^m) K_{\beta\alpha}^f(u_t^m)dt \\ &+ H_{(\gamma^{-1}\sigma)(u_t^m)}(u_t^m)[f]dW_t \\ &- d(H_{\gamma^{-1}(u_t^m)p_t^m}(u_t^m)[f]) - K_{\beta\alpha}^f(u_t^m) G_{\mu\nu}^{\beta\alpha}(u_t^m) d((p_t^m)^\nu (p_t^m)^\mu) \\ &+ K_{\beta\alpha}^f(u_t^m) G_{\mu\nu}^{\beta\alpha}(u_t^m) ((p_t^m)^\nu F^\mu(u_t^m) + (p_t^m)^\mu F^\nu(u_t^m)) dt \\ &+ K_{\beta\alpha}^f(u_t^m) G_{\mu\nu}^{\beta\alpha}(u_t^m) ((p_t^m)^\mu \sigma_\eta^\nu(u_t^m) + (p_t^m)^\nu \sigma_\eta^\mu(u_t^m)) dW_t^\eta. \end{aligned} \tag{6.21}$$

Based on the results of Sect. 5, we expect that terms involving p_t^m will converge to zero as $m \rightarrow 0$. Therefore, if it exists, we would expect the limiting process u_t to satisfy

$$df(u_t) = H_{(\gamma^{-1}F)(u_t)}(u_t)[f]dt + J^{\beta\alpha}(u_t) K_{\beta\alpha}^f(u_t)dt + H_{(\gamma^{-1}\sigma)(u_t)}(u_t)[f]dW_t \tag{6.22}$$

for every $f \in C^\infty(F_O(M))$. Note that by Lemma 3.5, the first term is the horizontal lift of the vector field $(\gamma^{-1}F)(x)$ on M to $F_O(M)$,

$$H_{(\gamma^{-1}F)(u)}(u) = (\gamma^{-1}F)^h(u). \tag{6.23}$$

To express Eq. (6.22) as an SDE on the manifold $F_O(M)$ we need to rewrite it using the Stratonovich integral. This is accomplished by the following lemma.

Lemma 6.1. *An $F_O(M)$ -valued semimartingale u_t satisfies the SDE*

$$\begin{aligned} du_t &= (\gamma^{-1}F)^h(u_t)dt + H_{(\gamma^{-1}\sigma)(u_t)}(u_t) \circ dW_t \\ &\quad - \frac{1}{2} \sum_{\alpha} (\gamma^{-1}(u_t))_{\mu}^{\eta} \sigma_{\alpha}^{\mu}(u_t) (\gamma^{-1}(u_t))_{\nu}^{\xi} H_{\eta}(u_t) [\sigma_{\alpha}^{\nu}] H_{\xi}(u_t) dt \\ &\quad - \frac{1}{2} ((\gamma^{-1}(u_t))_{\mu}^{\eta} J^{\mu\chi}(u_t) \gamma_{\chi}^{\nu}(u_t) - J^{\eta\nu}(u_t)) H_{\eta}(u_t) [(\gamma^{-1})_{\nu}^{\xi}] H_{\xi}(u_t) dt \\ &\quad - \frac{1}{2} J^{\xi\nu}(u_t) (\gamma^{-1}(u_t))_{\nu}^{\eta} [H_{\eta}, H_{\xi}](u_t) dt \end{aligned} \quad (6.24)$$

iff

$$df(u_t) = H_{(\gamma^{-1}F)(u_t)}(u_t)[f]dt + J^{\beta\alpha}(u_t)K_{\beta\alpha}^f(u_t)dt + H_{(\gamma^{-1}\sigma)(u_t)}(u_t)[f]dW_t \quad (6.25)$$

for every $f \in C^{\infty}(F_O(M))$. The sum over α in Eq. (6.24) is taken over the standard basis for \mathbb{R}^k .

Proof. By Lemma 4.2, u_t satisfies the SDE (6.24) iff

$$\begin{aligned} df(u_t) &= H_{(\gamma^{-1}F)(u_t)}(u_t)[f]dt + H_{(\gamma^{-1}\sigma)(u_t)}(u_t)[f]dW_t \\ &\quad + \frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \sigma_{\alpha}^{\mu}(u_t) H_{\eta}(u_t) [(\gamma^{-1})_{\nu}^{\xi} \sigma_{\beta}^{\nu} H_{\xi}[f]] d[W^{\alpha}, W^{\beta}]_t \\ &\quad - \frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \sigma_{\alpha}^{\mu}(u_t) (\gamma^{-1}(u_t))_{\nu}^{\xi} H_{\eta}(u_t) [\sigma_{\beta}^{\nu}] H_{\xi}[f](u_t) \delta^{\alpha\beta} dt \\ &\quad - \frac{1}{2} ((\gamma^{-1}(u_t))_{\mu}^{\eta} J^{\mu\chi}(u_t) \gamma_{\chi}^{\nu}(u_t) - J^{\eta\nu}(u_t)) H_{\eta}(u_t) [(\gamma^{-1})_{\nu}^{\xi}] H_{\xi}(u_t) [f] dt \\ &\quad - \frac{1}{2} J^{\xi\nu}(u_t) (\gamma^{-1}(u_t))_{\nu}^{\eta} [H_{\eta}, H_{\xi}](u_t) [f] dt \end{aligned} \quad (6.26)$$

for all $f \in C^{\infty}(F_O(M))$.

The Lyapunov equation, Eq. (6.20), implies $\gamma^{-1}\Sigma(\gamma^{-1})^T = \gamma^{-1}J + J(\gamma^{-1})^T$. Together with the symmetry of J and Σ , this yields the following for any $F_O(M)$ -valued semimartingale u_t and any $f \in C^{\infty}(F_O(M))$.

$$\begin{aligned} &\frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \sigma_{\alpha}^{\mu}(u_t) H_{\eta}(u_t) [(\gamma^{-1})_{\nu}^{\xi} \sigma_{\beta}^{\nu} H_{\xi}[f]] d[W^{\alpha}, W^{\beta}]_t \\ &= \frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \sigma_{\alpha}^{\mu}(u_t) \left(H_{\eta}(u_t) [(\gamma^{-1})_{\nu}^{\xi}] \sigma_{\beta}^{\nu}(u_t) H_{\xi}(u_t) [f] \right. \\ &\quad \left. + (\gamma^{-1}(u_t))_{\nu}^{\xi} H_{\eta}(u_t) [\sigma_{\beta}^{\nu}] H_{\xi}[f](u_t) + (\gamma^{-1}(u_t))_{\nu}^{\xi} \sigma_{\beta}^{\nu}(u_t) H_{\eta}(u_t) [H_{\xi}[f]] \right) \delta^{\alpha\beta} dt \end{aligned} \quad (6.27)$$

$$\begin{aligned} &= \frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \Sigma^{\mu\nu}(u_t) H_{\eta}(u_t) [(\gamma^{-1})_{\nu}^{\xi}] H_{\xi}(u_t) [f] dt \\ &\quad + \frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \sigma_{\alpha}^{\mu}(u_t) (\gamma^{-1}(u_t))_{\nu}^{\xi} H_{\eta}(u_t) [\sigma_{\beta}^{\nu}] H_{\xi}[f](u_t) \delta^{\alpha\beta} dt \\ &\quad + \frac{1}{4} (\gamma^{-1}(u_t))_{\mu}^{\eta} \Sigma^{\mu\nu}(u_t) (\gamma^{-1}(u_t))_{\nu}^{\xi} (H_{\eta}(u_t) [H_{\xi}[f]] + H_{\xi}(u_t) [H_{\eta}[f]]) dt \quad (6.28) \\ &= \frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \Sigma^{\mu\nu}(u_t) H_{\eta}(u_t) [(\gamma^{-1})_{\nu}^{\xi}] H_{\xi}(u_t) [f] dt \\ &\quad + \frac{1}{2} (\gamma^{-1}(u_t))_{\mu}^{\eta} \sigma_{\alpha}^{\mu}(u_t) (\gamma^{-1}(u_t))_{\nu}^{\xi} H_{\eta}(u_t) [\sigma_{\beta}^{\nu}] H_{\xi}[f](u_t) \delta^{\alpha\beta} dt \end{aligned}$$

$$\begin{aligned}
 & + \frac{1}{2} J^{\xi\nu}(u_t)(\gamma^{-1}(u_t))_{\nu}^{\eta}([H_{\eta}, H_{\xi}](u_t)[f] + 2H_{\xi}(u_t)[H_{\eta}[f]]) dt \tag{6.29} \\
 = & J^{\xi\nu}(u_t)K_{\nu\xi}^f(u_t)dt + \frac{1}{2}(\gamma^{-1}(u_t))_{\mu}^{\eta}\sigma_{\alpha}^{\mu}(u_t)(\gamma^{-1}(u_t))_{\nu}^{\xi}H_{\eta}(u_t)[\sigma_{\beta}^{\nu}]H_{\xi}[f](u_t)\delta^{\alpha\beta}dt \\
 & + \frac{1}{2}((\gamma^{-1}(u_t))_{\mu}^{\eta}\Sigma^{\mu\nu}(u_t) - 2J^{\eta\nu}(u_t))H_{\eta}(u_t)[(\gamma^{-1})_{\nu}^{\xi}]H_{\xi}(u_t)[f]dt \\
 & + \frac{1}{2}J^{\xi\nu}(u_t)(\gamma^{-1}(u_t))_{\nu}^{\eta}[H_{\eta}, H_{\xi}](u_t)[f]dt. \tag{6.30}
 \end{aligned}$$

Using the Lyapunov equation, Eq. (6.20), one more time gives

$$\gamma^{-1}\Sigma - 2J = \gamma^{-1}J\gamma^T - J. \tag{6.31}$$

Therefore, combining the result of Eq. (6.27) with Eq. (6.26) yields Eq. (6.25). \square

The proposed limiting SDE, Eq. (6.24), includes a drift term generated by the vector field $S(u) = S^h(u) + S^v(u)$, where

$$\begin{aligned}
 S^h(u) = & -\frac{1}{2}\sum_{\alpha}(\gamma^{-1}(u_t))_{\mu}^{\eta}\sigma_{\alpha}^{\mu}(u_t)(\gamma^{-1}(u_t))_{\nu}^{\xi}H_{\eta}(u_t)[\sigma_{\alpha}^{\nu}]H_{\xi}(u_t) \\
 & -\frac{1}{2}((\gamma^{-1}(u_t))_{\mu}^{\eta}J^{\mu\chi}(u_t)\gamma'_{\chi}(u_t) - J^{\eta\nu}(u_t))H_{\eta}(u_t)[(\gamma^{-1})_{\nu}^{\xi}]H_{\xi}(u_t), \tag{6.32}
 \end{aligned}$$

$$S^v(u) = -\frac{1}{2}J^{\xi\nu}(u_t)(\gamma^{-1}(u_t))_{\nu}^{\eta}[H_{\eta}, H_{\xi}](u_t). \tag{6.33}$$

This is on top of the horizontal lift of the deterministic force to the frame bundle, $(\gamma^{-1}F)^h(u)$. $S^h(u)$ is a linear combination of the $H_{\alpha}(u)$ s, hence is a horizontal vector field on $F_O(M)$. It corresponds to an additional “force” on the particle’s position, which we call the *noise-induced drift*. A nonzero noise-induced drift requires either a non-trivial state dependence of the noise coefficients, σ_{ν}^{μ} , or non-trivial state dependence of the drag, together with $\gamma^{-1}J\gamma^T \neq J$, but it does not require any deterministic forcing, F , to be present in the original system. An analogous phenomenon was derived in Euclidean space of arbitrary dimension in [13]. See [34] for a recent review of noise-induced drift in systems with multiplicative noise.

For a torsion-free connection on M , such as the Levi-Civita connection employed here, the $[H_{\eta}, H_{\xi}]$ are vertical vector fields on $F_O(M)$ that can be expressed in terms of the curvature tensor [8]. Therefore, the S^v is a vertical vector field on $F_O(M)$ that results in an additional rotational “force” on the particle’s frame.

Note that if γ is a scalar then γ and J commute, hence the second term of S^h vanishes. S^v also vanishes in this case, due to the symmetry of $J^{\xi\eta}$ combined with the antisymmetry of $[H_{\eta}, H_{\xi}]$.

6.1. Some Special Cases

Before we prove convergence to the proposed limiting equation, Eq. (6.24), we will study its form in several cases of interest. For all these cases, we make the assumption that $k = n$ and σ_{ν}^{μ} comes from a $\binom{1}{1}$ -tensor field, $\sigma(x)$, on M , as

in Eq. (4.16). The following lemma will allow us to obtain a geometric formula for the noise-induced drift in this case. A similar study could be done in the case of Eq. (4.15), but we do not pursue this here.

Lemma 6.2. *Let σ, τ, κ , be $\binom{1}{1}$ -tensor fields on M and define the matrix-valued functions $\sigma(u), \tau(u), \kappa(u)$ as done in Eq. (4.16), i.e., $\sigma(u) = u^{-1}\sigma(\pi(u))u$ and similarly for τ, κ . Let Y be the vector field on M defined by*

$$Y^\chi(x) = g^{\nu\rho}(x)\tau_\nu^\beta(x)\kappa_\delta^\chi(x)(\nabla_\beta\sigma_\rho^\delta)(x). \quad (6.34)$$

The horizontal lift of Y is given by

$$Y^h(u) = \sum_\eta \tau_\eta^\nu(u)H_\nu(u)[\sigma_\eta^\xi]\kappa_\xi^\chi(u)H_\chi(u). \quad (6.35)$$

Proof. We work in the domain of a coordinate chart and local o.n. frame E_α and use the notation defined in Sect. 3.2. Let $\sigma_\nu^\mu(x)$ be the components of σ in the frame E_α . Therefore,

$$\sigma_\xi^\nu(u) = (h^{-1})_\mu^\nu\sigma_\alpha^\mu(x)h_\xi^\alpha \quad (6.36)$$

and similarly for τ, κ . Using Sect. 3.2 and the fact that $h \in O(\mathbb{R}^n)$, we obtain

$$\begin{aligned} \sum_\eta \tau_\eta^\nu(u)H_\nu(u)[\sigma_\eta^\xi] &= \sum_\eta \tau_\eta^\nu(u)h_\nu^\beta((\Lambda^{-1})_\beta^i(x)\partial_i\sigma_\eta^\xi(u) - h_\mu^\kappa A_{\beta\kappa}^\delta(x)\partial_{h_\mu^\delta}\sigma_\eta^\xi(u)) \\ &= \sum_\eta \tau_\nu^\beta(x)h_\eta^\nu((\Lambda^{-1})_\beta^i(x)\partial_i((h^{-1})_\chi^\xi\sigma_\rho^\chi(x)h_\eta^\rho) \\ &\quad - h_\mu^\kappa A_{\beta\kappa}^\delta(x)\partial_{h_\mu^\delta}((h^{-1})_\chi^\xi\sigma_\rho^\chi(x)h_\eta^\rho)) \end{aligned} \quad (6.37)$$

$$\begin{aligned} &= \tau_\nu^\beta(x) \left(\sum_\eta h_\eta^\rho h_\eta^\nu (\Lambda^{-1})_\beta^i(x) (h^{-1})_\chi^\xi \partial_i (\sigma_\rho^\chi)(x) \right. \\ &\quad \left. - \sum_\eta h_\eta^\nu h_\mu^\kappa A_{\beta\kappa}^\delta(x) \partial_{h_\mu^\delta} ((h^{-1})_\chi^\xi) \sigma_\rho^\chi(x) h_\eta^\rho \right. \\ &\quad \left. - \sum_\eta h_\eta^\nu h_\mu^\kappa A_{\beta\kappa}^\delta(x) (h^{-1})_\chi^\xi \sigma_\rho^\chi(x) \partial_{h_\mu^\delta} (h_\eta^\rho) \right) \end{aligned} \quad (6.38)$$

$$\begin{aligned} &= \tau_\nu^\beta(x) \left(\delta^{\rho\nu} (\Lambda^{-1})_\beta^i(x) (h^{-1})_\chi^\xi \partial_i (\sigma_\rho^\chi)(x) \right. \\ &\quad \left. - \delta^{\rho\nu} h_\mu^\kappa A_{\beta\kappa}^\delta(x) ((-h^{-1})\partial_{h_\mu^\delta}(h)(h^{-1})_\chi^\xi) \sigma_\rho^\chi(x) \right. \\ &\quad \left. - \sum_\eta h_\eta^\nu h_\mu^\kappa A_{\beta\kappa}^\delta(x) (h^{-1})_\chi^\xi \sigma_\rho^\chi(x) \delta_\delta^\rho \delta_\eta^\mu \right) \end{aligned} \quad (6.39)$$

$$\begin{aligned} &= \tau_\nu^\beta(x) \delta^{\nu\rho} ((\Lambda^{-1})_\beta^i(x) (h^{-1})_\chi^\xi \partial_i (\sigma_\rho^\chi)(x) \\ &\quad + h_\mu^\kappa A_{\beta\kappa}^\delta(x) (h^{-1})_\phi^\xi \partial_{h_\mu^\delta} (h_\psi^\phi) (h^{-1})_\chi^\psi \sigma_\rho^\chi(x) \\ &\quad - A_{\beta\rho}^\delta(x) (h^{-1})_\chi^\xi \sigma_\delta^\chi(x)) \end{aligned} \quad (6.40)$$

$$\begin{aligned}
 &= \tau_\nu^\beta(x) \delta^{\nu\rho} ((\Lambda^{-1})^i_\beta(x) (h^{-1})^\xi_\chi \partial_i (\sigma_\rho^\chi)(x) \\
 &\quad + h^\kappa_\mu A^\delta_{\beta\kappa}(x) (h^{-1})^\xi_\delta (h^{-1})^\mu_\chi \sigma_\rho^\chi(x) \\
 &\quad - A^\delta_{\beta\rho}(x) (h^{-1})^\xi_\chi \sigma_\delta^\chi(x)) \tag{6.41}
 \end{aligned}$$

$$\begin{aligned}
 &= \tau_\nu^\beta(x) \delta^{\nu\rho} ((\Lambda^{-1})^i_\beta(x) \partial_i (\sigma_\rho^\chi)(x) \\
 &\quad + A^\chi_{\beta\kappa}(x) \sigma_\rho^\kappa(x) - A^\delta_{\beta\rho}(x) \sigma_\delta^\chi(x)) (h^{-1})^\xi_\chi. \tag{6.42}
 \end{aligned}$$

In the last line, the terms in parentheses are the components of the covariant derivative of σ in the direction E_β ; therefore,

$$\sum_\eta \tau_\nu^\nu(u) H_\nu(u) [\sigma_\eta^\xi] = \tau_\nu^\beta(x) \delta^{\nu\rho} (\nabla_\beta \sigma)_\rho^\chi(x) (h^{-1})^\xi_\chi. \tag{6.43}$$

All tensor components are taken in the o.n. frame E_α , hence $g^{\mu\nu} = \delta^{\mu\nu}$. Therefore,

$$\sum_\eta \tau_\eta^\nu(u) H_\nu(u) [\sigma_\eta^\xi] \kappa_\xi^\chi(u) e_\chi = u^{-1} (g^{\nu\rho} \tau_\nu^\beta(x) (\nabla_\beta \sigma)_\rho^\alpha(x) \kappa_\alpha^\mu(x) E_\mu) = u^{-1} Y(x). \tag{6.44}$$

This proves

$$\sum_\eta \tau_\eta^\nu(u) H_\nu(u) [\sigma_\eta^\xi] \kappa_\xi^\chi(u) H_\chi(u) = H_{u^{-1}Y(\pi(u))} = Y^h(u). \tag{6.45}$$

□

Several applications of the above lemma can be used to show that, for tensor σ , the noise-induced drift is a horizontal lift.

Corollary 6.1. *Define the smooth tensor fields on M ,*

$$\Sigma^{\xi\eta}(x) = \sigma_\alpha^\xi(x) g^{\alpha\beta}(x) \sigma_\beta^\eta(x), \tag{6.46}$$

$$J^{\mu\nu}(x) = \int_0^\infty (e^{-y\gamma(x)})^\mu_\delta \Sigma^{\delta\eta}(x) (e^{-y\gamma(x)})^\nu_\eta dy, \tag{6.47}$$

$$L^{\mu\nu}(x) = \int_0^\infty (\gamma^{-1}(x) e^{-y\gamma(x)})^\mu_\alpha \Sigma^{\alpha\beta}(x) (\gamma(x) e^{-y\gamma(x)})^\nu_\beta dy. \tag{6.48}$$

Then the noise-induced drift vector field, S^h , is the horizontal lift of the vector field, Y , on M defined by

$$Y^\chi = -\frac{1}{2} g^{\nu\rho} (\gamma^{-1} \sigma)_\nu^\beta (\gamma^{-1})^\chi_\delta \nabla_\beta \sigma_\rho^\delta - \frac{1}{2} (L^{\beta\rho} - J^{\beta\rho}) \nabla_\beta (\gamma^{-1})^\chi_\rho. \tag{6.49}$$

An important special case is when the drag and noise satisfy the fluctuation–dissipation relation.

Corollary 6.2. (A particle satisfying the fluctuation–dissipation relation) *Suppose the fluctuation–dissipation relation is satisfied,*

$$\gamma_\nu^\mu(x) = \frac{1}{2k_B T} \sigma_\alpha^\mu(x) g^{\alpha\beta}(x) \sigma_\beta^\delta(x) g_{\delta\nu}(x), \tag{6.50}$$

where T is the temperature and k_B is the Boltzmann constant. Then $\gamma_\nu^\mu(u) = \frac{1}{2k_B T} \Sigma^{\mu\alpha}(u) \delta_{\alpha\nu}$, $J^{\mu\nu} = k_B T \delta^{\mu\nu}$, $S^v = 0$, and S^h is the horizontal lift of

$$Y^\chi = -k_B T g^{\beta\mu} (\sigma^{-1})_\mu^\rho (\gamma^{-1})_\delta^\chi \nabla_\beta \sigma_\rho^\delta. \tag{6.51}$$

Next, we specialize to scalar drag.

Corollary 6.3. (A particle with scalar drag) *If $\gamma_\nu^\mu(x) = \gamma(x) \delta_\nu^\mu$ for some $\gamma \in C^\infty(M)$ then $S^v = 0$ and*

$$S^h = -\frac{1}{2} (\gamma^{-2} Y)^h, \quad Y^\chi = g^{\nu\rho} \sigma_\nu^\beta \nabla_\beta \sigma_\rho^\chi. \tag{6.52}$$

The proposed limiting SDE is then

$$du_t = (\gamma^{-1} F)^h(u_t) dt - \frac{1}{2} (\gamma^{-2} Y)^h(u_t) dt + \gamma^{-1} (\pi(u_t)) H_{\sigma(u_t)}(u_t) \circ dW_t. \tag{6.53}$$

Next, we consider the case where both γ and σ are scalars.

Corollary 6.4. (A particle with scalar drag and noise) *Specializing further to $\gamma_\nu^\mu(x) = \gamma(x) \delta_\nu^\mu$ and $\sigma_\nu^\mu(x) = \sigma(x) \delta_\nu^\mu$ for some $\gamma, \sigma \in C^\infty(M)$ we obtain $S^v = 0$,*

$$S^h = -\frac{1}{2} (\gamma^{-2} \sigma \nabla \sigma)^h, \tag{6.54}$$

and hence the proposed limiting SDE is

$$du_t = (\gamma^{-1} F)^h(u_t) dt - \frac{1}{2} (\gamma^{-2} \sigma \nabla \sigma)^h(u_t) dt + (\gamma^{-1} \sigma) (\pi(u_t)) H(u_t) \circ dW_t. \tag{6.55}$$

In particular, if σ is a constant then the noise-induced drift vanishes.

Finally, we arrive at the case that leads to Brownian motion in the limit.

Corollary 6.5. (Brownian motion) *If $\gamma = \sigma$ are constant scalars and $F = 0$ then the proposed limiting SDE is*

$$du_t = H(u_t) \circ dW_t, \tag{6.56}$$

whose solution is the lift of a Brownian motion on M to the frame bundle [8].

Once we prove convergence in the next section, this last corollary will complete the objective set forth in the introduction, namely deriving Brownian motion on the manifold as the small mass limit of a noisy inertial system with drag.

6.2. Behavior Under Change of Frame

The transformation properties of the proposed limiting equation Eq. (6.24) under right multiplication by an orthogonal matrix are given by the following lemma.

Lemma 6.3. *Let $h \in O(\mathbb{R}^n)$ and u_t be the solution to Eq. (6.24) corresponding to the initial condition u_0 . Suppose σ is of the form Eq. (4.16). Then*

$$\tilde{u}_t = u_t h \tag{6.57}$$

is the solution to Eq. (6.24) corresponding to the initial condition $u_0 h$ with the Wiener process W_t replaced by the Wiener process $\tilde{W}_t = h^{-1} W_t$.

Proof. \tilde{u}_t is a semimartingale with initial condition u_0h . Right multiplication by h is a diffeomorphism, and hence by Lemma 4.1, \tilde{u}_t is a solution to

$$d\tilde{u}_t = ((R_h)_*(\gamma^{-1}F)^h + (R_h)_*S^h + (R_h)_*S^v)(\tilde{u}_t)dt + (R_h)_*(H_{\gamma^{-1}\sigma})(\tilde{u}_t) \circ dW_t. \tag{6.58}$$

By Lemma 3.5, horizontal lifts are invariant under right translation. The vertical term is also right invariant by the following computation:

$$\begin{aligned} ((R_h)_*S^v)(\tilde{u}_t) &= (R_h)_*(S^v(u_t)) \\ &= -\frac{1}{2}J^{\xi\nu}(u_t)(\gamma^{-1}(u_t))_\nu^\eta[(R_h)_*H_\eta, (R_h)_*H_\xi](\tilde{u}_t) \\ &= -\frac{1}{2}(h^{-1})_\xi^\beta J^{\xi\nu}(u_t)(\gamma^{-1}(u_t))_\nu^\eta (h^{-1})_\eta^\alpha [H_\alpha, H_\beta](\tilde{u}_t). \end{aligned} \tag{6.59}$$

Using the definition of $\gamma(u)$ along with Eq. (6.19), one can show that

$$(h^{-1})_\xi^\beta J^{\xi\nu}(u_t)(\gamma^{-1}(u_t))_\nu^\eta (h^{-1})_\eta^\alpha = J^{\beta\nu}(\tilde{u}_t)(\gamma^{-1}(\tilde{u}_t))_\nu^\alpha \tag{6.60}$$

and hence

$$((R_h)_*S^v)(\tilde{u}_t) = S^v(\tilde{u}_t). \tag{6.61}$$

The last term in Eq. (6.58) is

$$(R_h)_*(H_{\gamma^{-1}\sigma})(\tilde{u}_t) = H_{h^{-1}(\gamma^{-1}\sigma)(u_t)}(\tilde{u}_t) = H_{(\gamma^{-1}\sigma)(\tilde{u}_t)h^{-1}}(\tilde{u}_t). \tag{6.62}$$

Hence, we have

$$d\tilde{u}_t = (\gamma^{-1}F)^h(\tilde{u}_t) + S^h(\tilde{u}_t) + S^v(\tilde{u}_t) + (H_{\gamma^{-1}\sigma})(\tilde{u}_t) \circ d\tilde{W}_t \tag{6.63}$$

as claimed. □

We end this section by showing that if one is only interested in the statistics of the position process, then the vertical drift term, S^v , can be neglected.

Proposition 6.1. *Let u_t be the solution to the proposed limiting SDE, Eq. (6.24). Suppose σ is of the form Eq. (4.16). Then there exists an $O(\mathbb{R}^n)$ -valued semimartingale, h_t , with $h_0 = I$, the identity matrix, such that $\tilde{u}_t = u_t h_t$ is a solution to the limiting SDE, minus the vertical component of the drift, i.e., a solution to*

$$\begin{aligned} d\tilde{u}_t &= (\gamma^{-1}F)^h(\tilde{u}_t)dt + H_{(\gamma^{-1}\sigma)(\tilde{u}_t)}(\tilde{u}_t) \circ d\tilde{W}_t \\ &\quad - \frac{1}{2} \sum_\alpha (\gamma^{-1}(\tilde{u}_t))_\mu^\eta \sigma_\alpha^\mu(\tilde{u}_t) (\gamma^{-1}(\tilde{u}_t))_\nu^\xi H_\eta(\tilde{u}_t) [\sigma_\alpha^\nu] H_\xi(\tilde{u}_t) dt \\ &\quad - \frac{1}{2} ((\gamma^{-1}(\tilde{u}_t))_\mu^\eta J^{\mu\chi}(\tilde{u}_t) \gamma_\chi^\nu(\tilde{u}_t) - J^{\eta\nu}(\tilde{u}_t)) H_\eta(\tilde{u}_t) [(\gamma^{-1})_\nu^\xi] H_\xi(\tilde{u}_t) dt, \end{aligned} \tag{6.64}$$

where the Wiener process $\tilde{W}_t = \int_0^t h_s^{-1} dW_s$ is used in the Stratonovich integral.

In particular, since $u_t h_t$ and u_t have the same position process and uniqueness in law holds for an SDE on a compact manifold (as can be seen by employing a smooth embedding in some \mathbb{R}^l along with the corresponding result in [1]), the distribution of the position process is unchanged by the vertical component of the drift, even if one does not make a change to the Wiener process \tilde{W}_t .

Proof. Note that for any $O(\mathbb{R}^n)$ -valued semimartingale, h_t , the process $\tilde{W}_t = \int_0^t h_s^{-1} dW_s$ is a continuous \mathbb{R}^n -valued local martingale with quadratic covariation

$$[\tilde{W}^\alpha, \tilde{W}^\beta]_t = \int_0^t (h_s^{-1})^\alpha_\delta (h_s^{-1})^\beta_\eta \delta^{\delta\eta} ds = t\delta^{\alpha\beta}. \tag{6.65}$$

Hence, \tilde{W}_t is a Wiener process by Lévy’s theorem [1].

We will show more generally that given a vector field, $Y(x)$, on M , a $\binom{1}{1}$ -tensor field, $\tau(x)$, on M , and a vertical vector field, $V(u)$, on $F_O(M)$ (not to be confused with the forcing, Eq. (4.12)) one can go from a solution of the SDE

$$du_t = (Y)^h(u_t)dt + V(u_t)dt + H_{\tau(u_t)}(u_t) \circ dW_t \tag{6.66}$$

to a solution of

$$d\tilde{u}_t = (Y)^h(\tilde{u}_t)dt + H_{\tau(\tilde{u}_t)}(\tilde{u}_t) \circ d\tilde{W}_t \tag{6.67}$$

with the same initial condition in the manner described above.

Let $\phi : F_O(M) \times O(\mathbb{R}^n) \rightarrow F_O(M)$ be the right action and, for $u \in F_O(M)$, define $\phi_u(h) = \phi(u, h)$. These are both smooth maps. For each $A \in \mathfrak{o}(\mathbb{R}^n)$, the Lie algebra of $O(\mathbb{R}^n)$, we obtain a smooth vertical vector field V_A on $F_O(M)$ defined by

$$V_A(u) = (\phi_u)_* A. \tag{6.68}$$

If we let A_i be a basis for $\mathfrak{o}(\mathbb{R}^n)$ then $V_i(u) \equiv V_{A_i}(u)$ form a basis for the vertical subspace at each u [8]. Therefore, we can write $V(u) = V^i(u)V_i(u)$ for some smooth functions V^i on $F_O(M)$.

Let X_i be the smooth right-invariant vector fields on $O(\mathbb{R}^n)$ defined by $X_i(h) = (R_h)_* A_i$. Consider the SDE on the compact manifold $F_O(M) \times O(\mathbb{R}^n)$,

$$du_t = (Y)^h(u_t)dt + V(u_t)dt + H_{\tau(u_t)}(u_t) \circ dW_t, \tag{6.69}$$

$$dh_t = -V^i(u_t)X_i(h)dt. \tag{6.70}$$

By compactness, a unique solution, (u_t, h_t) , corresponding to the initial condition (u_0, I) exists for all $t \geq 0$. Note that the first component, u_t , is a solution to Eq. (6.66) with the initial condition u_0 .

Next we see how the vector fields of this SDE behave under the pushforward.

$$\begin{aligned} &\phi_*((Y)^h(u) + V(u), -V^i(u)X_i(h)) \\ &= (R_h)_*((Y)^h(u) + V(u)) - (\phi_u)_*(V^i(u)X_i(h)) \\ &= (Y)^h(uh) + (R_h)_*(V(u)) - V^i(u)(\phi_u \circ R_h)_* A_i \\ &= (Y)^h(uh) + (R_h)_* V(u) - V^i(u)(R_h)_* V_i(u) \\ &= (Y)^h(uh) \end{aligned} \tag{6.71}$$

and

$$\begin{aligned} \phi_*(H_{\tau(u)e_\alpha}(u), 0) &= (R_h)_* H_{\tau(u)e_\alpha}(u) \\ &= H_{h^{-1}\tau(u)e_\alpha}(uh) = (h^{-1})^\beta_\alpha H_{\tau(uh)e_\beta}(uh). \end{aligned} \tag{6.72}$$

Using Eqs. (6.71) and (6.72), we can derive the SDE satisfied by the semimartingale $\tilde{u}_t = u_t h_t = \phi(u_t, h_t)$. For any $f \in C^\infty(F_O(M))$, $\tilde{f} = f \circ \phi$ is a smooth function on $F_O(M) \times O(\mathbb{R}^n)$, hence

$$f(\tilde{u}_t) = \tilde{f}(u_0, I) + \int_0^t ((Y)^h(u_s) + V(u_s), -V^i(u_s)X_i(h_s))[\tilde{f}]ds + \int_0^t H_{\tau(u_s)}(u_s)[\tilde{f}] \circ dW_s \tag{6.73}$$

$$= f(u_0) + \int_0^t \phi_*((Y)^h(u_s) + V(u_s), -V^i(u_s)X_i(h_s))[f]ds + \int_0^t \phi_* H_{\tau(u_s)}(u_s)[f] \circ dW_s \tag{6.74}$$

$$= f(u_0) + \int_0^t (Y)^h(\tilde{u}_s)[f]ds + \int_0^t H_{\tau(\tilde{u}_s)e_\beta}(\tilde{u}_s)[f](h_s^{-1})^\beta_\alpha \circ dW_s^\alpha = f(u_0) + \int_0^t (Y)^h(\tilde{u}_s)[f]ds + \int_0^t H_{\tau(\tilde{u}_s)}(\tilde{u}_s)[f] \circ d\tilde{W}_s, \tag{6.75}$$

where we used the fact that h_t has locally bounded variation, and hence

$$\int_0^t h_s^{-1} \circ dW_s = \int_0^t h_s^{-1} dW_s = \tilde{W}_t. \tag{6.76}$$

Therefore, \tilde{u}_t solves the SDE (6.67) with the initial condition u_0 , as claimed. Applying this result to the proposed limiting SDE, Eq. (6.24), completes the proof. □

7. Existence of the Zero-Mass Limit

We are now in a position to prove convergence of the solutions of the SDE with mass, Eqs. (4.13)–(4.14), to the solution of the proposed limiting SDE, Eq. (6.24), as $m \rightarrow 0$. First we need a pair of lemmas that relate metric distance on $F_O(M)$ to smooth functions. These lemmas will allow us to prove convergence globally on the manifold $F_O(M)$, without explicitly patching together computations in local coordinates. Both lemmas make important use of compactness of the manifold in question.

Lemma 7.1. *Let (M, g) be a compact, connected Riemannian manifold and d be the metric on M induced by g . For any $f \in C^\infty(M)$ there exists a $C > 0$ such that*

$$|f(x) - f(y)| \leq Cd(x, y) \tag{7.1}$$

for all $x, y \in M$.

Proof. The result follows from writing

$$f(y) - f(x) = \int_0^1 \dot{\eta}_t[f]dt = \int_0^1 g_t(\dot{\eta}_t, \nabla f_t)dt \tag{7.2}$$

for any piecewise smooth curve η from x to y , taking the absolute value, and using the definition of d together with compactness of M . □

Lemma 7.2. *Let (M, g) be a compact, connected Riemannian manifold with metric d . For any $p > 0$ there exist finitely many smooth functions f_i such that*

$$d(x, y)^p \leq \sum_i |f_i(x) - f_i(y)|^p. \tag{7.3}$$

Proof. For each $x \in M$ there exists a coordinate chart (U_x, ϕ_x) about x and a constant C_x such that for $y, z \in U_x$ we have $d(y, z) \leq C_x \|\phi_x(y) - \phi_x(z)\|$ where the right-hand side is the Euclidean norm (see [35, 36]). Shrinking the charts if necessary, we can assume that ϕ_x extends smoothly to all of M .

Take another open set, V_x , containing x , with V_x having compact closure in U_x . By compactness of M there exists a finite subcover $V_i \equiv V_{x_i}$ and there also exists $R > 0$ such that $d(y, z) \leq R$ for all $y, z \in M$. For each i let ψ_i be a smooth function equal to R on V_i with compact support in $U_i \equiv U_{x_i}$ and define $C_i \equiv C_{x_i}, \phi_i \equiv \phi_{x_i}$.

Take $y, z \in M$. $y \in V_i$ for some i . If $z \in U_i$ then

$$d(y, z)^2 \leq C_i^2 \|\phi_i(y) - \phi_i(z)\|^2 = \sum_j (C_i \phi_i^j(y) - C_i \phi_i^j(z))^2. \tag{7.4}$$

Otherwise, $z \notin U_i$, hence

$$d(y, z)^2 \leq R^2 = (\psi_i(y) - \psi_i(z))^2. \tag{7.5}$$

So taking the collection of functions $C_i \phi_i^j, \psi_i$ gives the result for $p = 2$.

For $p > 0$, if we let N be the number of functions we obtained when $p = 2$ then raising the $p = 2$ result to the power $p/2$ gives

$$\begin{aligned} d(x, y)^p &\leq \left(\sum_i |f_i(x) - f_i(y)|^2 \right)^{p/2} \leq N^{p/2} \sum_i |f_i(x) - f_i(y)|^p \\ &= \sum_i |g_i(x) - g_i(y)|^p \end{aligned} \tag{7.6}$$

where $g_i = N^{1/2} f_i$. □

With these two lemmas, we can now prove the convergence result.

Theorem 7.1. *Fix $(u_0, v_0) \in N$. Let (u_t^m, v_t^m) be the unique solution to Eqs. (4.13)–(4.14) with initial condition (u_0, v_0) and let u_t be the unique solution to Eq. (6.24) with initial condition u_0 . Fix $T > 0$ and a Riemannian metric tensor field on $F_O(M)$. Let d be the associated metric on the connected component of $F_O(M)$ that contains u_0 . Then for any $q > 0$ and any $0 < \kappa < q/2$ we have*

$$E \left[\sup_{t \in [0, T]} d(u_t^m, u_t)^q \right] = O(m^\kappa) \quad \text{as } m \rightarrow 0. \tag{7.7}$$

We emphasize that this result is heavily reliant on Assumption 1; the existence of a positive lower bound on the eigenvalues of the symmetric part of the damping tensor γ .

Proof. As discussed in Lemma 4.4, a unique solution (u_t^m, v_t^m) is defined for all $t \geq 0$. Since Eq. (6.24) is an SDE on the compact manifold $F_O(M)$, it also has a unique solution, u_t , defined for all $t \geq 0$. Both u_t^m and u_t are continuous, so they stay in the same connected component as u_0 .

For $f \in C^\infty(F_O(M))$, define

$$M^f(u) = H_{(\gamma^{-1}F)(u)}(u)[f] + J^{\beta\alpha}(u)K_{\beta\alpha}^f(u), \quad Q_\eta^f(u) = H_{(\gamma^{-1}\sigma)(u)e_\eta}(u)[f] \tag{7.8}$$

where K^f and J were defined in Eqs. (6.7) and (6.19), respectively. M^f and Q_η^f are smooth functions of u .

Remark 3. We again emphasize that the indices appearing in Eq. (7.8) and in the subsequent computations represent the components in the standard basis of functions on $F_O(M)$ that are valued in the product of some \mathbb{R} 's. Though we employ the summation convention, these expressions do not represent (contractions of) tensors on M or $F_O(M)$. In particular, the following convergence proof is global in nature and does not directly utilize any computations in local coordinates on the base manifold or its frame bundle.

Using Eqs. (6.21) and (6.25) and rearranging we obtain

$$f(u_t^m) - f(u_t) = \int_0^t (M^f(u_s^m) - M^f(u_s)) ds + \int_0^t (Q_\eta^f(u_s^m) - Q_\eta^f(u_s)) dW_s^\eta + R^f(t), \tag{7.9}$$

where we have grouped the momentum-dependent terms in the quantity

$$\begin{aligned} R^f(t) = & - \int_0^t K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) d((p_s^m)^\nu (p_s^m)^\mu) \\ & + \int_0^t K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\nu F^\mu(u_s^m) + (p_s^m)^\mu F^\nu(u_s^m)) ds \\ & + \int_0^t K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\mu \sigma_\eta^\nu(u_s^m) + (p_s^m)^\nu \sigma_\eta^\mu(u_s^m)) dW_s^\eta \\ & - H_{\gamma^{-1}(u_t^m)p_t^m}(u_t^m)[f] + H_{\gamma^{-1}(u_0)p_0}(u_0)[f]. \end{aligned} \tag{7.10}$$

For now, restrict to $q = 2p$ with $p > 1$. For any $t \leq T$,

$$\begin{aligned} E \left[\sup_{0 \leq s \leq t} |f(u_s^m) - f(u_s)|^{2p} \right] \leq & 3^{2p-1} \left(E \left[\left(\int_0^t |M^f(u_s^m) - M^f(u_s)| ds \right)^{2p} \right] \right. \\ & + E \left[\sup_{0 \leq s \leq t} \left| \int_0^s Q_\eta^f(u_r^m) - Q_\eta^f(u_r) dW_r^\eta \right|^{2p} \right] \\ & \left. + E \left[\sup_{0 \leq s \leq t} |R^f(s)|^{2p} \right] \right). \end{aligned} \tag{7.11}$$

Applying the Hölder’s inequality with exponents $2p$ and $2p/(2p - 1)$ to the first term and the Burkholder–Davis–Gundy inequality to the second we get

$$\begin{aligned}
 E \left[\sup_{0 \leq s \leq t} |f(u_s^m) - f(u_s)|^{2p} \right] &\leq 3^{2p-1} \left(T^{2p-1} E \left[\int_0^t |M^f(u_s^m) - M^f(u_s)|^{2p} ds \right] \right. \\
 &\quad \left. + CE \left[\left(\int_0^t \sum_{\eta} |Q_{\eta}^f(u_s^m) - Q_{\eta}^f(u_s)|^2 ds \right)^p \right] \right. \\
 &\quad \left. + E \left[\sup_{0 \leq s \leq t} |R^f(s)|^{2p} \right] \right). \tag{7.12}
 \end{aligned}$$

We have assumed $p > 1$, so we can use Hölder’s inequality with indices p and $p/(p - 1)$ on the second term.

$$\begin{aligned}
 E \left[\left(\int_0^t \sum_{\eta} |Q_{\eta}^f(u_s^m) - Q_{\eta}^f(u_s)|^2 ds \right)^p \right] \\
 \leq T^{p-1} E \left[\int_0^t \left(\sum_{\eta} |Q_{\eta}^f(u_s^m) - Q_{\eta}^f(u_s)|^2 \right)^p ds \right]. \tag{7.13}
 \end{aligned}$$

Therefore,

$$\begin{aligned}
 E \left[\sup_{0 \leq s \leq t} |f(u_s^m) - f(u_s)|^{2p} \right] &\leq 3^{2p-1} \left(T^{2p-1} E \left[\int_0^t |M^f(u_s^m) - M^f(u_s)|^{2p} ds \right] \right. \\
 &\quad \left. + C(kT)^{p-1} \sum_{\eta} E \left[\int_0^t |Q_{\eta}^f(u_s^m) - Q_{\eta}^f(u_s)|^{2p} ds \right] \right. \\
 &\quad \left. + E \left[\sup_{0 \leq s \leq t} |R^f(s)|^{2p} \right] \right) \tag{7.14}
 \end{aligned}$$

where k is the number of terms in the sum over η .

The integrands of the terms involving M^f and Q_{η}^f are of the form $|g(u_s^m) - g(u_s)|^{2p}$ for some smooth functions g , depending on f . Therefore, for a fixed f , by Lemma 7.1 there exists $C_f > 0$ such that

$$\begin{aligned}
 E \left[\sup_{0 \leq s \leq t} |f(u_s^m) - f(u_s)|^{2p} \right] &\leq 3^{2p-1} (T^{2p-1} + Ck^p T^{p-1}) C_f \\
 &\quad \times \int_0^t E [d(u_s^m, u_s)^{2p}] ds + 3^{2p-1} E \left[\sup_{0 \leq s \leq t} |R^f(s)|^{2p} \right]. \tag{7.15}
 \end{aligned}$$

$F_O(M)$ is compact and we have equipped it with a Riemannian metric tensor. Therefore, by Lemma 7.2, there exist finitely many smooth functions, f_i , such that for any u_1, u_2 in the connected component of $F_O(M)$ containing u_0 we have

$$d(u_1, u_2)^{2p} \leq \sum_i |f_i(u_1) - f_i(u_2)|^{2p}. \tag{7.16}$$

Hence, applying Eq. (7.15) to each of the functions f_i , we obtain

$$\begin{aligned}
 E \left[\sup_{0 \leq s \leq t} d(u_s^m, u_s)^{2p} \right] &\leq \sum_i E \left[\sup_{0 \leq s \leq t} |f_i(u_s^m) - f_i(u_s)|^{2p} \right] \\
 &\leq \sum_i C_i \int_0^t E [d(u_s^m, u_s)^{2p}] ds \\
 &\quad + 3^{2p-1} \sum_i E \left[\sup_{0 \leq s \leq t} |R^{f_i}(s)|^{2p} \right] \tag{7.17}
 \end{aligned}$$

for some constants C_i . Therefore, there exists a $C > 0$ such that

$$\begin{aligned}
 E \left[\sup_{0 \leq s \leq t} d(u_s^m, u_s)^{2p} \right] &\leq C \int_0^t E \left[\sup_{0 \leq r \leq s} d(u_r^m, u_r)^{2p} \right] ds \\
 &\quad + 3^{2p-1} \sum_i E \left[\sup_{0 \leq s \leq T} |R^{f_i}(s)|^{2p} \right] \tag{7.18}
 \end{aligned}$$

for all $0 \leq t \leq T$. We apply Gronwall’s inequality to obtain

$$E \left[\sup_{0 \leq s \leq T} d(u_s^m, u_s)^{2p} \right] \leq 3^{2p-1} \sum_i E \left[\sup_{0 \leq s \leq T} |R^{f_i}(s)|^{2p} \right] e^{CT}. \tag{7.19}$$

Fix $0 < \kappa < p$. The sum in Eq. (7.19) has finitely many terms, so, if we can prove that

$$E \left[\sup_{0 \leq s \leq T} |R^f(s)|^{2p} \right] = O(m^\kappa) \quad \text{as } m \rightarrow 0 \tag{7.20}$$

for every smooth f , then the claim will follow for $q = 2p, p > 1$.

Fix $f \in C^\infty(F_O(M))$ and compute

$$\begin{aligned}
 &E \left[\sup_{0 \leq t \leq T} |R^f(t)|^{2p} \right] \\
 &\leq 5^{2p-1} \left(E \left[\sup_{0 \leq t \leq T} \left| \int_0^t K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) d((p_s^m)^\nu (p_s^m)^\mu) \right|^{2p} \right] \right. \\
 &\quad + E \left[\left(\int_0^T |K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\nu F^\mu(u_s^m) + (p_s^m)^\mu F^\nu(u_s^m))| ds \right)^{2p} \right] \\
 &\quad + E \left[\sup_{0 \leq t \leq T} \left| \int_0^t K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\nu \sigma_\eta^\mu(u_s^m) + (p_s^m)^\mu \sigma_\eta^\nu(u_s^m)) dW_s^\eta \right|^{2p} \right] \\
 &\quad \left. + E \left[\sup_{0 \leq t \leq T} |H_{\gamma^{-1}(u_t^m) p_t^m}(u_t^m)[f]|^{2p} \right] + E \left[|H_{\gamma^{-1}(u_0) p_0}(u_0)[f]|^{2p} \right] \right). \tag{7.21}
 \end{aligned}$$

We will consider each term individually. Proposition 5.3 implies that

$$E \left[\sup_{0 \leq t \leq T} \left| \int_0^t K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) d((p_s^m)^\nu (p_s^m)^\mu) \right|^{2p} \right] = O(m^p). \tag{7.22}$$

Using Hölder’s inequality, boundedness of continuous functions on the compact manifold $F_O(M)$, and Proposition 5.2, the second term can be bounded as follows:

$$\begin{aligned}
 E & \left[\left(\int_0^T |K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\nu F^\mu(u_s^m) + (p_s^m)^\mu F^\nu(u_s^m))| ds \right)^{2p} \right] \\
 & \leq T^{2p-1} \int_0^T E \left[\left| \left((K_{\beta\alpha}^f G_{\nu\mu}^{\beta\alpha} F^\nu)(u_s^m) + (K_{\beta\alpha}^f G_{\mu\nu}^{\beta\alpha} F^\nu)(u_s^m) \right) (p_s^m)^\mu \right|^{2p} \right] ds \\
 & \leq CT^{2p} \sup_{0 \leq s \leq T} E [\|p_s^m\|^{2p}] = O(m^p). \tag{7.23}
 \end{aligned}$$

The third term can be bounded using the Burkholder–Davis–Gundy inequality, similarly to the way the second term in Eq. (7.14) was estimated, together with Proposition 5.2.

$$\begin{aligned}
 E & \left[\sup_{0 \leq t \leq T} \left| \int_0^t K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) ((p_s^m)^\nu \sigma_\eta^\mu(u_s^m) + (p_s^m)^\mu \sigma_\eta^\nu(u_s^m)) dW_s^\eta \right|^{2p} \right] \\
 & \leq C(kT)^{p-1} \sum_\eta E \left[\int_0^T \left| \left(K_{\beta\alpha}^f(u_s^m) G_{\nu\mu}^{\beta\alpha}(u_s^m) \sigma_\eta^\nu(u_s^m) \right. \right. \right. \\
 & \quad \left. \left. \left. + K_{\beta\alpha}^f(u_s^m) G_{\mu\nu}^{\beta\alpha}(u_s^m) \sigma_\eta^\mu(u_s^m) \right) (p_s^m)^\mu \right|^{2p} ds \right] \tag{7.24}
 \end{aligned}$$

$$\leq \tilde{C}T^{p-1} \int_0^T E [\|p_s^m\|^{2p}] ds \leq \tilde{C}T^p \sup_{0 \leq s \leq T} E [\|p_s^m\|^{2p}] = O(m^p). \tag{7.25}$$

Compactness of $F_O(M)$ and Proposition 5.1 imply a bound on the fourth term:

$$\begin{aligned}
 E \left[\sup_{0 \leq t \leq T} |H_{\gamma^{-1}(u_t^m)p_t^m}(u_t^m)[f]|^{2p} \right] & = E \left[\sup_{0 \leq t \leq T} |H_{\gamma^{-1}(u_t^m)e_\nu}(u_t^m)[f](p_t^m)^\nu|^{2p} \right] \\
 & \leq CE \left[\sup_{0 \leq t \leq T} \|p_t^m\|^{2p} \right] = O(m^\kappa). \tag{7.26}
 \end{aligned}$$

Finally, the last term is

$$E[|H_{\gamma^{-1}(u_0)p_0}(u_0)[f]|^{2p}] = m^{2p} E[|H_{\gamma^{-1}(u_0)v_0}(u_0)[f]|^{2p}] = O(m^{2p}). \tag{7.27}$$

This completes the proof for $q = 2p, p > 1$. Similarly to Proposition 5.2, an application of Hölder’s inequality gives the result for all $q > 0$. \square

As a corollary, we get convergence in probability and in law on compact time intervals.

Corollary 7.1. *For any $T > 0$, the mass-dependent process restricted to the compact time interval $[0, T], u^m|_{[0, T]}$, converges in probability (and hence also in law) to $u|_{[0, T]}$ as $m \rightarrow 0$.*

Convergence in probability on the path space $C([0, T], F_O^i(M))$, where $F_O^i(M)$ is a connected component of $F_O(M)$, is defined through the metric

$$d_T(u, \tau) = \sup_{t \in [0, T]} d(u(t), \tau(t)) \quad (7.28)$$

where d is the metric on $F_O^i(M)$ induced by a choice of Riemannian metric tensor on $F_O(M)$.

Acknowledgments

The authors are grateful to K. Gawędzki, P. Grzegorzcyk, X.-M. Li, and A. McDaniel for stimulating discussions. We thank an editor of AHP for pointing out several references. The research of JW was supported in part by US NSF Grant DMS-131271. A part of his work on this article was supported by US NSF Grant DMS-1440140 while he was in residence at the Mathematical Sciences Research Institute in Berkeley during the Fall 2015 semester.

References

- [1] Karatzas, I., Shreve, S.: *Brownian Motion and Stochastic Calculus*. Graduate Texts in Mathematics. Springer, New York (2014)
- [2] Nelson, E.: *Dynamical Theories of Brownian Motion*, Mathematical Notes. Princeton University Press, Princeton (1967)
- [3] Casuso, I., Khao, J., Chami, M., Paul-Gilloteaux, P., Husain, M., Duneau, J.-P., Stahlberg, H., Sturgis, J.N., Scheuring, S.: Characterization of the motion of membrane proteins using high-speed atomic force microscopy. *Nat. Nanotechnol.* **7**(8), 525–529 (2012)
- [4] Kärger, J., Ruthven, D., Theodorou, D.: *Diffusion in Nanoporous Materials*. Wiley, New York (2012)
- [5] Barkai, E., Garini, Y., Metzler, R.: Strange kinetics of single molecules in living cells. *Phys. Today* **65**(8), 29 (2012)
- [6] Manzo, C., Torreno-Pina, J.A., Massignan, P., Lapeyre, G.J., Lewenstein, M., Garcia Parajo, M.F.: Weak ergodicity breaking of receptor motion in living cells stemming from random diffusivity. *Phys. Rev. X* **5**, 011021 (2015)
- [7] Ramaswamy, S.: The mechanics and statistics of active matter. *Annu. Rev. Condens. Matter Phys.* **1**, 323–345 (2010)
- [8] Hsu, E.: *Stochastic Analysis on Manifolds*, Contemporary Mathematics. American Mathematical Society, Providence (2002)
- [9] Stroock, D.W.: *An Introduction to the Analysis of Paths on a Riemannian Manifold* No. 74. American Mathematical Society, Providence (2005)
- [10] Van Kampen, N.: Brownian motion on a manifold. *J. Stat. Phys.* **44**(1–2), 1–24 (1986)
- [11] Kubo, R., Toda, M., Hashitsume, N.: *Statistical Physics II: Nonequilibrium Statistical Mechanics*. Springer Series in Solid-State Sciences. Springer, Berlin (2012)
- [12] Polettini, M.: Generally covariant state-dependent diffusion. *J. Stat. Mech. Theory Exp.* **2013**(07), P07005 (2013)

- [13] Hottovy, S., McDaniel, A., Volpe, G., Wehr, J.: The smoluchowski-kramers limit of stochastic differential equations with arbitrary state-dependent friction. *Commun. Math. Phys.* **336**(3), 1259–1283 (2014)
- [14] Smoluchowski, M.V.: Drei vortrage uber diffusion, brownsche bewegung und koagulation von kolloidteilchen. *Z. Phys.* **17**, 557–585 (1916)
- [15] Pavliotis, G.A., Stuart, A.M.: White noise limits for inertial particles in a random field. *Multiscale Model. Simul.* **1**(4), 527–553 (2003)
- [16] Chevalier, C., Debbasch, F.: Relativistic diffusions: a unifying approach. *J. Math. Phys.* **49**(4), 383 (2008)
- [17] Bailleul, I.: A stochastic approach to relativistic diffusions. *Annales de l’Institut Henri Poincaré (B)* **46**, 760–795 (2010)
- [18] Pinsky, M.A.: Isotropic transport process on a Riemannian manifold. *Trans. Am. Math. Soc.* **218**, 353–360 (1976)
- [19] Pinsky, M.A.: Homogenization in stochastic differential geometry. *Publ. Res. Inst. Math. Sci.* **17**(1), 235–244 (1981)
- [20] JØrgensen, E.: Construction of the brownian motion and the ornstein-uhlenbeck process in a riemannian manifold on basis of the gangolli-mc.kean injection scheme. *uZeitschrift fr Wahrscheinlichkeitstheorie Und Verwandte Gebiete.* **44**(1), 71–87 (1978)
- [21] Dowell, R.M.: Differentiable approximations to Brownian motion on manifolds, Ph.D. thesis. University of Warwick (1980)
- [22] Li, X.-M.: Random perturbation to the geodesic equation. *Ann. Probab.* **44**(1), 544–566 (2016)
- [23] Angst, J., Bailleul, I., Tardif, C.: Kinetic brownian motion on Riemannian manifolds. [arXiv:1501.03679](https://arxiv.org/abs/1501.03679), (2015, arXiv preprint)
- [24] Bismut, J.-M.: The hypoelliptic laplacian on the cotangent bundle. *J. Am. Math. Soc.* **18**(2), 379–476 (2005)
- [25] Bismut, J.-M.: Hypoelliptic laplacian and probability. *J. Math. Soc. Japan* **67**(4), 1317–1357 (2015)
- [26] Herzog, D.P., Hottovy, S., Volpe, G.: The small-mass limit for Langevin dynamics with unbounded coefficients and positive friction. [arXiv:1510.04187](https://arxiv.org/abs/1510.04187) (2015, arXiv preprint)
- [27] Seifert, U.: Stochastic thermodynamics, fluctuation theorems and molecular machines. *Rep. Prog. Phys.* **75**(12), 126001 (2012)
- [28] Kobayashi, S., Nomizu, K.: *Foundations of Differential Geometry Set.* Wiley Classics Library, Wiley, New York (2009)
- [29] Teschl, G.: *Ordinary Differential Equations and Dynamical Systems.* Graduate Studies in Mathematics. American Mathematical Society, Providence (2012)
- [30] Kedem, G.: A posteriori error bounds for two-point boundary value problems. *SIAM J. Numer. Anal.* **18**(3), 431–448 (1981)
- [31] Freidlin, M.: Some remarks on the Smoluchowski–Kramers approximation. *J. Stat. Phys.* **117**(3), 617–634 (2004)
- [32] Ortega, J.: *Matrix Theory: A Second Course.* University Series in Mathematics. Springer US (2013)
- [33] Wilcox, R.M.: Exponential operators and parameter differentiation in quantum physics. *J. Math. Phys.* **8**(4), 962 (1967)

- [34] Volpe, G., Wehr, J.: Effective drifts in dynamical systems with multiplicative noise: A review of recent progress. *Rep. Prog. Phys.* **79**(5), 053901 (2016)
- [35] Lee, J.: *Introduction to Smooth Manifolds*. Graduate Texts in Mathematics. Springer, New York (2013)
- [36] Lee, J.: *Riemannian Manifolds: An Introduction to Curvature*. Graduate Texts in Mathematics. Springer, New York (2006)

Jeremiah Birrell
Department of Mathematics
University of Arizona
Tucson, AZ 85721
USA
e-mail: jbirrell@math.arizona.edu

Scott Hottovy
Department of Mathematics
University of Wisconsin-Madison
Madison, WI 53706
USA
e-mail: shottovy@math.wisc.edu

Giovanni Volpe
Soft Matter Lab
Department of Physics
Bilkent University
Ankara 06800
Turkey

and

UNAM, National Nanotechnology Research Center
Bilkent University
Ankara 06800
Turkey
e-mail: giovanni.volpe@fen.bilkent.edu.tr

Jan Wehr
Department of Mathematics
University of Arizona
Tucson, AZ 85721
USA
e-mail: wehr@math.arizona.edu

Communicated by Christian Maes.

Received: April 1, 2016.

Accepted: June 15, 2016.