

ON PLANE SEXTICS WITH DOUBLE SINGULAR POINTS

ALEX DEGTYAREV

We compute the fundamental groups of five maximizing sextics with double singular points only; in four cases, the groups are as expected. The approach used would apply to other sextics as well, given their equations.

1. Introduction

The fundamental group $\pi_1 := \pi_1(\mathbb{P}^2 \setminus D)$ of a plane curve $D \subset \mathbb{P}^2$, introduced by O. Zariski [1929], is an important topological invariant of the curve. Apart from distinguishing the connected components of the equisingular moduli spaces, this group can be used as a seemingly inexpensive way of studying algebraic surfaces, the curve serving as the branch locus of a projection of the surface onto \mathbb{P}^2 .

At present, the fundamental groups of all curves of degree up to five are known, and the computation of the groups of irreducible curves of degree six (sextics) is close to its completion; see [Degtyarev 2012] for the principal statements and further references. In higher degrees, little is known: there are a few general theorems, usually bounding the complexity of the group of a curve with sufficiently “moderate” singularities, and a number of sporadic examples scattered in the literature. For further details on this fascinating subject, we refer the reader to the recent surveys [Artal-Bartolo et al. 2008; Libgober 2007a; 2007b].

1A. Principal results. If a sextic $D \subset \mathbb{P}^2$ has a singular point P of multiplicity three or higher, then, projecting from this point, we obtain a trigonal (or, even better, bi- or monogonal) curve in a Hirzebruch surface; see Section 3A. By means of the so-called *dessins d’enfants*, such curves and their topology can be studied in purely combinatorial terms, as certain graphs in the plane. The classification of such curves and the computation of their fundamental groups were completed in [Degtyarev 2012]. If all singular points are double, the best that one can obtain is a tetragonal curve, which is a much more complicated object. (A reduction of tetragonal curves to trigonal curves in the presence of a section is discussed in Section 3B; see Remark 3.6. It is the extra section that makes the problem difficult.) At present, I do not know how the group of a tetragonal curve can be computed

MSC2000: primary 14H45; secondary 14H30, 14H50.

Keywords: plane sextic, torus type, fundamental group, tetragonal curve.

unless the curve is real and its defining equation is known (and, even then, the approach suggested in the paper may still fail; cf. [Remark 2.1](#)).

There is a special class of irreducible sextics, the so-called \mathbb{D}_{2n} -*sextics* and, in particular, sextics of *torus type* (see [Section 2A](#) for the precise definitions), for which the fundamental group is nonabelian for some simple homological reasons; see [[Degtyarev 2008](#)]. (The fact that a sextic is of torus type is usually indicated by the presence of a pair of parentheses in the notation; their precise meaning is explained in [Section 2A](#).) On the other hand, thanks to the special structures and symmetries of these curves, their explicit equations are known; see [[Degtyarev 2009b](#); [Degtyarev and Oka 2009](#); [Oka and Pho 2002](#)]. In this paper, we almost complete the computation of the fundamental groups of \mathbb{D}_{2n} -sextics (with one pair of complex conjugate sextics of torus type left). Our principal results can be stated as follows.

Theorem 1.1. *The fundamental group of the \mathbb{D}_{14} -special sextic with the set of singularities $3A_6 \oplus A_1$, [line 37](#) in [Table 1](#), is $\mathbb{Z}_3 \times \mathbb{D}_{14}$.*

Theorem 1.2. *The fundamental groups of the irreducible sextics of torus type with the sets of singularities $(A_{14} \oplus A_2) \oplus A_3$, [line 8](#), $(A_{14} \oplus A_2) \oplus A_2 \oplus A_1$, [line 9](#), and $(A_{11} \oplus 2A_2) \oplus A_4$, [line 17](#) in [Table 1](#), are isomorphic to $\Gamma := \mathbb{Z}_2 * \mathbb{Z}_3$. The group of the curve with the set of singularities $(A_8 \oplus 3A_2) \oplus A_4 \oplus A_1$, [line 33](#), is*

$$(1.3) \quad \pi_1 = \langle \alpha_2, \alpha_3, \alpha_4 \mid [\alpha_3, \alpha_4] = \{\alpha_2, \alpha_3\}_3 = \{\alpha_2, \alpha_4\}_9 = 1, \\ \alpha_4 \alpha_2 \alpha_3^{-1} \alpha_4 \alpha_2 \alpha_4 (\alpha_4 \alpha_2)^{-2} \alpha_3 = (\alpha_2 \alpha_4)^2 \alpha_3^{-1} \alpha_2 \alpha_4 \alpha_3 \alpha_2 \rangle,$$

where $\{\alpha, \beta\}_{2k+1} := (\alpha\beta)^k \alpha (\alpha\beta)^{-k} \beta^{-1}$.

[Theorem 1.1](#) is proved in [Section 4C](#), and [Theorem 1.2](#) is proved in [Sections 4E–4H](#), one curve at a time. I do not know whether the last group (1.3) is isomorphic to Γ : all “computable” invariants seem to coincide (see [Remark 4.7](#)), but the presentations obtained resist all simplification attempt. The quotient of (1.3) by the extra relation $\{\alpha_2, \alpha_4\}_3 = 1$ is Γ .

The next proposition is proved in [Section 4I](#). (The perturbation $3A_6 \oplus A_1 \rightarrow 3A_6$ excluded in the statement results in a \mathbb{D}_{14} -special sextic and the fundamental group equals $\mathbb{Z}_3 \times \mathbb{D}_{14}$; see [[Degtyarev and Oka 2009](#)].)

Proposition 1.4. *Let D' be a nontrivial perturbation of a sextic as in [Theorem 1.1](#) or [1.2](#). Unless the set of singularities of D' is $3A_6$, the group $\pi_1(\mathbb{P}^2 \setminus D')$ is Γ or \mathbb{Z}_6 , depending on whether D' is, or, respectively, is not, of torus type.*

With [Theorem 1.1](#) in mind, the fundamental groups of all \mathbb{D}_{2n} -special sextics, $n \geq 5$, are known; see [[Degtyarev 2012](#)]. Modulo the feasible conjecture that any sextic of torus type degenerates to a maximizing one, the only such sextic whose group remains unknown is $(A_8 \oplus A_5 \oplus A_2) \oplus A_4$, [line 32](#) in [Table 1](#). (This

conjecture has been proved, and *all* groups except the one just mentioned are indeed known; details will appear elsewhere.) Most of these groups are isomorphic to Γ ; see [Degtyarev 2012] for details and further references.

I would like to mention an alternative approach (see [Artal Bartolo et al. 2002]) reducing a plane sextic with large Milnor number to a trigonal curve equipped with a number of sections, all but one splitting in the covering elliptic surface. It was used in [Artal Bartolo et al. 2002] to handle the curves in lines 1–6 in Table 1. This approach is also used in a forthcoming paper to produce the defining equations of most sextics listed in Table 1; then, the fundamental groups of most *real* ones can be computed using Theorem 3.16. All groups that could be found are abelian. Together with the classification of sextics, which is also almost completed, this fact implies that, with very few exceptions, the fundamental group of a nonspecial irreducible simple sextic is abelian.

1B. Idea of the proof (see Section 4A for more details). We use the classical Zariski–van Kampen method (see Theorem 3.16), expressing the fundamental group of a curve in terms of its braid monodromy with respect to an appropriate pencil of lines. The curves and pencils considered are *real*, and the braid monodromy in a neighborhood of the real part of the pencil is computed in terms of the real part of the curve. (This approach originates in topology of real algebraic curves; historically, it goes back to Viro, Fiedler, Kharlamov, Rokhlin, and Klein.) Our main contribution is the description of the monodromy along a real segment where all four branches of the curve are nonreal; see Proposition 3.12. Besides, the curves are not required to be *strongly real*; i.e., nonreal singular fibers are allowed. Hence, we follow [Orevkov 1999] and attempt to extract information about such nonreal fibers from the real part of the curve. The outcome is Theorem 3.16, which gives us an “upper bound” on the fundamental group in question. The applicability issues and a few other common tricks are discussed in Section 4A.

1C. Contents of the paper. In Section 2, we introduce the terminology related to plane sextics, list the sextics that are still to be investigated, and discuss briefly the few known results. In Section 3, we outline an approach to the (partial) computation of the braid monodromy of a real tetragonal curve and state an appropriate version of the Zariski–van Kampen theorem. Finally, in Section 4 the results of Section 3 and known equations are used to prove Theorems 1.1 and 1.2 and Proposition 1.4.

1D. Conventions. All group actions are *right*. Given a right action $X \times G \rightarrow X$ and a pair of elements $x \in X$, $g \in G$, the image of (x, g) is denoted by $x \uparrow g \in X$. The same postfix notation and multiplication convention is often used for maps: it is under this convention that the monodromy $\pi_1(\text{base}) \rightarrow \text{Aut}(\text{fiber})$ of a locally trivial fibration is a homomorphism rather than an antihomomorphism.

The assignment symbol $:=$ is used as a shortcut for “is defined as”.

We use the conventional symbol \square to mark the ends of the proofs. Some statements are marked with \triangleleft , meaning that the proof has already been explained (for example, most corollaries).

2. Preliminaries

2A. *Special classes of sextics.* A plane sextic $D \in \mathbb{P}^2$ is called *simple* if all its singularities are simple, i.e., those of type $A-D-E$. The total Milnor number μ of a simple sextic D does not exceed 19; see [Persson 1985]; if $\mu = 19$, then D is called *maximizing*. Maximizing sextics are always defined over algebraic number fields and their moduli spaces are discrete: two such sextics are equisingular deformation equivalent if and only if they are related by a projective transformation of \mathbb{P}^2 .

A sextic D is said to be of *torus type* if its equation can be represented in the form $f_2^3 + f_3^2 = 0$, where f_2 and f_3 are some polynomials of degree 2 and 3, respectively. The points of intersection of the conic $\{f_2 = 0\}$ and cubic $\{f_3 = 0\}$ are always singular for D . These singular points play a very special rôle; they are called the *inner* singularities (with respect to the given torus structure). For the vast majority of curves, a torus structure is unique, and in this case it is common to parenthesize the inner singularities in the notation.

An irreducible sextic D is called \mathbb{D}_{2n} -*special* if its fundamental group $\pi_1(\mathbb{P}^2 \setminus D)$ admits a dihedral quotient $\mathbb{D}_{2n} := \mathbb{Z}_n \rtimes \mathbb{Z}_2$. According to [Degtyarev 2008], only \mathbb{D}_6 -, \mathbb{D}_{10} -, and \mathbb{D}_{14} -special sextics exist, and an irreducible sextic is of torus type if and only if it is \mathbb{D}_6 -special. (In particular, torus type is a topological property.)

Any sextic D of torus type is a degeneration of Zariski’s six-cuspidal sextic, which is obtained from a generic pair (f_2, f_3) . It follows that the fundamental group of D factors to the modular group $\Gamma := SL(2, \mathbb{Z}) = \mathbb{Z}_2 * \mathbb{Z}_3 = \mathbb{B}_3 / (\sigma_1 \sigma_2 \sigma_1)^2$; see [Zariski 1929]; in particular, this group is infinite. Conjecturally, the fundamental groups of all other irreducible simple sextics are finite.

2B. *Sextics to be considered.* It is expected that, with few explicit exceptions (e.g., $9A_2$), any simple sextic degenerates to a maximizing one. (The proof of this conjecture, which relies upon the theory of $K3$ -surfaces, is currently a work in progress. In fact, most curves degenerate to one of those whose groups are already known.) Hence, it is essential to compute the fundamental groups of the maximizing sextics; the others would follow. The groups of all irreducible sextics with a singular point of multiplicity three or higher are known (see [Degtyarev 2012] for a summary of the results), and those with A type singularities only are still to be investigated.

A list of irreducible maximizing sextics with A type singular points only can be compiled using the results of [Yang 1996] (a list of the sets of singularities realized

#	Singularities	(r, c)	Equation, π_1 , remarks
1.	A_{19}	$(2, 0)$	$\pi_1 = \mathbb{Z}_6$, see [Artal Bartolo et al. 2002]
2.	$A_{18} \oplus A_1$	$(1, 1)$	$\pi_1 = \mathbb{Z}_6$, see [ibid.]
3.	$(A_{17} \oplus A_2)$	$(1, 0)^*$	$\pi_1 = \Gamma$, see [ibid.] and [Degtyarev 2009a] (torus type)
4.	$A_{16} \oplus A_3$	$(2, 0)$	$\pi_1 = \mathbb{Z}_6$, see [Artal Bartolo et al. 2002]
5.	$A_{16} \oplus A_2 \oplus A_1$	$(1, 1)$	$\pi_1 = \mathbb{Z}_6$, see [ibid.]
6.	$A_{15} \oplus A_4$	$(0, 1)^*$	$\pi_1 = \mathbb{Z}_6$, see [ibid.]
7.	$A_{14} \oplus A_4 \oplus A_1$	$(0, 3)$	
8.	$(A_{14} \oplus A_2) \oplus A_3$	$(1, 0)$	$\pi_1 = \Gamma$, see Section 4E (torus type)
9.	$(A_{14} \oplus A_2) \oplus A_2 \oplus A_1$	$(1, 0)$	$\pi_1 = \Gamma$, see Section 4F (torus type)
10.	$A_{13} \oplus A_6$	$(0, 2)$	
11.	$A_{13} \oplus A_4 \oplus A_2$	$(2, 0)$	
12.	$A_{12} \oplus A_7$	$(0, 1)$	
13.	$A_{12} \oplus A_6 \oplus A_1$	$(1, 1)$	
14.	$A_{12} \oplus A_4 \oplus A_3$	$(1, 0)$	
15.	$A_{12} \oplus A_4 \oplus A_2 \oplus A_1$	$(1, 1)$	
16.	$A_{11} \oplus 2A_4$	$(2, 0)$	
17.	$(A_{11} \oplus 2A_2) \oplus A_4$	$(1, 0)$	$\pi_1 = \Gamma$, see Section 4G (torus type)
18.	$A_{10} \oplus A_9$	$(2, 0)^*$	
19.	$A_{10} \oplus A_8 \oplus A_1$	$(1, 1)$	
20.	$A_{10} \oplus A_7 \oplus A_2$	$(2, 0)$	
21.	$A_{10} \oplus A_6 \oplus A_3$	$(0, 1)$	
22.	$A_{10} \oplus A_6 \oplus A_2 \oplus A_1$	$(1, 1)$	
23.	$A_{10} \oplus A_5 \oplus A_4$	$(2, 0)$	
24.	$A_{10} \oplus 2A_4 \oplus A_1$	$(1, 1)$	
25.	$A_{10} \oplus A_4 \oplus A_3 \oplus A_2$	$(1, 0)$	
26.	$A_{10} \oplus A_4 \oplus 2A_2 \oplus A_1$	$(2, 0)$	
27.	$A_9 \oplus A_6 \oplus A_4$	$(1, 1)^*$	
28.	$A_9 \oplus 2A_4 \oplus A_2$	$(1, 0)^*$	$\pi_1 = (2, 2)$, see [Degtyarev 2009b] (\mathbb{D}_{10} -sextic)
29.	$(2A_8) \oplus A_3$	$(1, 0)$	$\pi_1 = \Gamma$, see [Degtyarev 2009a] (torus type)
30.	$A_8 \oplus A_7 \oplus A_4$	$(0, 1)$	
31.	$A_8 \oplus A_6 \oplus A_4 \oplus A_1$	$(1, 1)$	
32.	$(A_8 \oplus A_5 \oplus A_2) \oplus A_4$	$(0, 1)$	nt104 in [Oka and Pho 2002] (torus type)
33.	$(A_8 \oplus 3A_2) \oplus A_4 \oplus A_1$	$(1, 0)$	$\pi_1 = (1, 3)$, see Section 4H (torus type)
34.	$A_7 \oplus 2A_6$	$(0, 1)$	
35.	$A_7 \oplus A_6 \oplus A_4 \oplus A_2$	$(2, 0)$	
36.	$A_7 \oplus 2A_4 \oplus 2A_2$	$(1, 0)$	
37.	$3A_6 \oplus A_1$	$(1, 0)$	$\pi_1 = \mathbb{Z}_3 \times \mathbb{D}_{14}$, see Section 4C (\mathbb{D}_{14} -sextic)
38.	$2A_6 \oplus A_4 \oplus A_2 \oplus A_1$	$(2, 0)$	
39.	$A_6 \oplus A_5 \oplus 2A_4$	$(2, 0)$	

An * marks sets of singularities that are realized by reducible sextics as well.

There are 42 real and 20 pairs of complex conjugate curves.

Table 1. Irreducible maximizing sextics with A type singularities.

by such sextics) and [Shimada 2007] (a description of the moduli spaces). We represent the result in Table 1, where the column (r, c) shows the number of classes: r is the number of real sextics, and c is the number of pairs of complex conjugate ones. The approach developed further in the paper lets one compute (or at least estimate) the fundamental group of a sextic with A type singularities, provided that its equation is known. In the literature, I could find explicit equations for lines 1–6, 8, 9, 17, 28, 29, 32, 33, and 37. With the results of this paper (Theorems 1.1 and 1.2) taken into account, the groups of all these sextics except $(A_8 \oplus A_5 \oplus A_2) \oplus A_4$, line 32 (which is not real), are known.

Remark 2.1. Unfortunately, our approach does not always work even if the curve is real. Thus, each of the two sextics with the set of singularities A_{19} , line 1, has a single real point (the isolated singular point of type A_{19} ; see [Artal Bartolo et al. 2002] for the equations) and Theorem 3.16 does not provide enough relations to compute the group.

2C. Known results. The fundamental group of the \mathbb{D}_{10} -special sextic with the set of singularities $A_9 \oplus 2A_4 \oplus A_2$, line 28 in Table 1, can be described as follows; see [Degtyarev 2009b] (where $'$ temporarily stands for the commutant of a group):

$$(2.2) \quad \pi_1/\pi_1'' = \mathbb{Z}_3 \times \mathbb{D}_{10}, \quad \pi_1'' = SL(2, \mathbb{k}_9),$$

where \mathbb{k}_9 is the field of nine elements. The fundamental groups of the first twelve sextics, lines 1–6, have been found in [Artal Bartolo et al. 2002]: with the exception of $(A_{17} \oplus A_2)$, line 3 (sextic of torus type, $\pi_1 = \Gamma$), they are all abelian. To my knowledge, the groups not mentioned in Table 1 have not been computed yet.

3. The braid monodromy

3A. Hirzebruch surfaces. A Hirzebruch surface Σ_d , $d > 0$, is a geometrically ruled rational surface with a (unique) exceptional section E of self-intersection $-d$. Typically, we use affine coordinates (x, y) in Σ_d such that E is given by $y = \infty$; then, x can be regarded as an affine coordinate in the base of the ruling. (The line $\{x = \infty\}$ plays no special rôle; usually, it is assumed sufficiently generic.) The fiber of the ruling over a point x in the base is denoted by F_x , and the affine fiber over x is $F_x^\circ := F_x \setminus E$. This is an affine space over \mathbb{C} ; in particular, one can speak about convex hulls in F_x° .

An n -gonal curve is a reduced curve $C \subset \Sigma_d$ intersecting each fiber at n points, i.e., such that the restriction to C of the ruling $\Sigma_d \rightarrow \mathbb{P}^1$ is a map of degree n . A singular fiber of an n -gonal curve C is a fiber F of the ruling intersecting $C + E$ geometrically at fewer than $(n + 1)$ points. A singular fiber F is proper if C does not pass through $F \cap E$. The curve C is proper if so are all its singular fibers. In other words, C is proper if it is disjoint from E .

In affine coordinates (x, y) as above an n -gonal curve $C \subset \Sigma_d$ is given by a polynomial of the form $\sum_{i=0}^n a_i(x)y^i$, where $\deg a_i \leq m + d(n - i)$ for some $m \geq 0$ (in fact, $m = C \cdot E$) and at least one polynomial a_i does have the prescribed degree (so that C does not contain the fiber $\{x = \infty\}$). The curve is proper if and only if $m = 0$; in this case $a_n(x) = \text{const}$.

A proper n -gonal curve $C \subset \Sigma_d$ defines a distinguished *zero section* $Z \subset \Sigma_d$, sending each point $x \in \mathbb{P}^1$ to the barycenter of the n points of $F_x^\circ \cap C$. Certainly, this section does not need to coincide with $\{y = 0\}$, which depends on the choice of the coordinates.

3B. The cubic resolvent. Consider a reduced *real* quartic polynomial

$$(3.1) \quad f(x, y) := y^4 + p(x)y^2 + q(x)y + r(x),$$

so that its roots y_1, y_2, y_3, y_4 (at each point x) satisfy $y_1 + y_2 + y_3 + y_4 = 0$, and consider the (*modified*) *cubic resolvent* of f

$$(3.2) \quad y^3 - 2p(x)y^2 + b_1(x)y + q(x)^2, \quad b_1 := p^2 - 4r,$$

and its reduced form

$$(3.3) \quad \bar{y}^3 + g_2(x)\bar{y} + g_3(x)$$

obtained by the substitution $y = \bar{y} + \frac{2}{3}p$. The discriminants of (3.1)–(3.3) are equal:

$$(3.4) \quad D = 16p^4r - 4p^3q^2 - 128p^2r^2 + 144pq^2r - 27q^4 + 256r^3.$$

Recall that $D = 0$ if and only if (3.1), or, equivalently, (3.2) or (3.3), has a multiple root. Otherwise, $D < 0$ if and only if exactly two roots of (3.1) are real. The roots of (3.2) are

$$(3.5) \quad \begin{aligned} \alpha &:= (y_1 + y_2)(y_3 + y_4) = -(y_1 + y_2)^2, \\ \beta &:= (y_1 + y_3)(y_2 + y_4) = -(y_1 + y_3)^2, \\ \gamma &:= (y_1 + y_4)(y_2 + y_3) = -(y_1 + y_4)^2, \end{aligned}$$

and those of (3.3) are obtained from (3.5) by shifting the barycenter $\frac{1}{3}(\alpha + \beta + \gamma)$ to zero.

Remark 3.6. If $\{f(x, y) = 0\}$ is a proper tetragonal curve in a Hirzebruch surface Σ_d , then (3.2) defines a proper trigonal curve $C' \subset \Sigma_{2d}$ and a distinguished section $S := \{y = 0\}$ (in general, *other* than the zero section) which is tangent (more precisely, has even intersection index at each intersection point) to C' . Conversely, (3.1) can be recovered from (3.2) (together with the section $S = \{y = 0\}$) uniquely up to the automorphism $y \mapsto -y$, which takes q to $-q$.

Remark 3.7. One has

$$q = -(y_1 + y_2)(y_1 + y_3)(y_1 + y_4);$$

hence, q vanishes if and only if two of the roots of (3.1) are opposite. If all roots are nonreal, $y_{1,2} = \alpha \pm \beta'i$, $y_{3,4} = -\alpha \pm \beta''i$, $\alpha, \beta', \beta'' \in \mathbb{R}$, then

$$q = 2\alpha(\beta'^2 - \beta''^2), \quad b_1 = -8\alpha^2(\beta'^2 + \beta''^2) + (\beta'^2 - \beta''^2)^2.$$

Hence, $q(x) = 0$ if and only if either $\alpha = 0$ (and then $b_1(x) > 0$, assuming that $D(x) \neq 0$) or $\beta' = \pm\beta''$ (and then $b_1(x) < 0$). If $y_1 = y_2$, i.e., $\beta' = 0$, then $q(x) > 0$ if and only if one has the inequality $y_1 < \operatorname{Re} y_3 = \operatorname{Re} y_4$ equivalent to $y_1 < 0$.

Remark 3.8. Observe also that, if $y_1 = y_2$, then g_3 takes the form

$$g_3 = \frac{2}{27}(y_1 - y_4)^3(y_1 - y_3)^3.$$

Hence, $g_3(x) < 0$ if and only if the two other roots are real and separated by the double root $y_1 = y_2$. Otherwise, either $y_1 < \operatorname{Re} y_3, \operatorname{Re} y_4$ or $y_1 > \operatorname{Re} y_3, \operatorname{Re} y_4$, and, in view of Remark 3.7, the former holds if and only if $q(x) > 0$.

3C. The real monodromy. Choose affine coordinates (x, y) in the Hirzebruch surface Σ_d so that the exceptional section E is $\{y = \infty\}$. Consider a *real* proper tetragonal curve $C \subset \Sigma_d$; it is given by a real polynomial $f(x, y)$ as in (3.1). Over a generic real point $x \in \mathbb{R}$, the four points y_1, \dots, y_4 of the intersection $C \cap F_x^\circ$ can be ordered lexicographically, according to the *decreasing* of $\operatorname{Re} y$ first and $\operatorname{Im} y$ second. We always assume this ordering. Then, choosing a real reference point $y \gg 0$, we have a *canonical* geometric basis $\{\alpha_1, \dots, \alpha_4\}$ for the fundamental group $\pi(x) := \pi_1(F_x^\circ \setminus C, y)$; see Figure 1.

Let x_1, \dots, x_r be all *real* singular fibers of C , ordered by increasing. For each i , consider a pair of nonsingular fibers $x_i^- := x_i - \epsilon$ and $x_i^+ := x_i + \epsilon$, where ϵ is a sufficiently small positive real number; see Figure 2. Define $x_0 = x_{r+1} = \infty$ and,




Figure 1. The canonical basis.




Figure 2. The monodromies β_i and γ_j .

assuming the fiber $x = \infty$ is nonsingular, pick also a pair of real nonsingular fibers $x_{r+1}^- = x_\infty^- := R \gg 0$ and $x_0^+ = x_\infty^+ := -R$. Identify all groups $\pi(x_i^\pm)$ with the free group \mathbb{F}_4 by means of their respective canonical bases. (All reference points are chosen in a real section $y = \text{const} \gg 0$, which is assumed disjoint from the fiberwise convex hull of C over the disk $|x| \leq R$.) Consider the semicircles $t \mapsto x_i + \epsilon e^{i\pi(1-t)}$, $t \in [0, 1]$, and the line segments $t \mapsto t$, $t \in [x_j^+, x_{j+1}^-]$; see Figure 2. These paths give rise to the *monodromy isomorphisms*

$$\beta_i : \pi(x_i^-) \rightarrow \pi(x_i^+), \quad \gamma_j : \pi(x_j^+) \rightarrow \pi(x_{j+1}^-),$$

$i = 1, \dots, r$, $j = 0, \dots, r$. In addition, we also have the monodromy $\beta_0 = \beta_\infty = \beta_{r+1} : \pi(x_\infty^-) \rightarrow \pi(x_\infty^+)$ along the semicircle $t \mapsto Re^{i\pi t}$, $t \in [0, 1]$, and the *local monodromies*

$$\mu_i : \pi(x_i^+) \rightarrow \pi(x_i^+), \quad i = 1, \dots, r$$

along the circles $t \mapsto x_i + \epsilon e^{2\pi it}$, $t \in [0, 1]$. Using the identifications $\pi(x_i^\pm) = \mathbb{F}_4$ fixed above, all β_i, μ_i, γ_j can be regarded as elements of the automorphism group $\text{Aut } \mathbb{F}_4$, and as such they belong to the braid group \mathbb{B}_4 . Recall (see [Artin 1947]) that Artin’s *braid group* $\mathbb{B}_4 \subset \text{Aut}\langle \alpha_1, \dots, \alpha_4 \rangle$ is the subgroup consisting of the automorphisms taking each generator α_i to a conjugate of a generator and preserving the product $\alpha_1\alpha_2\alpha_3\alpha_4$. It is generated by the three braids

$$\sigma_i : \alpha_i \mapsto \alpha_i\alpha_{i+1}\alpha_i^{-1}, \quad \alpha_{i+1} \mapsto \alpha_i, \quad i = 1, 2, 3,$$

the defining relations being $\{\sigma_1, \sigma_2\}_3 = \{\sigma_2, \sigma_3\}_3 = [\sigma_1, \sigma_3] = 1$.

3D. The computation. The braids β_i, μ_i , and γ_j introduced in the previous section are easily computed from the real part $C_\mathbb{R} \subset \mathbb{R}^2$ of the curve. In the figures, we use the following notation:

- real branches of C are represented by solid bold lines;

- pairs y_i, y_{i+1} of complex conjugate branches are represented by dotted lines (showing the common real part $\text{Re } y_i = \text{Re } y_{i+1}$);
- relevant fibers of Σ_d are represented by vertical dotted grey lines.

Certainly, the dotted lines are not readily seen in the figures; however, in most cases, it is only the intersection indices that matter, and the latter are determined by the indexing of the branches at the starting and ending positions.

We summarize the results in the next three statements. The first one is obvious: essentially, one speaks about the link of the singularity $y^4 - x^{4d}$.

Lemma 3.9. *Assume that $R \gg 0$ is so large that the disk $\{|x| < R\}$ contains all singular fibers of C . Then one has $\beta_\infty = \Delta^d$, where $\Delta := \sigma_1\sigma_2\sigma_3\sigma_1\sigma_2\sigma_1 \in \mathbb{B}_4$ is the Garside element.* ◁

The following lemma is easily proved by considering the local normal forms of the singularities. (In the simplest case of a vertical tangent, the circumventing braids β are computed, e.g., in [Orevkov 1999]; the general case is completely similar.) For the statement, we extend the standard notation $A_m, m \geq 1$, to A_0 to designate a simple tangency of C and the fiber.

Lemma 3.10. *The braids β_j and μ_j about a singular fiber x_j of type $A_m, m \geq 0$, depend only on m and the pair $(i, i + 1)$ of indices of the branches that merge at the singular point. They are as shown in Figure 3.* ◁

Remark 3.11. At a point of type A_{2k-1} , it is *not* important whether the two branches of C at this point are real or complex conjugate. On the other hand, at a point of type A_{2k} it *does* matter whether the number of real branches increases or decreases. If a fiber contains two double points, with indices $(1, 2)$ and $(3, 4)$, then the powers of σ_1 and σ_3 contributed to β or μ by each of the points are multiplied; since σ_1 and σ_3 commute, the order is not important.

The following statement is our principal technical tool, most important being Figure 4, right, describing the behavior of the “invisible” branches. (Note that the




Figure 3. The braids β and μ .




Figure 4. The braids γ .

two dotted lines in the figure may cross; the permutation of the branches depends on the parity of the twist parameter t introduced in the statement.)

Proposition 3.12. *Let I be a real segment in the x -axis free of singular fibers of C . Then the monodromy γ over I is*

- identity, if all four branches of C over I are real, and
- as shown in [Figure 4](#) otherwise.

Here, $\tau := \sigma_2^{-1} \sigma_3 \sigma_1^{-1} \sigma_2$ and the twist parameter t in [Figure 4](#), right, is the number of roots $x' \in I$ of the coefficient $q(x)$ (see (3.1)) such that $b_1(x') > 0$ (see (3.2)) and q changes sign at x' ; each root x' contributes $+1$ or -1 depending on whether q is increasing or decreasing at x' , respectively.

Proof. The only case that needs consideration, viz. that of four nonreal branches (see [Figure 4](#), right), is given by [Remark 3.7](#). Indeed, the canonical basis in the fiber F_x° over $x \in I$ changes when the real parts of all four branches vanish, and this happens when $q(x) = 0$ and $b_1(x) > 0$. This change contributes $\tau^{\pm 1}$ to γ , and the sign ± 1 (the direction of rotation) depends on whether q increases or decreases. \square

Remark 3.13. A longer segment I with exactly two real branches of C over it can be divided into smaller pieces I_1, I_2, \dots , each containing a single crossing point as in [Figure 4](#); then, the monodromy γ over I is the product of the contributions of each piece. In fact, as explained above, the precise position and number of crossings is irrelevant; what only matters is the final permutation between the endpoints of I . For example, to minimize the number of elementary pieces, one can always assume that the branches, both bold and dotted, are monotonous.

3E. The Zariski–van Kampen theorem. We are interested in the fundamental group $\pi_1 := \pi_1(\Sigma_{\tilde{d}} \setminus (\tilde{C} \cup E))$, where $\tilde{C} \subset \Sigma_{\tilde{d}}$ is a real tetragonal curve, possibly improper, and $E \subset \Sigma_{\tilde{d}}$ is the exceptional section. To compute π_1 , we consider the proper model $C \subset \Sigma_d$, obtained from \tilde{C} by blowing up all points of intersection $\tilde{C} \cap E$ and blowing down the corresponding fibers. In addition to the braids β_i, μ_i , and γ_j introduced in [Section 3C](#), to each real singular fiber x_i of C we assign its local slope $\varkappa_i \in \pi(x_i^+)$, which depends on the type of the corresponding singular fiber of

the original curve \tilde{C} . Roughly, consider a small analytic disk $\Phi \subset \Sigma_d$ transversal to the fiber F_{x_i} and disjoint from C and E , and a similar disk $\tilde{\Phi} \subset \Sigma_{\tilde{d}}$ with respect to \tilde{C} . Let $\tilde{\Phi}' \subset \Sigma_d$ be the image of $\tilde{\Phi}$, and assume that the boundaries $\partial\Phi$ and $\partial\tilde{\Phi}'$ have a common point in the fiber over x_i^+ . Then the loop $[\partial\tilde{\Phi}'] \cdot [\partial\Phi]^{-1}$ is homotopic to a certain class $\varkappa_i \in \pi(x_i^+)$, well defined up to a few moves irrelevant in the sequel. This class is the slope.

Roughly, the slope measures (in the form of the twisted monodromy; see the definitions prior to [Theorem 3.16](#)) the deviation of the braid monodromy of an improper curve \tilde{C} from that of its proper model C . Slopes appear in the relation at infinity as well, compensating for the fact that, near improper singular fibers, the curve intersects *any* section of $\Sigma_{\tilde{d}}$. Details and further properties are found in [[Degtyarev 2012](#), Section 5.1.3]; in this paper, slopes are used in [Theorem 3.16](#).

Remark 3.14. In all examples considered below, $\tilde{C} \subset \Sigma_{d-1}$ has a single improper fiber F , where \tilde{C} has a singular point of type \tilde{A}_m , $m \geq 1$, maximally transversal to both E and F . If $F = \{x = 0\}$, such a curve \tilde{C} is given by a polynomial \tilde{f} of the form $\sum_{i=0}^4 y^i a_i(x)$ with $a_4(x) = x^2$ and $x \mid a_3(x)$, and the defining polynomial of its transform $C \subset \Sigma_d$ is $f_{nr}(x, y) := x^2 \tilde{f}(x, y/x)$. The corresponding singular fiber of C has a node A_1 at $(0, 0)$ and another double point A_{m-2} (assuming $m \geq 2$).

Thus, the only nontrivial example relevant in the sequel is the one described below. (By the very definition, at each singular fiber x_i proper for \tilde{C} the slope is $\varkappa_i = 1$.) A great deal of other examples of both computing the slopes and using them in the study of the fundamental group are found in [[Degtyarev 2012](#)].

Example 3.15. At the only improper fiber $x_i = 0$ described in [Remark 3.14](#) the slope is the class of $\alpha_j \alpha_{j+1}$, where $(j, j + 1)$ are the two branches merging at the node; see [Figure 3](#). This fact can easily be seen using a local model. In a small neighborhood of $x = 0$, one can assume that \tilde{C} is given by $(y - a)(y - b) = 0$. Let $\tilde{\Phi} \subset \Sigma_{\tilde{d}}$ and $\Phi \subset \Sigma_d$ be the disk $\{y = c, |x| \leq 1\}$, $c \in \mathbb{R}$ and $c \gg |a|, |b|$. Then, the relevant part of C is the node $(y - ax)(y - bx) = 0$, and $\tilde{\Phi}$ projects onto the disk $\tilde{\Phi}' = \{y = cx, |x| \leq 1\}$, which meets Φ at $(1, c)$. Now, consider one full turn $x = \exp(2\pi it)$, $t \in [0, 1]$, and follow the point (x, cx) in $\partial\tilde{\Phi}'$: it describes the circle $y = c \exp(2\pi it)$ encompassing once the two points of the intersection $C \cap F_1^\circ$. The class $\alpha_j \alpha_{j+1}$ of this circle is the slope. Even more precisely, one should start with the constant path $[0, 1] \rightarrow (1, c)$ and homotope this path in $F_x^\circ \setminus C$, keeping one end in Φ and the other in $\tilde{\Phi}'$. In the terminal position, the path is a loop again, and its class $\alpha_j \alpha_{j+1}$ is the slope.

Define the *twisted local monodromy* $\tilde{\mu}_i := \mu_i \cdot \text{inn } \varkappa_i$, where $\text{inn} : G \rightarrow \text{Aut } G$ is the homomorphism sending an element g of a group G to the inner automorphism $\text{inn } g : h \mapsto g^{-1}hg$. Thus, $\tilde{\mu}_i : \pi(x_i^+) \rightarrow \pi(x_i^+)$ is the map $\alpha \mapsto \varkappa_i^{-1}(\alpha \uparrow \mu_i)\varkappa_i$. In

general, $\tilde{\mu}_i$ is not a braid. Take $x_0^+ = x_\infty^+$ for the reference fiber and consider the braids

$$\rho_i := \prod_{j=1}^i \gamma_{j-1} \beta_j : \pi(x_0^+) \rightarrow \pi(x_i^+), \quad i = 1, \dots, r+1 = \infty$$

(left to right product), the (global) slopes

$$\tilde{\varkappa}_i := \varkappa_i \uparrow \rho_1^{-1} \in \pi(x_0^+), \quad i = 1, \dots, r,$$

and the twisted monodromy homomorphisms

$$\tilde{m}_i := \rho_i \tilde{\mu}_i \rho_i^{-1} : \pi(x_0^+) \rightarrow \pi(x_0^+), \quad i = 1, \dots, r.$$

The following theorem is essentially due to Zariski and van Kampen [van Kampen 1933], and the particular case of improper curves in Hirzebruch surfaces, treated by means of the slopes, is considered in detail in [Degtyarev 2012, Section 5.1.3]. Here, we state and outline the proof of a very special case of this approach, incorporating the (partial) computation of the braid monodromy of a real tetragonal curve in terms of its real part.

We use the following common convention: given an automorphism β of the free group $\langle \alpha_1, \dots, \alpha_4 \rangle$, the *braid relation* $\beta = \text{id}$ stands for the quadruple of relations $\alpha_j \uparrow \beta = \alpha_j$, $j = 1, \dots, 4$. Note that, since β is an automorphism, this is equivalent to the infinitely many relations $\alpha = \alpha \uparrow \beta$, $\alpha \in \langle \alpha_1, \dots, \alpha_4 \rangle$.

Theorem 3.16. *In the notation above, the inclusion of the reference fiber induces an epimorphism*

$$\pi(x_0^+) = \langle \alpha_1, \dots, \alpha_4 \rangle \twoheadrightarrow \pi_1,$$

and the relations $\tilde{m}_i = \text{id}$, $i = 1, \dots, r$, hold in π_1 . If the fiber $x = \infty$ is nonsingular and all nonreal singular fibers are proper for \tilde{C} , then one also has the relations at infinity $\rho_\infty = \text{id}$ and $(\alpha_1 \cdots \alpha_4)^d = \tilde{\varkappa}_r \cdots \tilde{\varkappa}_1$. If, in addition, C has at most one pair of conjugate nonreal singular fibers, then the relations listed define π_1 .

Proof. The assertion is a restatement of the classical Zariski–van Kampen theorem modified for the case of improper curves; see [Degtyarev 2012, Theorem 5.50]. The relation at infinity $(\alpha_1 \cdots \alpha_4)^d = \tilde{\varkappa}_r \cdots \tilde{\varkappa}_1$ holds in π_1 whenever all slopes not accounted for, namely those at the nonreal fibers, are known to be trivial. The automorphism $\rho_{r+1} : \pi(x_0^+) \rightarrow \pi(x_{r+1}^+) = \pi(x_0^+)$ is the monodromy along the “boundary” of the upper half-plane $\text{Im } x > 0$ (see Figure 2), i.e., the product of the monodromies about all singular fibers in this half-plane; if the slopes at these fibers are all trivial, then $\rho_{r+1} = \text{id}$ in π_1 . Finally, if \tilde{C} has at most one pair of conjugate nonreal singular fibers, then all but possibly one braid relations are present and hence they define the group; see [Degtyarev 2012, Lemma 5.59]. \square

4. The computation

4A. The strategy. We start with a plane sextic $D \subset \mathbb{P}^2$ and choose homogeneous coordinates $(z_0 : z_1 : z_2)$ so that D has a singular point of type A_m , $m \geq 3$, at $(0 : 0 : 1)$ tangent to the axis $\{z_1 = 0\}$. Then, in the affine coordinates $x := z_1/z_0$, $y := z_2/z_0$, the curve D is given by a polynomial \tilde{f} as in [Remark 3.14](#), and the same polynomial \tilde{f} defines a certain tetragonal curve $\tilde{C} \subset \Sigma_1$, viz. the proper transform of D under the blow-up of $(0 : 0 : 1)$. The common fundamental group

$$\pi_1 := \pi_1(\mathbb{P}^2 \setminus D) = \pi_1(\Sigma_1 \setminus (\tilde{C} \cup E))$$

is computed using [Theorem 3.16](#) applied to \tilde{C} and its transform $C \subset \Sigma_2$, with the only nontrivial slope $\varkappa = \alpha_1\alpha_2$ or $\alpha_3\alpha_4$ over $x = 0$ given by [Example 3.15](#). (Here, $E \subset \Sigma_1$ is the exceptional section, i.q. the exceptional divisor over the point $(0 : 0 : 1)$ blown up.) *A priori*, [Theorem 3.16](#) may only produce a certain group \mathfrak{g} that surjects onto π_1 rather than π_1 itself; however, in most cases this group \mathfrak{g} is “minimal expected” (see [Section 4D](#) below) and we do obtain π_1 .

The assumption that the fiber $x = \infty$ is nonsingular is not essential as long as the singularity over ∞ is taken into consideration: one can always move ∞ to a generic point by a real projective change of coordinates. To keep the defining equations as simple as possible, we assume such a change of coordinates implicitly. Furthermore, it is only the cyclic order of the singular fibers in the circle $\mathbb{P}_{\mathbb{R}}^1$ that matters, and sometimes we reorder the fibers by applying a cyclic permutation to their “natural” indices. In other words, the braid $\beta_{\infty} = \Delta^2$ is in the center of \mathbb{B}_4 and, hence, it can be inserted at any place in the relation $\gamma_0\beta_1\gamma_1 \cdots \gamma_r\beta_{\infty} = \text{id}$.

To compute the braids, we outline the real (bold lines) and imaginary (dotted lines) branches of C in the figures. Recall that it is only the mutual position of the real branches and their intersection indices with the imaginary ones that matters; see [Remark 3.13](#). The “special” node that contributes the only nontrivial slope (the blow-up center in the passage from C to \tilde{C} ; see [Remark 3.14](#)) is marked with a white dot; the other singular points of C (including those of type A_0) are marked with black dots. The shape of the curve can mostly be recovered using [Remarks 3.7](#) and [3.8](#); however, it is usually easier to determine the mutual position of the roots directly via `Maple`. The braids β_i , μ_i , and γ_j are computed from the figures as explained in [Section 3D](#).

Warning 4.1. The polynomial f_{nr} given by [Remark 3.14](#) is used to determine the slope and mutual position of the two singular points over $x = 0$: the “special” node is always at $(0, 0)$. For all other applications, e.g., for [Proposition 3.12](#), this polynomial should be converted to the reduced form [\(3.1\)](#).

4B. Relations. Recall that a braid relation $\tilde{m}_i = \text{id}$ stands for a quadruple of relations $\alpha_j \uparrow \tilde{m}_i = \alpha_j$, $j = 1, \dots, 4$. Alternatively, this can be regarded as an

infinite sequence of relations $\alpha \uparrow \tilde{m}_i = \alpha$, $\alpha \in \mathbb{F}_4$, or, equivalently, as a quadruple of relations $\alpha'_j \uparrow \tilde{m}_i = \alpha'_j$, $j = 1, \dots, 4$, where $\alpha'_1, \dots, \alpha'_4$ is any basis for \mathbb{F}_4 . For this reason, in the computation below we start with the braid relations $\alpha'_j \uparrow \tilde{\mu}_i = \alpha'_j$ in the canonical basis over x_i^+ and translate them to x_0^+ via ρ_i^{-1} . In the most common case $\tilde{\mu}_i = \sigma_r^p$, $r = 1, 2, 3$, $p \in \mathbb{Z}$, the whole quadruple is equivalent to the single relation $\{\alpha'_r, \alpha'_{r+1}\}_p = 1$, where

$$\{\alpha, \beta\}_{2k} := (\alpha\beta)^k (\beta\alpha)^{-k}, \quad \{\alpha, \beta\}_{2k+1} := (\alpha\beta)^k \alpha (\alpha\beta)^{-k} \beta^{-1}.$$

Remark 4.2. The braid relations about the fiber $x_k = 0$ with the only nontrivial slope (see [Example 3.15](#)) can also be presimplified. Let $\alpha'_1, \dots, \alpha'_4$ be the canonical basis in x_k^+ . If $\varkappa_k = \alpha'_1 \alpha'_2$ and $\mu_k = \sigma_1^2 \sigma_3^p$, the braid relations $\tilde{\mu}_k = \text{id}$ and relation at infinity $(\alpha'_1 \cdots \alpha'_4)^2 = \varkappa_k$ together are equivalent to

$$\alpha'_1 \alpha'_2 (\alpha'_3 \alpha'_4)^2 = \{\alpha'_3, \alpha'_4\}_{p+4} = 1.$$

Similarly, if $\varkappa_k = \alpha'_3 \alpha'_4$ and $\mu_k = \sigma_1^p \sigma_3^2$, we obtain

$$(\alpha'_1 \alpha'_2)^2 \alpha'_3 \alpha'_4 = \{\alpha'_1, \alpha'_2\}_{p+4} = 1.$$

Certainly, these relations should be translated back to x_0^+ via ρ_k^{-1} . Note, though, that we do not use this simplification in the sequel.

Remark 4.3. In some cases, simpler relations are obtained if another point x_i^+ , $i > 0$, is taken for the reference fiber. To do so, one merely replaces the braids ρ_j , $j = 1, \dots, r + 1 = \infty$, with $\rho'_j := \rho_i^{-1} \rho_j$.

All computations below were performed using GAP [[GAP Group 2008](#)], with the help of the simple braid manipulation routines contained in [[Degtyarev 2012](#)]. The GAP code can be found at <http://www.fen.bilkent.edu.tr/~degt/papers/papers.htm>. The processing is almost fully automated, the input being the braids β_i, μ_i, γ_j and the only nontrivial slope $\varkappa_k = \alpha_1 \alpha_2$ or $\alpha_3 \alpha_4$, which are read off from the diagrams depicting the curves.

4C. The set of singularities $3A_6 \oplus A_1$, [line 37](#). Any sextic with this set of singularities is \mathbb{D}_{14} -special (see [[Degtyarev 2008](#)]), and, according to [[Degtyarev and Oka 2009](#)], any \mathbb{D}_{14} -special sextic can be given by an equation of the form

$$\begin{aligned} & 2t(t^3 - 1)(z_0^4 z_1 z_2 + z_1^4 z_2 z_0 + z_2^4 z_0 z_1) \\ & + (t^3 - 1)(z_0^4 z_1^2 + z_1^4 z_2^2 + z_2^4 z_0^2) + t^2(t^3 - 1)(z_0^4 z_2^2 + z_1^4 z_0^2 + z_2^4 z_1^2) \\ & + 2t(t^3 + 1)(z_0^3 z_1^3 + z_1^3 z_2^3 + z_2^3 z_0^3) + 4t^2(t^3 + 2)(z_0^3 z_1^2 z_2 + z_1^3 z_2^2 z_0 + z_2^3 z_0^2 z_1) \\ & + 2(t^6 + 4t^3 + 1)(z_0^3 z_1 z_2^2 + z_1^3 z_2 z_0^2 + z_2^3 z_0 z_1^2) + t(t^6 + 13t^3 + 10)z_0^2 z_1^2 z_2^2, \end{aligned}$$




Figure 5. The set of singularities $3A_6 \oplus A_1$, [line 37](#).

$t^3 \neq 1$. The set of singularities of this curve is $3A_6 \oplus A_1$ if and only if $t^3 = -27$; we use the real value $t = -3$. After the substitution $z_0 = 1$, $z_1 = x + \frac{1}{3}$, and $z_2 = y/x$ the equation is brought to the form considered in [Remark 3.14](#). Up to a positive factor, the discriminant (3.4) with respect to y is

$$-x^5(27x^3 - 648x^2 + 6363x + 7)(3x - 2)^2(3x + 1)^7,$$

which has real roots

$$x_1 = -\frac{1}{3}, \quad x_2 \approx -0.001, \quad x_3 = 0, \quad x_4 = \frac{2}{3}, \quad x_5 = \infty$$

and two simple imaginary roots. Hence, [Theorem 3.16](#) does compute the group.

The only root of q on the real segment $[-\infty, x_1]$ is $x' \approx -3.48$, and $b_1(x') < 0$; hence, one has $\gamma_0 = \text{id}$; see [Proposition 3.12](#). The other braids β_i, γ_j are easily found from [Figure 5](#), and, using [Theorem 3.16](#) and GAP, we obtain a group of order 42. This concludes the proof of [Theorem 1.1](#). \square

4D. Sextics of torus type. All maximal, in the sense of degeneration, sextics of torus type are described in [\[Oka and Pho 2002\]](#), where a sextic D is represented by a pair of polynomials $f_2(x, y), f_3(x, y)$ of degree 2 and 3, respectively, so that the defining polynomial of D is $f_{\text{tor}} := f_2^3 + f_3^2$. (Below, these equations are cited in a slightly simplified form: I tried to clear the denominators by linear changes of variables and appropriate coefficients.) Each curve (at least, each of those considered below) has a type $A_m, m \geq 3$, singularity at $(0, 0)$ tangent to the y -axis. Hence, we start with the substitution $\tilde{f}(x, y) := y^6 f_{\text{tor}}(x/y, 1/y)$ to obtain a polynomial \tilde{f} as in [Remark 3.14](#); then we proceed as in [Section 4A](#).

To identify the group g given by [Theorem 3.16](#) as Γ , we use the following GAP code, which was suggested to me by E. Artal:

```
(4.4) P := PresentationNormalClosure(g, Subgroup(g, a));
      SimplifyPresentation(P);
```

here, a is an appropriate ratio $\alpha_i \alpha_j^{-1}$ which normally generates the commutant of g . If the resulting presentation has two generators and no relations, we conclude

that $g = \pi_1 = \Gamma$, even when the statement of [Theorem 3.16](#) does not guarantee a complete set of relations. Indeed, a priori we have epimorphisms $g \twoheadrightarrow \pi_1 \twoheadrightarrow \Gamma$ (the latter follows from the fact that the curve is assumed to be of torus type), which induce epimorphisms $[g, g] \twoheadrightarrow [\pi_1, \pi_1] \twoheadrightarrow [\Gamma, \Gamma] = \mathbb{F}_2$ of the commutants. If $[g, g] = \mathbb{F}_2$, both these epimorphisms are isomorphisms (since \mathbb{F}_2 is Hopfian) and the 5-lemma implies that $g \twoheadrightarrow \pi_1 \twoheadrightarrow \Gamma$ are also isomorphisms.

In some cases (e.g., in Sections 4E and 4F), the call `SimplifiedFpGroup(g)` returns a recognizable presentation of Γ .

4E. The set of singularities $(A_{14} \oplus A_2) \oplus A_3$, [line 8](#). The curve in question is nt139 in [\[Oka and Pho 2002\]](#):

$$f_2 = 80(-36y^2 + 120xy - 82x^2 + 2x),$$

$$f_3 = 100(-1512y^3 + 7794y^2x - 18y^2 - 11664yx^2 + 144xy + 5313x^3 - 194x^2 + x).$$

Up to a positive coefficient, the discriminant of f_{nr} is

$$x^{13}(5120x^4 + 36864x^3 + 3456x^2 - 2160x - 405)(x - 1)^3.$$

It has five real roots, which we reorder cyclically as follows:

$$x_1 = 0, \quad x_2 \approx 0.27, \quad x_3 = 1, \quad x_4 = \infty, \quad x_5 \approx -7.1.$$

Besides, there are two conjugate imaginary singular fibers, which are of type A_0 .

The curve is depicted in [Figure 6](#), from which all braids β_i, γ_j are easily found. Taking x_0^+ for the reference fiber and using $a = \alpha_1\alpha_2^{-1}$ in [\(4.4\)](#), we obtain $\pi_1 = \Gamma$.

4F. The set of singularities $(A_{14} \oplus A_2) \oplus A_2 \oplus A_1$, [line 9](#). The curve is nt142 in [\[Oka and Pho 2002\]](#):

$$f_2 = -45y^2 - 240yx - 106x^2 + 90x,$$

$$f_3 = 1025y^3 + 6045y^2x - 375y^2 + 5490yx^2 - 4050yx + 1354x^3 - 2040x^2 + 750x.$$

Up to a positive coefficient, the discriminant of f_{nr} is

$$x^{13}(8x^3 - 10720x^2 + 14250x - 5625)(x + 1)^2(14x + 15)^3,$$




Figure 6. The set of singularities $(A_{14} \oplus A_2) \oplus A_3$, [line 8](#).




Figure 7. The set of singularities $(A_{14} \oplus A_2) \oplus A_2 \oplus A_1$, line 9.

and all its roots are real:

$$x_1 = -\frac{15}{14}, \quad x_2 = -1, \quad x_3 = 0, \quad x_4 \approx 1338, \quad x_5 = \infty.$$

The braids β_i, γ_j are found from Figure 7 and, using x_0^+ as the reference fiber and $a = \alpha_1 \alpha_2^{-1}$ in (4.4), we conclude that $\pi_1 = \Gamma$.

4G. The set of singularities $(A_{11} \oplus 2A_2) \oplus A_4$, line 17. This is nt118 in [Oka and Pho 2002]:

$$f_2 = \frac{1}{5}(-3456y^2 + 1200yx - 3005x^2 + 240x),$$

$$f_3 = \frac{1}{5}(-89856y^3 + 130464y^2x - 6912y^2 - 112680yx^2 + 8640yx + 91345x^3 - 13320x^2 + 480x).$$

Up to a positive coefficient, the discriminant of f_{nr} is

$$-x^{10}(25x^3 + 290x^2 + 360x + 162)(35x^2 - 384x + 1152)^3.$$

It has three real roots, which we reorder cyclically as follows:

$$x_1 = 0, \quad x_2 = \infty, \quad x_3 \approx -10.26.$$

In addition, there are two pairs of complex conjugate singular fibers, of types A_2 and A_0 . Thus, *a priori* Theorem 3.16 only gives us a certain epimorphism $g \rightarrow \pi_1$. However, using $a = \alpha_1 \alpha_2^{-1}$ in (4.4), we conclude that $g = \pi_1 = \Gamma$. (All braids are found from Figure 8 and the reference fiber is x_1^+ ; see Remark 4.3.)




Figure 8. The set of singularities $(A_{11} \oplus 2A_2) \oplus A_4$, line 17.

4H. *The set of singularities* $(A_8 \oplus 3A_2) \oplus A_4 \oplus A_1$, [line 33](#). This curve is nt83 in [\[Oka and Pho 2002\]](#):

$$(4.5) \quad \begin{aligned} f_2 &= -565y^2 - 14yx + 176y - 5x^2 + 104x - 16, \\ f_3 &= 13321y^3 + 3135y^2x - 6294y^2 + 207yx^2 - 3516yx + 1056y \\ &\quad + 25x^3 - 558x^2 + 624x - 64. \end{aligned}$$

Up to a positive coefficient, the discriminant of f_{nr} is

$$x^3(x+3)(x+9)^2(11915x^3 + 96579x^2 - 14823x + 729)^3(x-9)^9.$$

It has five real roots, which we reorder cyclically as follows:

$$x_1 = 0, \quad x_2 = 9, \quad x_3 = -9, \quad x_4 \approx -8.26, \quad x_5 = -3.$$

We conclude that the curve has only two nonreal singular fibers, which are cusps. Hence, [Theorem 3.16](#) gives us a complete presentation of π_1 .

In the interval (x_5, x_1) , where f has four imaginary branches, q has four roots

$$x'_1 \approx -2.93, \quad x'_2 = -1.92, \quad x'_3 \approx -0.79, \quad x'_4 \approx -0.14,$$

with b_1 negative at x'_1, x'_3 and positive at x'_2, x'_4 ; at the latter two points one also has $q' < 0$. Hence, $\gamma_0 = \tau^{-2}$; see [Proposition 3.12](#). All other braids are easily found from [Figure 9](#).

Remark 4.6. For a further simplification, observe that the braid ρ_∞ appearing in [Theorem 3.16](#) equals

$$\sigma_2^{-1} \sigma_1 \sigma_3^{-1} \sigma_1 \sigma_3^{-1} \sigma_2 \cdot \sigma_1^{-1} \cdot \sigma_2^{-1} \sigma_1 \cdot \sigma_2^{-4} \cdot \sigma_3^{-1} \cdot \sigma_2^{-2} \cdot \sigma_3^{-1} \sigma_2 \cdot \sigma_1^{-1} \cdot (\sigma_3 \sigma_1 \sigma_2)^4,$$

and one can check that $\rho_\infty = \rho_{im}^{-1} \sigma_1^3 \rho_{im}$, where $\rho_{im} := \sigma_2 \sigma_1^{-1} \sigma_3^2 \sigma_2$. (Note that ρ_∞ represents the monodromy about a single imaginary cusp of the curve; hence, it is expected to be conjugate to σ_1^3 .) Thus, we can replace the quadruple of relations $\rho_\infty = \text{id}$ with a single relation $\{\alpha_1, \alpha_2\}_3 \uparrow \rho_{im} = 1$; see [Section 4B](#).

Now, taking x_3^+ for the reference fiber (see [Remark 4.3](#)), using [Remark 4.6](#), and applying `SimplifiedFpGroup(g)`, we arrive at [\(1.3\)](#). This presentation has three




Figure 9. The set of singularities $(A_8 \oplus 3A_2) \oplus A_4 \oplus A_1$, [line 33](#), projected from A_4 .




Figure 10. The set of singularities $(A_8 \oplus 3A_2) \oplus A_4 \oplus A_1$, line 33, projected from A_8 .

generators and four relations of total length 48. Together with the previous sections, this concludes the proof of Theorem 1.2. \square

Remark 4.7. The Alexander module of the group π_1 considered in this section is $\mathbb{Z}[t, t^{-1}]/(t^2 - t + 1)$, and the finite quotients π_1/α_2^p , $p = 2, 3, 4$, are isomorphic to the similar quotients of Γ . My laptop failed to compute the order of π_1/α_2^5 .

Remark 4.8. In (4.5), the singular point at the origin is of type A_4 . One can start with a change of variables $x \mapsto y + 9$, $y \mapsto x + 1$ and resolve the type A_8 point instead. The tetragonal model is depicted in Figure 10, and the computation becomes slightly simpler, but the resulting presentation is of the same complexity, even with the additional observation that $\rho_\infty = \rho_{\text{im}}^{-1} \sigma_1^3 \rho_{\text{im}}$, where $\rho_{\text{im}} := \sigma_2 \sigma_1^{-1} \sigma_3 \sigma_2$; see Remark 4.6.

4I. Proof of Proposition 1.4. For the sets of singularities $(A_{14} \oplus A_2) \oplus A_3$, line 8, $(A_{14} \oplus A_2) \oplus A_2 \oplus A_1$, line 9, and $(A_{11} \oplus 2A_2) \oplus A_4$, line 17, the statement is an immediate consequence of [Degtyarev 2012, Theorem 7.48]. For $3A_6 \oplus A_1$, line 37, the only proper quotient of the commutant $[\pi_1, \pi_1] = \mathbb{Z}_7$ is trivial; hence, the group π'_1 of any perturbation D' is either abelian, $\pi'_1 = \mathbb{Z}_6$, or isomorphic to π_1 , the latter being the case if and only if D' is \mathbb{D}_{14} -special; see [Degtyarev 2008].

For the remaining set of singularities $(A_8 \oplus 3A_2) \oplus A_4 \oplus A_1$, line 33, we proceed as follows. Any proper perturbation factors through a maximal one, where a single singular point P of type A_m splits into two points $A_{m'}$, $A_{m''}$, so that $m' + m'' = m - 1$. Assume that $P \neq (0 : 0 : 1)$; see Section 4A. Then this point corresponds to a certain singular fiber x_i of the tetragonal model C and gives rise to a braid relation $\{\alpha_k, \alpha_{k+1}\}_{m+1} \uparrow \rho_i^{-1} = 1$; see Section 4B. For the new curve D' , this relation changes to $\{\alpha_k, \alpha_{k+1}\}_s \uparrow \rho_i^{-1} = 1$, where $s := \text{g.c.d.}(m' + 1, m'' + 1)$.

For any perturbation of any point P , we have $s = 3$ if P is of type A_8 or A_2 and the result is still of torus type, and $s = 1$ otherwise. Now, the statement is easily proved by repeating the computation with the braid $\mu_i = \sigma_k^{m+1}$ replaced with σ_k^s . (If it is the type A_4 point that is perturbed, one can use the alternative tetragonal model given by Remark 4.8.) \square

Acknowledgements

This paper was written during my sabbatical stay at *l'Institut des Hautes Études Scientifiques* and *Max-Planck-Institut für Mathematik*; I would like to thank these institutions for their support and hospitality. I am also grateful to M. Oka, who kindly clarified for me the results of [Oka and Pho 2002], to V. Kharlamov, who patiently introduced me to the history of the subject, and to the anonymous referee of the paper, who made a number of valuable suggestions and checked and confirmed the somewhat unexpected result of Section 4H.

References

- [Artal Bartolo et al. 2002] E. Artal Bartolo, J. Carmona Ruber, and J. I. Cogolludo Agustín, “On sextic curves with big Milnor number”, pp. 1–29 in *Trends in singularities*, edited by A. Libgober and M. Tibăr, Birkhäuser, Basel, 2002. MR 2003d:14034 Zbl 1018.14009
- [Artal-Bartolo et al. 2008] E. Artal-Bartolo, J. I. Cogolludo, and H.-o. Tokunaga, “A survey on Zariski pairs”, pp. 1–100 in *Algebraic geometry in East Asia* (Hanoi, 2005), edited by K. Konno and V. Nguyen-Khac, Adv. Stud. Pure Math. **50**, Math. Soc. Japan, Tokyo, 2008. MR 2009g:14030 Zbl 1141.14015
- [Artin 1947] E. Artin, “Theory of braids”, *Ann. of Math.* (2) **48** (1947), 101–126. MR 8,367a Zbl 0030.17703
- [Degtyarev 2008] A. Degtyarev, “Oka’s conjecture on irreducible plane sextics”, *J. Lond. Math. Soc.* (2) **78**:2 (2008), 329–351. MR 2009f:14054 Zbl 1158.14026
- [Degtyarev 2009a] A. Degtyarev, “Fundamental groups of symmetric sextics, II”, *Proc. Lond. Math. Soc.* (3) **99**:2 (2009), 353–385. MR 2011c:14091 Zbl 1173.14019
- [Degtyarev 2009b] A. Degtyarev, “On irreducible sextics with non-abelian fundamental group”, pp. 65–91 in *Singularities* (Toyama, 2007), edited by J.-P. Brasselet et al., Adv. Stud. Pure Math. **56**, Math. Soc. Japan, Tokyo, 2009. MR 2011e:14051 Zbl 1193.14037
- [Degtyarev 2012] A. Degtyarev, *Topology of algebraic curves: An approach via dessins d’enfants*, de Gruyter Studies in Mathematics **44**, de Gruyter, Berlin, 2012. MR 2952675 Zbl 06055189
- [Degtyarev and Oka 2009] A. Degtyarev and M. Oka, “A plane sextic with finite fundamental group”, pp. 93–108 in *Singularities* (Toyama, 2007), edited by J.-P. Brasselet et al., Adv. Stud. Pure Math. **56**, Math. Soc. Japan, Tokyo, 2009. MR 2011g:14075 Zbl 1193.14038
- [GAP Group 2008] GAP Group, *GAP – groups, algorithms, and programming, version 4.4.12*, 2008, <http://www.gap-system.org>.
- [van Kampen 1933] E. R. van Kampen, “On the fundamental group of an algebraic curve”, *Amer. J. Math.* **55**:1-4 (1933), 255–267. MR 1506962 Zbl 0006.41502
- [Libgober 2007a] A. Libgober, “Lectures on topology of complements and fundamental groups”, pp. 71–137 in *Singularity theory*, edited by D. Chéniot et al., World Scientific, Hackensack, NJ, 2007. MR 2008i:14043 Zbl 1125.14014
- [Libgober 2007b] A. Libgober, “Problems in topology of the complements to plane singular curves”, pp. 370–387 in *Singularities in geometry and topology* (Trieste, 2005), edited by J.-P. Brasselet et al., World Scientific, Hackensack, NJ, 2007. MR 2008c:14044 Zbl 1123.14018
- [Oka and Pho 2002] M. Oka and D. T. Pho, “Classification of sextics of torus type”, *Tokyo J. Math.* **25**:2 (2002), 399–433. MR 2003k:14030 Zbl 1062.14036

- [Orevkov 1999] S. Y. Orevkov, “Link theory and oval arrangements of real algebraic curves”, *Topology* **38**:4 (1999), 779–810. [MR 2000b:14066](#) [Zbl 0923.14032](#)
- [Persson 1985] U. Persson, “Double sextics and singular K -3 surfaces”, pp. 262–328 in *Algebraic geometry* (Barcelona, 1983), edited by E. Casas-Alvero et al., Lecture Notes in Math. **1124**, Springer, Berlin, 1985. [MR 87i:14036](#) [Zbl 0603.14023](#)
- [Shimada 2007] I. Shimada, “On the connected components of the moduli of polarized $K3$ surfaces”, preprint, 2007, <http://tinyurl.com/shimada-connK3>.
- [Yang 1996] J.-G. Yang, “Sextic curves with simple singularities”, *Tohoku Math. J. (2)* **48**:2 (1996), 203–227. [MR 98e:14026](#) [Zbl 0866.14014](#)
- [Zariski 1929] O. Zariski, “On the problem of existence of algebraic functions of two variables possessing a given branch curve”, *Amer. J. Math.* **51**:2 (1929), 305–328. [MR 1506719](#) [JFM 55.0806.01](#)

Received July 10, 2012. Revised December 29, 2012.

ALEX DEGTYAREV
DEPARTMENT OF MATHEMATICS
BILKENT UNIVERSITY
06800 ANKARA
TURKEY
degt@fen.bilkent.edu.tr

PACIFIC JOURNAL OF MATHEMATICS

msp.org/pjm

Founded in 1951 by E. F. Beckenbach (1906–1982) and F. Wolf (1904–1989)

EDITORS

V. S. Varadarajan (Managing Editor)
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
pacific@math.ucla.edu

Paul Balmer
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
balmer@math.ucla.edu

Don Blasius
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
blasius@math.ucla.edu

Vijayanthi Chari
Department of Mathematics
University of California
Riverside, CA 92521-0135
chari@math.ucr.edu

Daryl Cooper
Department of Mathematics
University of California
Santa Barbara, CA 93106-3080
cooper@math.ucsb.edu

Robert Finn
Department of Mathematics
Stanford University
Stanford, CA 94305-2125
finn@math.stanford.edu

Kefeng Liu
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
liu@math.ucla.edu

Jiang-Hua Lu
Department of Mathematics
The University of Hong Kong
Pokfulam Rd., Hong Kong
jhlu@maths.hku.hk

Sorin Popa
Department of Mathematics
University of California
Los Angeles, CA 90095-1555
popa@math.ucla.edu

Jie Qing
Department of Mathematics
University of California
Santa Cruz, CA 95064
qing@cats.ucsc.edu

Paul Yang
Department of Mathematics
Princeton University
Princeton NJ 08544-1000
yang@math.princeton.edu

PRODUCTION

Silvio Levy, Scientific Editor, production@msp.org

SUPPORTING INSTITUTIONS

ACADEMIA SINICA, TAIPEI
CALIFORNIA INST. OF TECHNOLOGY
INST. DE MATEMÁTICA PURA E APLICADA
KEIO UNIVERSITY
MATH. SCIENCES RESEARCH INSTITUTE
NEW MEXICO STATE UNIV.
OREGON STATE UNIV.

STANFORD UNIVERSITY
UNIV. OF BRITISH COLUMBIA
UNIV. OF CALIFORNIA, BERKELEY
UNIV. OF CALIFORNIA, DAVIS
UNIV. OF CALIFORNIA, LOS ANGELES
UNIV. OF CALIFORNIA, RIVERSIDE
UNIV. OF CALIFORNIA, SAN DIEGO
UNIV. OF CALIF., SANTA BARBARA

UNIV. OF CALIF., SANTA CRUZ
UNIV. OF MONTANA
UNIV. OF OREGON
UNIV. OF SOUTHERN CALIFORNIA
UNIV. OF UTAH
UNIV. OF WASHINGTON
WASHINGTON STATE UNIVERSITY

These supporting institutions contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its contents or policies.

See inside back cover or msp.org/pjm for submission instructions.

The subscription price for 2013 is US \$400/year for the electronic version, and \$485/year for print and electronic. Subscriptions, requests for back issues and changes of subscribers address should be sent to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163, U.S.A. The Pacific Journal of Mathematics is indexed by [Mathematical Reviews](#), [Zentralblatt MATH](#), [PASCAL CNRS Index](#), [Referativnyi Zhurnal](#), [Current Mathematical Publications](#) and the [Science Citation Index](#).

The Pacific Journal of Mathematics (ISSN 0030-8730) at the University of California, c/o Department of Mathematics, 798 Evans Hall #3840, Berkeley, CA 94720-3840, is published twelve times a year. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices. POSTMASTER: send address changes to Pacific Journal of Mathematics, P.O. Box 4163, Berkeley, CA 94704-0163.

PJM peer review and production are managed by EditFLOW[®] from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2013 Mathematical Sciences Publishers

PACIFIC JOURNAL OF MATHEMATICS

Volume 265 No. 2 October 2013

Singularity removability at branch points for Willmore surfaces	257
YANN BERNARD and TRISTAN RIVIÈRE	
On Bach flat warped product Einstein manifolds	313
QIANG CHEN and CHENXU HE	
On plane sextics with double singular points	327
ALEX DEGTYAREV	
A computational approach to the Kostant–Sekiguchi correspondence	349
HEIKO DIETRICH and WILLEM A. DE GRAAF	
Landau–Toeplitz theorems for slice regular functions over quaternions	381
GRAZIANO GENTILI and GIULIA SARFATTI	
On surgery curves for genus-one slice knots	405
PATRICK M. GILMER and CHARLES LIVINGSTON	
Characterizing abelian varieties by the reduction of the Mordell–Weil group	427
CHRIS HALL and ANTONELLA PERUCCA	
Variation of complex structures and the stability of Kähler–Ricci Solitons	441
STUART J. HALL and THOMAS MURPHY	
On crossed homomorphisms of the volume preserving diffeomorphism groups	455
RYOJI KASAGAWA	
Regularity at the boundary and tangential regularity of solutions of the Cauchy–Riemann system	491
TRAN VU KHANH and GIUSEPPE ZAMPIERI	
On the Steinberg character of a semisimple p -adic group	499
JU-LEE KIM and GEORGE LUSZTIG	