

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/45881879>

Hurwitz equivalence of braid monodromies and extremal elliptic surfaces

Article in *Proceedings of the London Mathematical Society* · November 2009

DOI: 10.1112/plms/pdr013 · Source: arXiv

CITATIONS

6

READS

22

1 author:



Alex Degtyarev

Bilkent University

81 PUBLICATIONS 748 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Lines in $K3$ -surfaces [View project](#)



Projective spaces in Fermat varieties [View project](#)

All content following this page was uploaded by [Alex Degtyarev](#) on 27 March 2014.

The user has requested enhancement of the downloaded file.

HURWITZ EQUIVALENCE OF BRAID MONODROMIES AND EXTREMAL ELLIPTIC SURFACES

ALEX DEGTYAREV

ABSTRACT. We discuss the equivalence between the categories of certain ribbon graphs and subgroups of the modular group Γ and use it to construct exponentially large families of not Hurwitz equivalent simple braid monodromy factorizations of the same element. As an application, we also obtain exponentially large families of *topologically* distinct algebraic objects such as extremal elliptic surfaces, real trigonal curves, and real elliptic surfaces.

1. INTRODUCTION

Strictly speaking, principal results of the paper concern extremal elliptic surfaces, see Subsection 1.3. However, we start with discussing a few applications to the braid monodromy, which seems to be a subject of more general interest.

1.1. Braid monodromy. Throughout the paper, we use the notation $[\cdot] = [\cdot]_G$ for the conjugacy class of an element $g \in G$ or a subgroup $H \subset G$ of a group G .

1.1.1. Definition. Given a group G , a (G -valued) *braid monodromy factorization* (*BM-factorization* for short) of length r is a sequence $\bar{\mathbf{m}} = (\mathbf{m}_1, \dots, \mathbf{m}_r)$ of elements of G . Two BM-factorizations are *strongly (Hurwitz) equivalent* if they are related by a finite sequence of *Hurwitz moves*

$$(\dots, \mathbf{m}_i, \mathbf{m}_{i+1}, \dots) \mapsto (\dots, \mathbf{m}_i^{-1} \mathbf{m}_{i+1} \mathbf{m}_i, \mathbf{m}_i, \dots)$$

and their inverse. Two BM-factorizations are *weakly equivalent* if they are related by a sequence of Hurwitz moves and their inverse and/or *global conjugation*

$$\bar{\mathbf{m}} = (\mathbf{m}_i) \mapsto g^{-1} \bar{\mathbf{m}} g := (g^{-1} \mathbf{m}_i g), \quad g \in G.$$

Often it is required that each element \mathbf{m}_i of a BM-factorization should belong to the union $\bigcup_j C_j$ of several conjugacy classes C_j fixed in advance. Thus, a \mathbb{B}_n -valued BM-factorization is called *simple* if each \mathbf{m}_i is conjugate to the Artin generator σ_1 , see Definition 5.1.3.

Note that we regard a braid monodromy as an *anti*-homomorphism, see 1.1.2 below. This convention explains the slightly unusual form of the Hurwitz moves and the fact that the order of multiplication is reversed in 1.1.3(1).

2000 *Mathematics Subject Classification.* 14J27, 14H57; 20F36, 11F06, 14P25.

Key words and phrases. Elliptic surface, braid monodromy, modular group, real trigonal curve, Lefschetz fibration, plane sextic, dessin d'enfant.

In this paper we mainly deal with the first nonabelian braid group \mathbb{B}_3 and the closely related groups $\tilde{\Gamma} := SL(2, \mathbb{Z})$ and $\Gamma := PSL(2, \mathbb{Z})$. A $\tilde{\Gamma}$ - or Γ -valued BM-factorization (\mathbf{m}_i) is called *simple* if each \mathbf{m}_i belongs to the corresponding conjugacy class $[[\mathbb{X}\mathbb{Y}]]$, see Subsection 2.1 for the notation. The classifications of simple BM-factorizations (up to weak/strong Hurwitz equivalence) in all three groups coincide, see Proposition 5.1.4.

1.1.2. A G -valued BM-factorization $\bar{\mathbf{m}} = (\mathbf{m}_1, \dots, \mathbf{m}_r)$ can be regarded as an anti-homomorphism $\langle \gamma_1, \dots, \gamma_r \rangle \rightarrow G$, $\gamma_i \mapsto \mathbf{m}_i$, $i = 1, \dots, r$. In this interpretation, Hurwitz moves generate the canonical action of the braid group \mathbb{B}_r on the free group $\langle \gamma_1, \dots, \gamma_r \rangle$, and the global conjugation represents the adjoint action of G on itself. Geometrically, anti-homomorphisms as above arise from locally trivial fibrations $X^\sharp \rightarrow B^\sharp$ over a punctured disk; then G is the (appropriately defined) mapping class group of the fiber over a fixed point $b \in \partial B^\sharp$ and $\langle \gamma_1, \dots, \gamma_r \rangle$ is a geometric basis for $\pi_1(B^\sharp, b)$. In this set-up, Hurwitz moves can be interpreted either as basis changes or as automorphisms of B^\sharp fixed on the boundary, see [3], and the topological classification of fibrations reduces to the purely algebraic classification of G -valued BM-factorizations up to weak Hurwitz equivalence. The best known examples are

- ramified coverings (the fiber is a finite set and $G = \mathbb{S}_n$, see [16]);
- algebraic or, more generally, pseudoholomorphic and Hurwitz curves in \mathbb{C}^2 (the fiber is a punctured plane and $G = \mathbb{B}_n$, see [29], [17], [6], [7], [21], [22], [20], [18], [24], [25]);
- (real) elliptic surfaces or, more generally, (real) Lefschetz fibrations of genus one (the fiber is an elliptic curve/topological torus and $G = \tilde{\Gamma}$, see [19], [28], [21], [5], [9], [13], [24], [25], [26]).

Last two subjects are quite popular and the reference lists are far from complete: I tried to cite the founding papers and a few recent results/surveys only.

Usually it is understood that the punctures of B^\sharp correspond to the singular fibers of a fibration $X \rightarrow B$ over a disk, the type of each singular fiber F being represented by the conjugacy class of the local monodromy about F . Thus, in the three examples above, simple BM-factorizations correspond to fibrations with simplest, not removable by a small deformation, singular fibers.

1.1.3. The following is a list of the most commonly used weak/strong equivalence invariants of a G -valued BM-factorization $\bar{\mathbf{m}}$:

- (1) the *monodromy at infinity* $\mathbf{m}_\infty(\bar{\mathbf{m}}) := \mathbf{m}_r \dots \mathbf{m}_1 \in G$ is a strong invariant; its conjugacy class $[[\mathbf{m}_\infty(\bar{\mathbf{m}})]]$ is a weak invariant;
- (2) the *monodromy group* $\text{Im}(\bar{\mathbf{m}}) := \langle \mathbf{m}_1, \dots, \mathbf{m}_r \rangle \subset G$ is a strong invariant; its conjugacy class $[[\text{Im}(\bar{\mathbf{m}})]]$ is a weak invariant;
- (3) for $G = SL(2, \mathbb{Z})$, the *transcendental lattice* $\mathcal{T}(\bar{\mathbf{m}})$, see Subsection 7.1 for the definition and generalizations, is a weak invariant;
- (4) for $G = \mathbb{B}_3$, define the (*affine*) *fundamental group* (see [29], [17])

$$\pi_1(\bar{\mathbf{m}}) := \langle \alpha_1, \alpha_2, \alpha_3 \mid \mathbf{m}_i(\alpha_j) = \alpha_j \text{ for } i = 1, \dots, r, j = 1, 2, 3 \rangle;$$

the homomorphism $\langle \alpha_1, \alpha_2, \alpha_3 \rangle \twoheadrightarrow \pi_1(\bar{\mathbf{m}})$ is a strong invariant; it depends on $\text{Im}(\bar{\mathbf{m}})$ only; the isomorphism class of the abstract group $\pi_1(\bar{\mathbf{m}})$ is a weak invariant; it depends on $[[\text{Im}(\bar{\mathbf{m}})]]$ only.

Due to Proposition 5.1.4, invariants (3) and (4) apply equally well to simple \mathbb{B}_3 -, $\tilde{\Gamma}$ -, and Γ -valued BM-factorizations. Note that often it is the group (4) that is the ultimate goal of computing the BM-factorization in the first place.

Geometrically, most important is the monodromy at infinity (1); in the set-up of 1.1.2, it corresponds to the monodromy along the boundary ∂B , and the BM-factorizations $\bar{\mathfrak{m}}$ with a given class $[[\mathfrak{m}_\infty(\bar{\mathfrak{m}})]] \subset G$ enumerate the extensions to B of a given fibration over ∂B . For this reason, a BM-factorization $\bar{\mathfrak{m}}$ is often regarded as a factorization of a given element $\mathfrak{m}_\infty(\bar{\mathfrak{m}})$ (which explains the term). The geometric importance of the extension problem, a number of partial results, and extensive experimental evidence give rise to the following two long standing questions.

1.1.4. Question. Is the weak/strong equivalence class of a simple \mathbb{B}_n -valued BM-factorization $\bar{\mathfrak{m}}$ determined by the monodromy at infinity $\mathfrak{m}_\infty(\bar{\mathfrak{m}})$? (Note that the length of $\bar{\mathfrak{m}}$ is determined by $\mathfrak{m}_\infty(\bar{\mathfrak{m}})$, see 5.1.5.)

1.1.5. Question. If two simple \mathbb{B}_n -valued BM-factorizations $\bar{\mathfrak{m}}_1, \bar{\mathfrak{m}}_2$ have the same monodromy at infinity and are weakly equivalent, are they also strongly equivalent? In other words, if a simple BM-factorization $\bar{\mathfrak{m}}$ is conjugated by an element of G commuting with $\mathfrak{m}_\infty(\bar{\mathfrak{m}})$, is the result strongly equivalent to $\bar{\mathfrak{m}}$?

The answer to Question 1.1.4 is in the affirmative if $n = 3$ and $\mathfrak{m}_\infty(\bar{\mathfrak{m}})$ is a central (see [21]) or, more generally, positive (with respect to the Artin basis, see [25]) element of \mathbb{B}_3 . Furthermore, for any n , two BM-factorizations sharing the same monodromy at infinity are known to be *stably equivalent*, see [18] or [20] for details.

The condition that $\bar{\mathfrak{m}}$ should be simple in Question 1.1.4 is crucial: in general, a BM-factorization is not unique. First example was essentially found in [29], and a great deal of other examples have been discovered since then. A few new examples are discussed in Subsections 5.5 and 5.6. In particular, we give a very simple, not computer aided, proof of the non-equivalence of the two BM-factorizations considered in [2].

1.2. Principal results. We answer Questions 1.1.4 and 1.1.5 in the negative for the braid group \mathbb{B}_3 (and related groups Γ and $\tilde{\Gamma}$, see Proposition 5.1.4). The inclusion $\mathbb{B}_3 \hookrightarrow \mathbb{B}_n$ implies a negative answer for the other braid groups as well, at least concerning the strong equivalence, see 5.1.7.

Let $T(k)$ be the number of isotopy classes of trees $\Xi \subset S^2$ with k trivalent vertices and $(k + 2)$ monovalent vertices (and no other vertices), see Section 4 and Corollary 4.2.2. Let $C(k) = \binom{2k}{k}/(k + 1)$ be the k -th Catalan number, and let $\tilde{T}(k) = (5k + 4)C(k)/(k + 2)$, see Subsection 4.2 and Corollary 4.2.2. Note that each of the three series grows faster than a^k for any $a < 4$. The first few values of $T(k)$ and $\tilde{T}(k)$ are listed in Table 1.

TABLE 1. A few values of $T(k)$ and $\tilde{T}(k)$

k	0	1	2	3	4	5	6	7	...	10	...	15
$T(k)$	1	1	1	1	4	6	19	49	...	1424	...	570285
$\tilde{T}(k)$	2	3	7	19	56	174	561	1859	...	75582	...	45052515

1.2.1. Theorem. For each integer $k \geq 0$, there is a set $\{\bar{\mathfrak{m}}_i\}$, $i = 1, \dots, \tilde{T}(k)$, of simple Γ -valued BM-factorizations of length $(k + 2)$ that share the same

– monodromy at infinity $\mathfrak{m}_\infty(\bar{\mathfrak{m}}_i) = (\mathbb{X}\mathbb{Y})^{-5k-4}$,

- transcendental lattice $\mathcal{T}(\bar{\mathfrak{m}}_i)$, see Example 7.2.3, and
- fundamental group $\pi_1(\bar{\mathfrak{m}}_i)$ (which is \mathbb{Z} for $k \geq 2$)

but are not strongly equivalent: the monodromy groups $\text{Im}(\bar{\mathfrak{m}}_i) \subset \Gamma$ are pairwise distinct subgroups of index $6(k+1)$.

1.2.2. Theorem. *For each k , the BM-factorizations $\bar{\mathfrak{m}}_i$ in Theorem 1.2.1 form $T(k)$ distinct weak equivalence classes: they are distinguished by the conjugacy classes $[\text{Im}(\bar{\mathfrak{m}}_i)]$ of the monodromy groups.*

Since $T(k) < \tilde{T}(k)$ for all $k \geq 0$, one has the following corollary.

1.2.3. Corollary. *For each integer $k \geq 0$, there is a pair of conjugate simple Γ -valued BM-factorizations of length $(k+2)$ that share the same monodromy at infinity $(\mathbb{X}\mathbb{Y})^{-5k-4}$ but are not strongly equivalent. \square*

Theorems 1.2.1 and 1.2.2 are proved in Subsection 5.2; the BM-factorizations in question are given by (5.2.2), and their \mathbb{B}_3 -valued counterparts are given by (5.3.1). The first example of weakly but not strongly equivalent \mathbb{B}_3 -valued BM-factorizations given by Corollary 1.2.3 has length two; it is as simple as

$$\bar{\mathfrak{m}}' = (\sigma_1^2 \sigma_2 \sigma_1^{-2}, \sigma_2), \quad \bar{\mathfrak{m}}'' = (\sigma_1 \sigma_2 \sigma_1^{-1}, \sigma_1^{-1} \sigma_2 \sigma_1),$$

see Example 5.3.3. The first example of non-equivalent BM-factorizations given by Theorem 1.2.2 has length six, see Example 5.3.2. In Subsection 5.4 we construct another example of not weakly equivalent BM-factorizations of length two; they also differ by the monodromy groups, which are of infinite index. A few other examples (not necessarily simple) are considered in Subsections 5.5 and 5.6.

1.3. Elliptic surfaces. Recall that an *extremal elliptic surface* can be defined as a Jacobian elliptic surface X of maximal Picard number, $\text{rk } NS(X) = h^{1,1}(X)$, and minimal Mordell-Weil rank, $\text{rk } MW(X) = 0$. (For an alternative description, in terms of singular fibers, see 2.2.3. Yet another characterization is the following: a Jacobian elliptic surface is extremal if and only if its transcendental lattice is positive definite, see [12].) Extremal elliptic surfaces are rigid (any small fiberwise equisingular deformation of such a surface X is isomorphic to X); they are defined over algebraic number fields.

In this paper, we mainly deal with elliptic surfaces with singular fibers of Kodaira types I_p and I_p^* . To shorten the statements, we call singular fibers of all other types, *i.e.*, Kodaira's II, III, IV and II^* , III^* , IV^* , *exceptional*. (These types are related to the exceptional simple singularities/Dynkin diagrams \mathbf{E}_6 , \mathbf{E}_7 , \mathbf{E}_8 .)

Given two elliptic surfaces X_1, X_2 , a fiberwise homeomorphism $\varphi: X_1 \rightarrow X_2$ is said to be *2-orientation preserving (reversing)* if it preserves (respectively, reverses) the complex orientation of the bases and the fibers of the two elliptic fibrations.

1.3.1. Theorem. *Two extremal elliptic surfaces without exceptional fibers are isomorphic if and only if they are related by a 2-orientation preserving fiberwise homeomorphism.*

Theorem 1.3.1 is not proved separately, as it is an immediate consequence of Theorem 2.5.3 below: the topological invariant distinguishing the surfaces is the conjugacy class in $\tilde{\Gamma}$ of the monodromy group of the homological invariant $\tilde{\mathfrak{h}}_X$, see 2.2.2. In fact, we show that appropriate subgroups of $\tilde{\Gamma}$ classify extremal elliptic surfaces without exceptional fibers, both analytically and topologically.

Two extensions of Theorem 1.3.1 to somewhat wider classes of surfaces are proved in Subsections 3.3 (see Remark 3.3.4) and 3.4.

As a by-product, we obtain exponentially large collections of non-homeomorphic elliptic surfaces sharing the same combinatorial type of singular fibers.

1.3.2. Theorem. *For each integer $k \geq 0$, there is a collection of $T(k)$ extremal elliptic surfaces that share the same combinatorial type of singular fibers, which is*

- $(k+2)I_1 \oplus I_{5k+4}^*$ if k is even, or
- $(k+2)I_1 \oplus I_{5k+4}$ if k is odd,

but are not related by a 2-orientation preserving fiberwise homeomorphism.

This theorem is proved in Subsection 4.3, and generalizations are discussed in Subsection 4.5. In fact, the surfaces were constructed in [8], and in [12] it was shown that they share as well such topological invariants as the transcendental lattice, see Example 7.2.3, and the fundamental group of the ramification locus.

The proof of Theorems 1.3.1 and 2.5.3 is based on an explicit computation of the monodromy group $\text{Im } \tilde{h}_X$ of an extremal elliptic surface X in terms of its skeleton Sk_X , see 2.2.4. In a sense, we show that Sk_X is $\text{Im } \tilde{h}_X$ (assuming that X has no type II^* singular fibers). As another consequence, we obtain an algebraic description of the reduced monodromy groups of such surfaces, see Subsection 3.5, and a few results (which may be known to the experts) on the subgroups of the modular group Γ ; to me, the most interesting seem Corollaries 3.2.5 and 3.6.2 describing the structure of subgroups and Proposition 4.4.1 characterizing monodromy groups of simple BM-factorizations.

1.4. Real trigonal curves and real elliptic surfaces. We consider a few other applications of the relation between ribbon graphs and subgroups of Γ , primarily to illustrate that some classification problems are wilder than they may seem.

Recall that the *Hirzebruch surface* is the geometrically ruled surface $\Sigma_k \rightarrow \mathbb{P}^1$, $k > 0$, with an exceptional section E of self-intersection $-k$. Up to isomorphism, there is a unique real structure (*i.e.*, anti-holomorphic involution) $\text{conj}: \Sigma_k \rightarrow \Sigma_k$ with nonempty real part $(\Sigma_k)_{\mathbb{R}} := \text{Fix conj}$. A curve $C \subset \Sigma_k$ is *real* if it is invariant under conj . A *trigonal curve* is a curve $C \subset \Sigma_k$ disjoint from E and intersecting each fiber of the ruling at three points. Such a curve is *generic* if all its singular fibers are of type I_1 (simple tangency of the curve and a fiber of the ruling). A generic curve is necessarily nonsingular.

1.4.1. Theorem. *For each integer $k \geq 0$, there is a collection of $T(k)$ generic real trigonal curves $C_i \subset \Sigma_{2k+2}$ such that all real parts $(C_i)_{\mathbb{R}} \subset (\Sigma_{2k+2})_{\mathbb{R}}$ are isotopic but the curves are not related by an equivariant 2-orientation preserving fiberwise auto-homeomorphism of Σ_{2k+2} preserving the orientation of the real part $\mathbb{P}_{\mathbb{R}}^1$ of the base of the ruling.*

Theorem 1.4.1 is proved in Subsection 6.2, and a generalization is discussed in Subsection 6.3. The real part of each curve C_i in Theorem 1.4.1 consists of a ‘long’ component L isotopic to $E_{\mathbb{R}}$ (see 6.1.3) and $(5k+4)$ ovals, all in the same connected component of $(\Sigma_{2k+2})_{\mathbb{R}} \setminus (L \cup E_{\mathbb{R}})$.

For each curve C_i as in Theorem 1.4.1, the double covering $X_i \rightarrow \Sigma_{2k+2}$ ramified at $C_i \cup E$ is a real Jacobian elliptic surface. Since the curves C_i are distinguished by the braid monodromy, one has the following corollary.

1.4.2. Corollary. *For each integer $k \geq 0$, there are two collections of $T(k)$ real Jacobian elliptic surfaces $X_i \rightarrow \mathbb{P}^1$ such that all real parts $(X_i)_{\mathbb{R}}$ are fiberwise homeomorphic but the surfaces are not related by an equivariant 2-orientation preserving fiberwise homeomorphism of Σ_{2k+2} preserving the orientation of the real part $\mathbb{P}_{\mathbb{R}}^1$ of the base of the elliptic pencil. \square*

In other words, each of the two collections consists of $T(k)$ pairwise non-isomorphic directed real Lefschetz fibrations of genus 1 in the sense of [26]. The real parts $(X_i)_{\mathbb{R}}$ can be described in terms of the *necklace diagrams*, see [26]: they are chains of $(5k+4)$ copies of the same stone, which is either $-\circ-$ or $-\square-$.

1.5. Contents of the paper. In Section 2, we introduce the basic objects and prove principal technical results relating extremal elliptic surfaces, 3-regular ribbon graphs, and geometric subgroups of Γ . Section 3 deals with a few generalizations of these results to wider classes of ribbon graphs/subgroups. In Section 4, we introduce *pseudo-trees*, which are ribbon graphs constructed from oriented rooted binary trees. It is this relation that is responsible for the exponential growth in most examples. Theorem 1.3.2 is proved here. In Sections 5 and 6, we prove the results concerning, respectively, simple BM-factorizations and real trigonal curves. Finally, in Section 7 we introduce the notion of transcendental lattice of a BM-factorization and consider a few examples.

2. ELLIPTIC SURFACES

In this section, we introduce basic notions and prove principal technical results: Corollary 2.3.5 and Theorem 2.4.5, establishing a connection between 3-regular ribbon graphs and geometric subgroups of Γ , and Theorems 2.5.2 and 2.5.3, relating extremal elliptic surfaces, their skeletons, and monodromy groups.

2.1. The modular group. Let $\mathcal{H} = \mathbb{Z}a \oplus \mathbb{Z}b$ be a rank 2 free abelian group with the skew-symmetric bilinear form $\wedge^2 \mathcal{H} \rightarrow \mathbb{Z}$ given by $a \cdot b = 1$. We fix the notation \mathcal{H} , a , b throughout the paper and define $\tilde{\Gamma} := SL(2, \mathbb{Z})$ as the group $\text{Sp } \mathcal{H}$ of symplectic auto-isometries of \mathcal{H} ; it is generated by the isometries $\mathbb{X}, \mathbb{Y}: \mathcal{H} \rightarrow \mathcal{H}$ given (in the basis $\{a, b\}$ above) by the matrices

$$\mathbb{X} = \begin{bmatrix} -1 & 1 \\ -1 & 0 \end{bmatrix}, \quad \mathbb{Y} = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix}.$$

One has $\mathbb{X}^3 = \text{id}$ and $\mathbb{Y}^2 = -\text{id}$. If $c = -a - b \in \mathcal{H}$, then \mathbb{X} acts *via*

$$(a, b) \xrightarrow{\mathbb{X}} (c, a) \xrightarrow{\mathbb{X}} (b, c) \xrightarrow{\mathbb{X}} (a, b).$$

The *modular group* $\Gamma := PSL(2, \mathbb{Z})$ is the quotient $\tilde{\Gamma}/\pm \text{id}$. We retain the notation \mathbb{X}, \mathbb{Y} for the generators of Γ . One has

$$\Gamma = \langle \mathbb{X} \mid \mathbb{X}^3 = 1 \rangle * \langle \mathbb{Y} \mid \mathbb{Y}^2 = 1 \rangle \cong \mathbb{Z}_3 * \mathbb{Z}_2.$$

A subgroup $H \subset \Gamma$ is called *geometric* if it is torsion free and of finite index. Since $\Gamma = \mathbb{Z}_3 * \mathbb{Z}_2$, the factors generated by \mathbb{X} and \mathbb{Y} , a subgroup $H \subset \Gamma$ is torsion free if and only if it is disjoint from the conjugacy classes $[\mathbb{X}]$ and $[\mathbb{Y}]$, or, equivalently, if both \mathbb{X} and \mathbb{Y} act freely on the quotient Γ/H .

Similarly, a subgroup $\tilde{H} \subset \tilde{\Gamma}$ is called *geometric* if it is torsion free and of finite index. A subgroup $\tilde{H} \subset \tilde{\Gamma}$ is torsion free if and only if $-\text{id} \notin \tilde{H}$ and the image of \tilde{H} in Γ is torsion free.

2.2. Extremal elliptic surfaces. In this subsection, we remind a few well known facts concerning Jacobian elliptic surfaces. The principal references are [14] or the original paper [19]. For more details concerning skeletons, we refer to [8].

A *Jacobian elliptic surface* is a compact complex surface X equipped with an elliptic fibration $\text{pr}: X \rightarrow B$ (*i.e.*, a fibration with all but finitely many fibers nonsingular elliptic curves) and a distinguished section $E \subset X$ of pr . (From the existence of a section it follows that X has no multiple fibers.) Throughout the paper we assume that surfaces are *relatively minimal*, *i.e.*, that fibers of the elliptic pencil contain no (-1) -curves.

2.2.1. Each nonsingular fiber of a Jacobian elliptic surface $\text{pr}: X \rightarrow B$ is an abelian group, and the multiplication by (-1) extends through the singular fibers of X . The quotient $X/\pm 1$ blows down to a geometrically ruled surface $\Sigma \rightarrow B$ over the same base B , and the double covering $X \rightarrow \Sigma$ is ramified over the exceptional section E of Σ and a certain *trigonal curve* $C \subset \Sigma$, *i.e.*, a curve disjoint from E and intersecting each generic fiber of the ruling at three points.

2.2.2. Denote by $B^\sharp \subset B$ the set of regular values of pr , and define the (*functional*) *j -invariant* $j_X: B \rightarrow \mathbb{P}^1$ as the analytic continuation of the function $B^\sharp \rightarrow \mathbb{C}^1$ sending each nonsingular fiber of pr to its classical j -invariant (divided by 12^3). The surface X is called *isotrivial* if $j_X = \text{const}$.

The monodromy $\tilde{h}_X: \pi_1(B^\sharp, b) \rightarrow \tilde{\Gamma} = \text{Sp } H_1(\text{pr}^{-1}(b))$, $b \in B^\sharp$, of the locally trivial fibration $\text{pr}^{-1}(B^\sharp) \rightarrow B^\sharp$ is called the *homological invariant* of X . Its reduction $h_X: \pi_1(B^\sharp) \rightarrow \Gamma$ is called the *reduced monodromy*; it is determined by the j -invariant. Together, j_X and \tilde{h}_X determine X up to isomorphism, and any pair (j, \tilde{h}) that agrees in the sense just described gives rise to a unique isomorphism class of Jacobian elliptic surfaces.

2.2.3. According to [23], a Jacobian elliptic surface X is extremal if and only if it satisfies the following conditions:

- (1) j_X has no critical values other than 0, 1, and ∞ ;
- (2) each point in $j_X^{-1}(0)$ has ramification index at most 3, and each point in $j_X^{-1}(1)$ has ramification index at most 2;
- (3) X has no singular fibers of types I_0^* , II, III, or IV.

2.2.4. The *skeleton* of a non-isotrivial elliptic surface $\text{pr}: X \rightarrow B$ is the embedded bipartite graph $\text{Sk}_X := j_X^{-1}[0, 1] \subset B$. The pull-backs of 0 and 1 are called \bullet - and \circ -vertices of Sk_X , respectively. (Thus, Sk_X is the *dessin d'enfants* of j_X in the sense of Grothendieck; however, we reserve the word ‘dessin’ for the more complicated graphs describing arbitrary, not necessarily extremal, surfaces, *cf.* Subsection 6.1.) *A priori*, j_X may have critical values in the open interval $(0, 1)$, hence the edges of Sk_X may meet at points other than \bullet - or \circ -vertices. However, by a small fiberwise equisingular deformation of X the skeleton Sk_X can be made *generic* in the sense that the edges of Sk_X meet only at \bullet - or \circ -vertices and the valency of each \bullet - (respectively, \circ -) vertex is ≤ 3 (respectively, ≤ 2).

The skeleton Sk_X of an extremal elliptic surface X is always generic. In addition, each region of Sk_X (*i.e.*, component of $B \setminus \text{Sk}_X$) is a topological disk; in particular, Sk_X is connected. Furthermore, each region contains a single critical point of j_X , the critical value being ∞ . Thus, in this case Sk_X can be regarded as an abstract ribbon graph: patching the cycles of Sk_X with disks, one recovers the topological surface B and ramified covering $j_X: B \rightarrow \mathbb{P}^1$; then, the analytic structure on B is

given by the Riemann existence theorem. It follows that the skeleton Sk_X of an extremal elliptic surface X determines its j -invariant $j_X: B \rightarrow \mathbb{P}^1$ (as an analytic function); hence the pair $(\text{Sk}_X, \tilde{h}_X)$ determines X .

2.2.5. The exceptional singular fibers of an elliptic surface X are in a one-to-one correspondence with the \bullet -vertices of Sk_X of valency $\neq 0 \pmod 3$ and its \circ -vertices of valency $\neq 0 \pmod 2$. Hence, if X is extremal and without exceptional fibers, all \bullet - and \circ -vertices of Sk_X are of valency 3 and 2, respectively. Since Sk_X is a bipartite graph, its \circ -vertices can be ignored, assuming that such a vertex is to be inserted at the middle of each edge connecting two \bullet -vertices. Under this convention, the skeleton of an extremal elliptic surface without exceptional fibers is a 3-regular ribbon graph. As explained above, each region of Sk_X is a disk containing a single singular fiber of X . Hence Sk_X is a strict deformation retract of B^\sharp , and the homological invariant can be regarded as an anti-homomorphism $\tilde{h}_X: \pi_1(\text{Sk}_X) \rightarrow \tilde{\Gamma}$. It is explained in [12] (see also Remark 2.5.6 below) that \tilde{h}_X can be encoded in terms of orientation of Sk_X .

2.3. Skeletons: another point of view. Following [12], we start with redefining a 3-regular ribbon graph as a set of ends of its edges. However, in the further exposition we will make no distinction between a graph in the sense of Definition 2.3.1 below and its geometric realization (defined in the obvious way). We will also redefine a few notions related to graphs (like connectedness, paths, *etc.*); each time, it is immediately obvious that the new notions are equivalent to their topological counterparts defined in terms of geometric realizations.

2.3.1. Definition. A *3-regular ribbon graph* is a collection $\text{Sk} = (\mathcal{E}, \text{op}, \text{nx})$, where $\mathcal{E} = \mathcal{E}_{\text{Sk}}$ is a finite set, $\text{op}: \mathcal{E} \rightarrow \mathcal{E}$ is a free involution, and $\text{nx}: \mathcal{E} \rightarrow \mathcal{E}$ is a free automorphism of order three. The orbits of op are called the *edges* of Sk , the orbits of nx are called its *vertices*, and the orbits of $\text{nx}^{-1} \text{op}$ are called its *faces* or *regions*. (Informally, op assigns to an end the other and of the same edge, and nx assigns the next end at the same vertex with respect to its cyclic order constituting the ribbon graph structure.)

A *pointed 3-regular ribbon graph* is a pair (Sk, e) , where $e \in \mathcal{E}_{\text{Sk}}$.

2.3.2. Remark. Alternatively, one can consider \mathcal{E}_{Sk} as the set of edges of Sk regarded as a bipartite ribbon graph, see 2.2.5. Then the orbits of op and nx represent, respectively, the \circ - and \bullet -vertices of Sk . Considering bipartite ribbon graph with the valency of \bullet - and \circ -vertices equal to (respectively, dividing) two given integers p and q , one can extend, almost literally, the material of this and next subsections (respectively, the generalizations found in Section 3) to the subgroups of the group $\langle x, y \mid x^p = y^q = 1 \rangle$. However, I do not know any interesting geometric applications of this group.

2.3.3. Given a 3-regular ribbon graph Sk , the set \mathcal{E}_{Sk} admits a canonical left Γ -action. To be precise, we define a homomorphism $\Gamma \rightarrow \mathbb{S}(\mathcal{E}_{\text{Sk}})$ to the group $\mathbb{S}(\mathcal{E}_{\text{Sk}})$ of permutations of \mathcal{E}_{Sk} via $\mathbb{X} \mapsto \text{nx}^{-1}$, $\mathbb{Y} \mapsto \text{op}$. According to this convention, the vertices, edges, and regions of Sk are the orbits of \mathbb{X} , \mathbb{Y} , and $\mathbb{X}\mathbb{Y}$, respectively. The graph Sk is *connected* if and only if the canonical Γ -action is transitive. A connected 3-regular ribbon graph is called a *3-skeleton*.

Given an element $e \in \mathcal{E}_{\text{Sk}}$, we denote by $\text{Stab } e \subset \Gamma$ its stabilizer. Stabilizers of all points of a 3-skeleton form a whole conjugacy class of subgroups of Γ ; it is denoted by $\llbracket \text{Stab } \text{Sk} \rrbracket$ and is called the *stabilizer* of Sk .

A *morphism* of 3-skeletons $\text{Sk}' = (\mathcal{E}', \text{op}, \text{nx})$ and $\text{Sk}'' = (\mathcal{E}'', \text{op}, \text{nx})$ is defined as a map $\mathcal{E}' \rightarrow \mathcal{E}''$ commuting with op and nx . In other words, it is a morphism of Γ -sets. A morphism of pointed 3-skeletons (Sk', e') and (Sk'', e'') is required, in addition, to take e' to e'' . The group of automorphisms of a 3-skeleton Sk is denoted Aut Sk ; we regard it as a subgroup of the symmetric group $\mathbb{S}(\mathcal{E}_{\text{Sk}})$.

The following two statements, although crucial for the sequel, are immediate consequences of the definitions.

2.3.4. Theorem. *The functors $(\text{Sk}, e) \mapsto \text{Stab } e$, $H \mapsto (\Gamma/H, H/H)$ establish an equivalence of the categories of*

- *pointed 3-skeletons and morphisms and*
- *geometric subgroups $H \subset \Gamma$ and inclusions.* \square

It follows that any morphism of 3-skeletons is a topological covering of their geometric realizations.

2.3.5. Corollary. *The maps $\text{Sk} \mapsto \llbracket \text{Stab Sk} \rrbracket$, $\llbracket H \rrbracket \mapsto \Gamma/H$ establish a canonical one-to-one correspondence between the sets of*

- *isomorphism classes of 3-skeletons and*
- *conjugacy classes of geometric subgroups $H \subset \Gamma$.* \square

If a 3-skeleton Sk is fixed, the isomorphism classes of pointed 3-skeletons (Sk, e) are naturally enumerated by the orbits of Aut Sk . Hence one has the following corollary, concerning properties of geometric subgroups.

2.3.6. Corollary. *The conjugacy class $\llbracket H \rrbracket$ of a geometric subgroup $H \subset \Gamma$ is in a one-to-one correspondence with the set of orbits of $\text{Aut}(\Gamma/H)$. Furthermore, there is an anti-isomorphism $\text{Aut}(\Gamma/H) = N(H)/H$, where $N(H)$ is the normalizer of H (acting on Γ/H by the right multiplication). \square*

2.3.7. Remark. Theorem 2.3.4, as well as its generalizations 3.2.1, 3.6.1 below, relating subgroups of Γ and ribbon graphs resemble the results of [4]. However, the two constructions differ: in [4], finite index subgroups of the congruence subgroup $\Gamma(2)$ are encoded using bipartite ribbon graphs with vertices of arbitrary valency. Our approach is closer to that of [5], where the modular j -function on a modular curve B (see [28] and Remark 2.5.5) is described in terms of a special triangulation of B . Theorem 2.4.5 below and its generalizations in Section 3 make the geometric relation between ribbon graphs and subgroups of Γ even more transparent.

2.4. Paths in a 3-skeleton. The treatment of paths found in [12] is not quite satisfactory for our purposes; we choose a slightly different approach here.

2.4.1. Definition. A *path* in a 3-skeleton $\text{Sk} = (\mathcal{E}, \text{op}, \text{nx})$ is a pair $\gamma = (e, w)$, where $e \in \mathcal{E}_{\text{Sk}}$ and w is a word in the alphabet $\{\text{op}, \text{nx}, \text{nx}^{-1}\}$. The *evaluation map* val sends a path $\gamma = (e, w)$ to the element $\text{val } \gamma \in \Gamma$ obtained by replacing $\text{op} \mapsto \mathbb{Y}$, $\text{nx}^{\pm 1} \mapsto \mathbb{X}^{\pm 1}$ in w and multiplying in Γ . The *starting* and *ending* points of γ are, respectively, $\gamma_0 := e \in \mathcal{E}_{\text{Sk}}$ and $\gamma_1 := (\text{val } \gamma)^{-1}e \in \mathcal{E}_{\text{Sk}}$. A path γ is a *loop* if $\gamma_0 = \gamma_1$. The *product* of two paths $\gamma' = (e', w')$ and $\gamma'' = (e'', w'')$ is defined whenever $\gamma_0'' = \gamma_1'$; it is $\gamma' \cdot \gamma'' := (e', w'w'')$, where $w'w''$ is the concatenation.

2.4.2. Remark. Intuitively, our definition of path represents the fact that, at each point $e \in \mathcal{E}_{\text{Sk}}$, one can choose among three directions: following the edge or walking around the vertex preserving or reversing the cyclic order. The inverse




FIGURE 1. A 3-skeleton Sk (black), auxiliary graph Sk° (bold grey), and space Sk^\bullet deformation equivalent to Sk (bold and light grey)

in the definition of γ_1 is due to the fact that the action of Γ is left rather than right, hence the order of elements of w should be reversed. (This is also one of the reasons why \mathbb{X} is defined to act *via* nx^{-1} .) Strictly speaking, what is defined is a geometric path (a chain of consecutive edges) in the auxiliary graph Sk° obtained from Sk by shortening each edge and replacing each vertex with a small circle (shown in bold grey lines in Figure 1). The vertices of Sk° are in a natural one-to-one correspondence with the elements of \mathcal{E}_{Sk} . When speaking about path homotopies, fundamental groups, *etc.*, we replace Sk° with the topological space Sk^\bullet obtained from Sk° by patching each circle with a disk (light grey in the figure) and consider the homomorphisms induced by the inclusion $\text{Sk}^\circ \hookrightarrow \text{Sk}^\bullet$ and the strict deformation retraction $\text{Sk}^\bullet \rightarrow \text{Sk}$.

The following two observations are also straightforward.

2.4.3. Lemma. *A path γ is a loop if and only if $\text{val } \gamma \in \text{Stab } \gamma_0$. Conversely, given $e \in \mathcal{E}_{\text{Sk}}$, any element of $\text{Stab } e$ has the form $\text{val } \gamma$ for some loop $\gamma = (e, w)$. \square*

2.4.4. Lemma. *Evaluation is multiplicative: $\text{val}(\gamma_1 \cdot \gamma_2) = \text{val } \gamma_1 \text{val } \gamma_2$. \square*

2.4.5. Theorem. *Given a pointed 3-skeleton (Sk, e) , the evaluation map restricts to a well defined isomorphism $\text{val}: \pi_1(\text{Sk}, e) \rightarrow \text{Stab } e$.*

Proof. Due to Lemmas 2.4.3 and 2.4.4, it suffices to show that val is well defined (*i.e.*, it takes equal values on homotopic loops) and $\text{Ker } \text{val} = \{1\}$. Both statements follow from comparing the cancellations in $\pi_1(\text{Sk}, e)$ and in Γ .

Since $\Gamma = \mathbb{Z}_3 * \mathbb{Z}_2$ is a free product, two words in $\{\mathbb{Y}, \mathbb{X}, \mathbb{X}^{-1}\}$ represent the same element of Γ if and only if they are obtained from each other by a sequence of cancellations of subwords of the form $\mathbb{Y}\mathbb{Y}$, $\mathbb{X}\mathbb{X}^{-1}$, $\mathbb{X}^{-1}\mathbb{X}$, $\mathbb{X}\mathbb{X}\mathbb{X}$, or $\mathbb{X}^{-1}\mathbb{X}^{-1}\mathbb{X}^{-1}$. The first three cancellations constitute the combinatorial definition of path homotopy in the auxiliary graph Sk° , see Remark 2.4.2: they correspond to cancelling an edge immediately followed by its inverse. The last two cancellations normally generate the kernel of the inclusion homomorphism $\pi_1(\text{Sk}^\circ, e) \rightarrow \pi_1(\text{Sk}^\bullet, e)$: they correspond to contracting circles in $\text{Sk}^\circ \subset \text{Sk}^\bullet$ to vertices of the original 3-skeleton Sk .

An alternative proof of the fact that val is well defined is given by Lemma 2.5.1 below, which provides an invariant geometric description of this map. \square

2.4.6. Corollary. *Any geometric subgroup $H \subset \Gamma$ (respectively, any geometric subgroup $\tilde{H} \subset \tilde{\Gamma}$) has index divisible by six, $[\Gamma : H] = 6k$ (respectively, divisible by twelve, $[\tilde{\Gamma} : \tilde{H}] = 12k$) and is isomorphic to a free group on $(k + 1)$ generators.*

Proof. Let $\text{Sk} = \Gamma/H$, see Theorem 2.3.4. Then $[\Gamma : H] = |\mathcal{E}_{\text{Sk}}|$. On the other hand, since Sk is a 3-regular graph, one has $|\mathcal{E}_{\text{Sk}}| = 6k$ and Sk has $2k$ vertices and $3k$ edges. Then $\chi(\text{Sk}) = -k$ and $\pi_1(\text{Sk})$ is a free group on $(k + 1)$ generators.

If $\tilde{H} \subset \tilde{\Gamma}$ is a geometric subgroup, then $\tilde{H} \not\cong -\text{id}$ and the projection $\tilde{H} \rightarrow \Gamma$ is an isomorphism onto its image, which is a geometric subgroup of Γ . \square

2.4.7. Remark. The universal covering of a 3-skeleton Sk is a 3-regular tree; hence it is the Farey tree. The automorphism group $\text{Aut } F$ of the Farey tree F can be identified with Γ : it is generated by the rotations about a vertex or the center of an edge. Thus, geometrically, $\text{Sk} = F/H$ for a finite index subgroup $H \subset \text{Aut } F$ acting freely on F , and Theorem 2.4.5 becomes a well known property of topological coverings. If the action of H on F is not free, one needs to consider the orbifold fundamental group $\pi_1^{\text{orb}}(F/H)$, see Subsection 3.2 below. If $[\Gamma : H] = \infty$, the quotient F/H is an infinite graph, see Subsections 3.1 and 3.6.

2.5. The homological invariant. Fix a Jacobian elliptic surface $\text{pr}: X \rightarrow B$ without exceptional fibers and let $\text{Sk} = \text{Sk}_X$ be the skeleton of X . Assume that Sk is generic, hence 3-regular. Consider the double covering $X \rightarrow \Sigma$ ramified at $C \cup E$, see 2.2.1. Pick a vertex v of Sk , let F_v be the fiber of X over v , and let \bar{F}_v be its projection to Σ . Then, F_v is the double covering of \bar{F}_v ramified at $\bar{F}_v \cap (C \cup E)$ (the three black points in Figure 2 and ∞).




FIGURE 2. The basis in $H_1(F_v)$

Recall that the three points of intersection $\bar{F}_v \cap C$ are in a canonical one-to-one correspondence with the three ends constituting v , see [8]. Choose one of the ends (a *marking* at v in the terminology of [8]) and let $\{\alpha_1, \alpha_2, \alpha_3\}$ be the canonical basis for the group $\pi_1(\bar{F}_v \setminus (C \cup E))$ defined by this end (see [8] and Figure 2; unlike [8], we take for the reference point the zero section of Σ , which is well defined in the presence of a trigonal curve; this choice removes the ambiguity in the definition of canonical basis). Then $H_1(F_v) = \pi_1(F_v)$ is generated by the lifts $a = \alpha_2 \alpha_1$ and $b = \alpha_1 \alpha_3$ (the two grey cycles in the figure). To be precise, one needs to choose one of the two pull-backs of the zero section and take it for the reference point for $\pi_1(F_v)$ (the grey point at the center of the figure). Thus, a choice of an end at v gives rise to an isometry $H_1(F_v) = \mathcal{H}$, which is canonical up to $\pm \text{id}$.

Now, consider a copy F_e of F_v for each end $e \in v$ and identify it with \mathcal{H} using e as the marker. (Alternatively, one can assume that a separate fiber is chosen over each vertex of the auxiliary graph Sk° , see Remark 2.4.2.) Under this identification, the monodromy $\tilde{\mathfrak{h}}_\gamma: H_1(F_{\gamma_0}) \rightarrow H_1(F_{\gamma_1})$ of the locally trivial fibration $\text{pr}^{-1}(\text{Sk}) \rightarrow \text{Sk}$ along a path γ in Sk reduces to a well defined element $\mathfrak{h}_\gamma \in \Gamma$.

2.5.1. Lemma. *In the notation above, one has $\mathfrak{h}_\gamma = (\text{val } \gamma)^{-1}$.*

Proof. Since both maps $\gamma \mapsto \mathfrak{h}_\gamma$ and $\gamma \mapsto (\text{val } \gamma)^{-1}$ reverse products, it suffices to prove the assertion for a path of length 1, *i.e.*, for a single edge of Sk° .

Circumventing a vertex of the original skeleton Sk in the positive direction is the change of basis induced by a change of the marker (rotation through $-2\pi/3$ about the center in Figure 2); its transition matrix is $\mathbb{X}^{-1} = (\text{val } \text{nx})^{-1}$. Following an edge of Sk is a lift of the monodromy $m_{1,1}$ in [8]: during the monodromy, the black ramification point surrounded by α_1 crosses the segment connecting the ramification points surrounded by α_2 and α_3 ; modulo $\pm \text{id}$, the corresponding linear operator is given by $\mathbb{Y} = (\text{val } \text{op})^{-1}$. \square

Let v be a vertex of Sk and let $e \in v$. We will use the notation $\pi_1(B^\sharp, e)$ for the group $\pi_1(B^\sharp, v)$, meaning that the fiber F_v is identified with \mathcal{H} using e as a marker. Thus, we will speak about the reduced monodromy $\mathfrak{h}_X: \pi_1(B^\sharp, e) \rightarrow \Gamma$.

2.5.2. Theorem. *Let X be an extremal elliptic surface without exceptional fibers, and let e be a representative of a vertex of Sk_X . Then the reduced monodromy $\mathfrak{h}_X: \pi_1(B^\sharp, e) \rightarrow \Gamma$ takes values in $\text{Stab } e$, both maps in the diagram*

$$\pi_1(\text{Sk}_X, e) \xrightarrow{\text{in}_*} \pi_1(B^\sharp, e) \xrightarrow{\mathfrak{h}_X} \text{Stab } e \subset \Gamma$$

are (anti-)isomorphisms, and the composed map is given by $\gamma \mapsto (\text{val } \gamma)^{-1}$.

Proof. Since Sk_X is a strict deformation retract of B^\sharp , see 2.2.5, the inclusion homomorphism $\text{in}_*: \pi_1(\text{Sk}_X) \rightarrow \pi_1(B^\sharp)$ is an isomorphism. The rest follows from Lemma 2.5.1 and Theorem 2.4.5. \square

2.5.3. Theorem. *The map $X \rightarrow [\text{Im } \tilde{\mathfrak{h}}_X]$ establishes a bijection between the set of isomorphism classes of extremal elliptic surfaces without exceptional fibers and the set of conjugacy classes of geometric subgroups of $\tilde{\Gamma}$.*

Proof. It suffices to show that a subgroup $\tilde{H} \subset \tilde{\Gamma}$ defines a unique extremal elliptic surface. Since \tilde{H} is geometric, in particular $-\text{id} \notin \tilde{H}$, the projection $\tilde{\Gamma} \rightarrow \Gamma$ induces an isomorphism of \tilde{H} to a geometric subgroup $H \subset \Gamma$. The latter determines a skeleton $\text{Sk} \subset B$, hence a j -invariant $j_X: B \rightarrow \mathbb{P}^1$ and corresponding reduced monodromy $\mathfrak{h}_X: \pi_1(B^\sharp) \rightarrow H$. Then, the inverse isomorphism $H \rightarrow \tilde{H}$ is merely a lift of \mathfrak{h}_X to a homological invariant $\tilde{\mathfrak{h}}_X$; together with j_X , it defines a unique isomorphism class of Jacobian elliptic surfaces, which are necessarily extremal due to [23], see 2.2.3. \square

Since the conjugacy class of the monodromy group of a fibration is obviously invariant under fiberwise homeomorphisms, Theorem 2.5.3 implies Theorem 1.3.1 in the introduction.

2.5.4. Remark. One can easily see that two extremal elliptic surfaces without exceptional singular fibers are anti-isomorphic if and only if their monodromy subgroups are conjugated by an element of $GL(2, \mathbb{Z}) \setminus \tilde{\Gamma}$. (This conjugation results in a homeomorphism of the skeletons reversing the cyclic order at each vertex.) In other words, surfaces are anti-isomorphic if and only if they are related by a 2-orientation reversing homeomorphism.

2.5.5. Remark. The inverse map sending a geometric subgroup $H \subset \tilde{\Gamma}$ to an extremal elliptic surface in Theorem 2.5.3 is equivalent to Shioda's construction [28] of modular elliptic surfaces, where the base B of the elliptic fibration is obtained as the quotient $\{z \in \mathbb{C} \mid \text{Im } z > 0\}/H$ and the j -invariant j_X is the descent of the modular j -invariant. A generalization of the results of this section to arbitrary finite

index subgroups of Γ is considered in Subsections 3.2 and 3.3, see Remark 3.3.4; such subgroups correspond to skeletons with monovalent \bullet - and \circ -vertices allowed. For a further generalization to arbitrary subgroups, see Subsections 3.1 and 3.6; finitely generated subgroups can still be encoded by finite ribbon graphs.

2.5.6. Remark. In [12] it is shown that, for an extremal elliptic surface X without exceptional singular fibers, the homological invariant \tilde{h}_X admits a simple geometric description in terms of an orientation of Sk_X : one defines the value $\tilde{h}_X(\gamma)$ on a loop γ in Sk_X to be $\pm(\text{val } \gamma)^{-1} \in \tilde{\Gamma}$, depending on the parity of the number of edges travelled by γ in the opposite direction. This correspondence is not one-to-one, as distinct orientations may give rise to the same homological invariant.

3. GENERALIZATIONS

In this section, we generalize some results of Section 2 to arbitrary subgroups of Γ : finitely generated subgroups can still be encoded by finite graphs. Proofs are merely sketched, as they repeat, almost literally, those in Section 2. The material of this section is not used in the proofs of the principal results of the paper stated in the introduction.

3.1. Infinite skeletons. In order to study subgroups of Γ of infinite index, we modify Definition 2.3.1 and define a *generalized 3-regular ribbon graph* as a triple $\text{Sk} = (\mathcal{E}_{\text{Sk}}, \text{op}, \text{nx})$, where \mathcal{E}_{Sk} is a set (not necessarily finite) and op and nx are free automorphisms of \mathcal{E}_{Sk} of order 2 and 3, respectively. A *generalized 3-skeleton* is a connected generalized 3-regular ribbon graph.

All notions introduced in Subsections 2.3 and 2.4 and most statements proved there extend to the general case with obvious changes. We restate Theorems 2.3.4 and 2.4.5.

3.1.1. Theorem. *The functors $(\text{Sk}, e) \mapsto \text{Stab } e$, $H \mapsto (\Gamma/H, H/H)$ establish an equivalence of the categories of*

- *pointed generalized 3-skeletons and morphisms and*
- *torsion free subgroups $H \subset \Gamma$ and inclusions.* \square

3.1.2. Theorem. *Given a pointed generalized 3-skeleton (Sk, e) , the evaluation map restricts to a well defined isomorphism $\text{val}: \pi_1(\text{Sk}, e) \rightarrow \text{Stab } e$.* \square

A generalized 3-skeleton Sk is called *almost contractible* if the group $\pi_1(\text{Sk})$ is finitely generated. Under Theorem 3.1.1, almost contractible skeletons correspond to finitely generated torsion free subgroups.

3.1.3. Proposition. *There is a one-to-one correspondence between the sets of*

- (1) *conjugacy classes of proper finitely generated torsion free subgroups $H \subset \Gamma$,*
- (2) *almost contractible 3-skeletons with at least one cycle, and*
- (3) *connected finite ribbon graphs with all vertices of valency 3 or 1 and such that distinct monovalent vertices are adjacent to distinct trivalent vertices.*

Under this correspondence $H \leftrightarrow \text{Sk} \leftrightarrow \text{Sk}^c$ one has (anti-)isomorphisms $N(H)/H = \text{Aut } \text{Sk} = \text{Aut } \text{Sk}^c$ and $H = \pi_1(\text{Sk}) = \pi_1(\text{Sk}^c)$; in fact, Sk^c is embedded to Sk as an induced subgraph and a strict deformation retract.

The finite ribbon graph Sk^c corresponding to an almost contractible 3-skeleton Sk under Proposition 3.1.3 is called the *compact part* of Sk . In the drawings, the

monovalent vertices of Sk^c (those that are to be extended to ‘half’ Farey trees) are represented by triangles \triangle , *cf.* Figure 8 in Subsection 5.4. The last condition in 3.1.3(3) is the requirement that Sk^c should admit no further contraction to a subgraph with all vertices of valency 3 or 1. This condition makes Sk^c canonical.

Proof. Each almost contractible 3-skeleton Sk contains an induced subgraph Sk' such that $\text{Sk} \setminus \text{Sk}'$ is a forest: one can pick a finite collection of loops representing a basis for $\pi_1(\text{Sk})$ and take for Sk' the induced subgraph generated by all vertices contained in at least one of the loops. (The notation $\text{Sk} \setminus \text{Sk}'$ stands for the induced subgraph generated by the vertices of Sk that are not in Sk' .) The complement $\text{Sk} \setminus \text{Sk}'$ is a finite disjoint union of *infinite branches*, each infinite branch being a tree with one bivalent vertex and all other vertices trivalent. Unless Sk is the Farey tree itself (corresponding to the trivial subgroup of Γ), each infinite branch is contained in a unique maximal one. The maximal infinite branches are pairwise disjoint, and contracting each such branch to its only bivalent vertex produces the compact part Sk^c as in the statement, the monovalent vertices of Sk^c corresponding to the maximal infinite branches contracted. (The last condition in 3.1.3(3) is due to the fact that, if two monovalent vertices u_1, u_2 were adjacent to the same vertex v then, together with v , the two infinite branches represented by u_1 and u_2 would form a larger infinite branch.)

Since the construction is canonical, any automorphism of Sk preserves Sk^c and hence restricts to an automorphism of Sk^c . Conversely, any automorphism of Sk^c extends to a unique automorphism of Sk : the uniqueness is due to the fact that *ribbon* graphs are considered; once an automorphism of such a graph fixes a vertex v and an edge adjacent to v , it is the identity. \square

3.2. Skeletons with monovalent vertices. As another generalization, we lift the requirement that op and nx should be free and define a $(3, 1)$ -*ribbon graph* as a triple $\text{Sk} = (\mathcal{E}_{\text{Sk}}, \text{op}, \text{nx})$, where \mathcal{E}_{Sk} is a finite set and op and nx are automorphisms of \mathcal{E}_{Sk} of order 2 and 3, respectively. A $(3, 1)$ -*skeleton* is a connected $(3, 1)$ -ribbon graph. Thus, a $(3, 1)$ -skeleton is allowed to have monovalent \bullet -vertices (which are the one element orbits of nx) and ‘hanging edges’ (one element orbits of op); the latter are represented in the figures by monovalent \circ -vertices attached to these edges, *cf.* Figure 3 below.

As above, all notions introduced in Subsections 2.3 and 2.4 extend to the case of $(3, 1)$ -skeletons. Theorem 2.3.4 takes the following form.

3.2.1. Theorem. *The functors $(\text{Sk}, e) \mapsto \text{Stab } e$, $H \mapsto (\Gamma/H, H/H)$ establish an equivalence of the categories of*

- *pointed $(3, 1)$ -skeletons and morphisms and*
- *finite index subgroups $H \subset \Gamma$ and inclusions.* \square

3.2.2. Denote by $D_1^2 \cong D^2$, $D_2^2 \cong \mathbb{P}_{\mathbb{R}}^1$, and D_3^2 the CW-complexes obtained by attaching a single 2-cell D^2 to a circle S^1 via a map $\partial D^2 \rightarrow S^1$ of degree 1, 2, or 3, respectively. Given $e \in \mathcal{E}_{\text{Sk}}$, define the *orbifold fundamental group* $\pi_1^{\text{orb}}(\text{Sk}, e)$ as the fundamental group $\pi_1(\text{Sk}^\bullet, e)$, where the space Sk^\bullet is obtained from Sk by replacing a neighborhood of each trivalent \bullet -vertex, monovalent \circ -vertex, or monovalent \bullet -vertex with a copy of D_1^2 , D_2^2 , or D_3^2 , respectively, *cf.* Figure 3. (Note that $\pi_1^{\text{orb}}(\text{Sk}, e)$ is indeed the orbifold fundamental group, with the orbifold structure given by declaring each monovalent \circ - or \bullet -vertex a ramification point of ramification index 2 or 3, respectively. With this convention, the universal

covering of Sk is again the Farey tree, *cf.* Remark 2.4.7.) Contracting a maximal tree not containing a monovalent vertex, one establishes a homotopy equivalence between Sk^\bullet and a wedge of circles and copies of D_2^2 and D_3^2 . Hence, $\pi_1^{\text{orb}}(\text{Sk}, e)$ is a free product

$$(3.2.3) \quad \pi_1^{\text{orb}}(\text{Sk}, e) = \otimes_{n_0} \mathbb{Z} * \otimes_{n_2} \mathbb{Z}_2 * \otimes_{n_3} \mathbb{Z}_3,$$

where n_2 and n_3 are the numbers of monovalent \circ - and \bullet -vertices, respectively, and $n_0 = 1 - \chi(\text{Sk}) = 1 - \chi(\text{Sk}^\bullet)$. Observe that $|\mathcal{E}_{\text{Sk}}| = 6n_0 + 3n_2 + 4n_3 - 6$ (a simple combinatorial computation of the Euler characteristic).




FIGURE 3. A $(3, 1)$ -skeleton Sk (black), auxiliary graph Sk° (bold grey), and space Sk^\bullet (bold and light grey)

Definition 2.4.1 of paths, loops, and the evaluation map extends literally to the case of $(3, 1)$ -skeletons. Thus, we are speaking about geometric paths in the auxiliary graph Sk° obtained by fattening the vertices of Sk as shown in Figure 3. (Note though that we disregard the direction of a path along the single edge replacing a \circ -vertex and the adjacent edge of Sk .) It is straightforward that $\pi_1^{\text{orb}}(\text{Sk})$ can be defined as the group of loops modulo an appropriate equivalence relation. Next statement is proved similar to Theorem 2.4.5.

3.2.4. Theorem. *Given a pointed $(3, 1)$ -skeleton (Sk, e) , the evaluation map val factors through a well defined isomorphism $\text{val}: \pi_1^{\text{orb}}(\text{Sk}, e) \rightarrow \text{Stab } e$. \square*

3.2.5. Corollary. *Any finite index subgroup $H \subset \Gamma$ is a free product (3.2.3), and one has $[\Gamma : H] = 6n_0 + 3n_2 + 4n_3 - 6$. \square*

3.3. Extremal elliptic surfaces without type II^* fibers. Using the concept of $(3, 1)$ -skeleton introduced in the previous section and the description of the braid monodromy of the ramification locus found in [8] (the monodromy $l_1(2) \mapsto \mathbb{Y}\mathbb{X}^{-1}\mathbb{Y}$ and $l_1(3) \mapsto \mathbb{Y}$ for monovalent \bullet - and \circ -vertices, respectively; as in Subsection 2.5, the homomorphism $\mathbb{B}_3 \rightarrow \Gamma$ is given by (5.1.1) below), one arrives at the following generalization of Theorem 2.5.2.

3.3.1. Theorem. *Let X be an extremal elliptic surface without type II^* fibers, and let $e \in \mathcal{E}$ be a representative of a vertex of the skeleton Sk_X . Then the reduced monodromy $\mathfrak{h}_X: \pi_1(B^\sharp, e) \rightarrow \Gamma$ factors as follows:*

$$\pi_1(B^\sharp, e) \longrightarrow \pi_1^{\text{orb}}(\text{Sk}_X, e) \xrightarrow{\cong} \text{Stab } e \subset \Gamma$$

where the rightmost anti-isomorphism is the map $\gamma \mapsto (\text{val } \gamma)^{-1}$. \square

3.3.2. Remark. In the presence of monovalent vertices, Sk_X is no longer a subspace of B^\sharp . The first arrow in Theorem 3.3.1 is the composition of the homomorphisms induced by the strict deformation retraction $B^\sharp \rightarrow \text{Sk}'$ and the inclusion $\text{Sk}' \hookrightarrow \text{Sk}^\bullet$, where Sk' is obtained from Sk° , see Figure 3, by patching with disks the circles surrounding the *trivalent* \bullet -vertices only.

3.3.3. Corollary. *The map $X \rightarrow [\text{Im } \tilde{\mathfrak{h}}_X]$ establishes a bijection between the set of isomorphism classes of extremal elliptic surfaces without type II^* or III^* fibers and the set of conjugacy classes of finite index subgroups $\tilde{H} \subset \tilde{\Gamma}$ such that $-\text{id} \notin \tilde{H}$.*

Proof. Let X be a surface as in the statement, let $\tilde{H} = \text{Im } \tilde{\mathfrak{h}}_X \subset \tilde{\Gamma}$ (with respect to some base point in B^\sharp), and let $H = \text{Im } \mathfrak{h}_X \subset \Gamma$ be the projection of \tilde{H} to Γ . Under the assumptions, Sk_X has no \circ -vertices and hence $\pi_1^{\text{orb}}(\text{Sk}_X) = H$ is a free product of copies of \mathbb{Z} and \mathbb{Z}_3 only. Furthermore, each order 3 generator of H represents the monodromy about a type IV^* singular fiber of X , see 2.2.3(3), and hence lifts to an order 3 element of \tilde{H} . Thus, the projection $\tilde{H} \rightarrow H$ admits a section and hence is an isomorphism. The rest of the proof follows that of Theorem 2.5.3. \square

3.3.4. Remark. Corollary 3.3.3 covers Shioda's construction [28] to full extent and generalizes Theorem 1.3.1 to surfaces with type IV^* fibers allowed. Apparently, considering the homological invariant itself rather than just its image, one can further generalize Theorem 1.3.1 to type III^* singular fibers. The special case of rational base is considered in Theorem 3.4.1 below.

3.3.5. Remark. Surprisingly, type II^* singular fibers do not fit into the approach of this paper at all, as they are represented by bivalent \bullet -vertices of the skeleton, *i.e.*, orbits of nx of length two. Possibly, such skeletons can be treated as homogeneous spaces of $\tilde{\Gamma}$ rather than Γ , but the precise statements are not quite clear at the moment. An attempt of considering such more general skeletons is made in [12].

3.4. The case of rational base. In this subsection, we assume that the base B of an elliptic fibration $X \rightarrow B$ is rational, $B \cong \mathbb{P}^1$. In this case, the homological invariant $\tilde{\mathfrak{h}}_X$ (lifting a given reduced monodromy \mathfrak{h}_X) can be defined in terms of a *type specification* of X , *i.e.*, a choice of one of the two possible types (whose local monodromies differ by $-\text{id}$) of each singular fiber. Moreover, the types of all but one singular fibers can be chosen arbitrary, whereas the type of the remaining fiber is determined by the requirement that the total multiplicity of all singular fibers, which equals the topological Euler characteristic $\chi(X)$, should be divisible by 12. (The multiplicities of the two lifts of a given element of Γ differ by 6, *cf.* 5.1.2.)

If X is extremal and has no type II^* singular fibers, its type specification can be described in terms of the reduced monodromy group $H = \text{Im } \mathfrak{h}_X$. Indeed, in view of condition 2.2.3(3), the types of the exceptional fibers of X are fixed. The non-exceptional singular fibers are in a one-to-one correspondence with the regions of Sk_X , equivalently, with the orbits of $\mathbb{X}\mathbb{Y}$, equivalently, with the H -conjugacy classes of maximal unipotent subgroups of H , and a type specification consists in assigning a lift $\langle \pm g^{-1}(\mathbb{X}\mathbb{Y})^n g \rangle \subset \tilde{\Gamma}$ to each such conjugacy class $[[g^{-1}(\mathbb{X}\mathbb{Y})^n g]]_H$.

3.4.1. Theorem. *Two extremal elliptic surfaces X_1, X_2 over the rational base $B = \mathbb{P}^1$ and without type II^* singular fibers are isomorphic if and only if they are related by a 2-orientation preserving fiberwise homeomorphism.*

Proof. The 'only if' part is obvious. For the 'if' part, it suffices to notice that a 2-orientation preserving homeomorphism $X_1 \rightarrow X_2$ induces an orientation preserving homeomorphism $B_1 \rightarrow B_2$ taking punctures to punctures, commuting with the homological invariants $\pi_1(B_1^\sharp) \rightarrow \Gamma \leftarrow \pi_1(B_2^\sharp)$ (and hence taking H_1 to H_2) and preserving the type specification (as distinct types of singular elliptic fibers differ topologically, for example by the local monodromy). Hence, X_1 and X_2 are isomorphic. \square

3.4.2. Remark. The extremality condition in Theorem 3.4.1 can be relaxed by replacing 2.2.3(3) by the requirement that the surface should have no singular fibers of type I_0^* , II^* , or IV . In this case, a type specification would also choose a lift $\langle \pm g^{-1}\mathbb{X}g \rangle$ for each conjugacy class $[[\langle g^{-1}\mathbb{X}g \rangle]]_H$ of order 3 subgroups of H (monovalent \bullet -vertices) and a lift $\langle \pm g^{-1}\mathbb{Y}g \rangle$ for each conjugacy class $[[\langle g^{-1}\mathbb{Y}g \rangle]]_H$ of order 2 subgroups of H (monovalent \circ -vertices).

3.4.3. Remark. The combinatorial type of singular fibers of an extremal (or more general as in Remark 3.4.2) elliptic surface X is determined by its type specification and the following combinatorial information about its skeleton Sk_X : the numbers of monovalent \bullet - and \circ -vertices and the shapes of the regions of Sk_X . Each monovalent \bullet - (respectively, \circ -) vertex gives rise to a singular fiber of type II or IV^* (respectively, III or III^*), and each n -gonal region gives rise to a singular fiber of type I_n or I_n^* . There are large numbers of skeletons sharing these data; some examples are considered in Subsections 4.3, 4.5, and 5.6 below.

3.5. The monodromy group of an elliptic surface. For an elliptic surface X , introduce the following fiber counts:

- n_{II} is the number of fibers of type II or IV^* ;
- n_{III} is the number of fibers of type III or III^* ;
- n_{IV} is the number of fibers of type IV or II^* ;
- t is the number of fibers of type I_p^* , $p \geq 0$, II^* , III^* , or IV^* .

Let, further, $\chi(X)$ be the topological Euler characteristic of X .

3.5.1. Theorem. *Let X be an extremal elliptic surface without type II^* singular fibers. Then the reduced monodromy group $\text{Im } \mathfrak{h}_X \subset \Gamma$ is a subgroup of index $\chi(X) - 6t - 2n_{II} - 3n_{III}$ isomorphic to the free product*

$$\otimes_n \mathbb{Z} * \otimes_{n_{III}} \mathbb{Z}_2 * \otimes_{n_{II}} \mathbb{Z}_3,$$

where $n = \frac{1}{6}\chi(X) - t - n_{II} - n_{III} + 1$.

Proof. The statement follows from Theorem 3.3.1, Corollary 3.2.5, and the fact that

$$(3.5.2) \quad \chi(X) = |\mathcal{E}_{\text{Sk}}| + 6t + 2n_{II} + 3n_{III} + 4n_{IV},$$

where $\text{Sk} = \text{Sk}_X$. (Here, we admit skeletons with bivalent \bullet -vertices as well.) For the latter, observe that $\chi(X)$ equals the total multiplicity of the singular fibers of X . Exceptional singular fibers are accounted for by the mono- and bivalent \bullet -vertices and monovalent \circ -vertices of Sk . Besides, there is one fiber of type I_p or I_p^* inside each p -gonal region of Sk . The sum of all indices p is the total number of corners of all regions of Sk , *i.e.*, $|\mathcal{E}_{\text{Sk}}|$. Finally, each $*$ -type fiber increases the total multiplicity by 6. \square

3.5.3. Theorem. *Let X be a non-isotrivial elliptic surface without type II^* or IV singular fibers. Then the index of the reduced monodromy group $\text{Im } \mathfrak{h}_X \subset \Gamma$ of X divides $\chi(X) - 6t - 2n_{II} - 3n_{III}$. In particular, it is finite.*

Proof. Let Sk be the skeleton of X . After a fiberwise equisingular deformation of X , not necessarily small, one can assume that Sk is generic and connected. (For the modifications of skeletons resulting in deformations of surfaces, see [8] or [13].)

Hence Sk is a $(3, 1)$ -skeleton. This time, each region of Sk may contain several singular fibers of X . Hence, instead of Theorem 3.3.1, one has a diagram

$$\pi_1(B^\sharp, e) \longleftarrow \pi_1(\text{Sk}', e) \longrightarrow \pi_1^{\text{orb}}(\text{Sk}_X, e) \xrightarrow{\cong} \text{Stab } e \subset \Gamma$$

(where Sk' is the auxiliary space introduced in Remark 3.3.2) and an inclusion $\text{Stab } e \subset \text{Im } \mathfrak{h}_X$. It remains to observe that $[\Gamma : \text{Stab } e] = |\mathcal{E}_{\text{Sk}}|$ and that (3.5.2) holds for any non-isotrivial surface X . \square

3.5.4. Remark. The reduced monodromy group $\text{Im } \mathfrak{h}_X$ of an isotrivial elliptic surface X is either trivial or conjugate to the subgroup generated by \mathbb{X} or \mathbb{Y} . In particular, $[\Gamma : \text{Im } \mathfrak{h}_X] = \infty$. At present, I do not know whether the index of $\text{Im } \mathfrak{h}_X$ is necessarily finite if X is a non-isotrivial surface with type II^* or IV singular fibers.

3.6. Further generalizations. Combined, the constructions of Subsections 3.1 and 3.2 give rise to the notion of *generalized* (i.e., possibly infinite) $(3, 1)$ -skeleton. Theorems 3.1.1 and 3.2.1 would combine to the following statement.

3.6.1. Theorem. *The functors $(\text{Sk}, e) \mapsto \text{Stab } e$, $H \mapsto (\Gamma/H, H/H)$ establish an equivalence of the categories of*

- pointed generalized $(3, 1)$ -skeletons and morphisms and
- subgroups $H \subset \Gamma$ and inclusions. \square

The orbifold fundamental group $\pi_1^{\text{orb}}(\text{Sk}, e)$ of a generalized $(3, 1)$ -skeleton Sk is defined as in 3.2.2, and Theorem 3.2.4 extends to this case literally. Since Sk^\bullet is still homotopy equivalent to a wedge of circles and copies of D_2^2 and D_3^2 , one obtains the following corollary.

3.6.2. Corollary. *Any subgroup of Γ is a free product (possibly infinite) of copies of cyclic groups \mathbb{Z} , \mathbb{Z}_2 , and \mathbb{Z}_3 .* \square

3.6.3. Under Theorem 3.6.1, finitely generated subgroups correspond to *almost contractible* $(3, 1)$ -skeletons, which are defined as those with finitely generated group $\pi_1^{\text{orb}}(\text{Sk})$. Following the proof of Proposition 3.1.3, one can easily show that any almost contractible $(3, 1)$ -skeleton Sk representing a finitely generated subgroup $H \subset \Gamma$, $H \neq \{1\}$ (so that Sk is not the Farey tree), admits a strict deformation retraction to a canonically defined finite induced subgraph $\text{Sk}^c \subset \text{Sk}$, called the *compact part* of Sk , with the following properties:

- (1) all vertices of Sk^c are of valency 3 or 1;
- (2) the monovalent vertices of Sk^c are divided into three types: \circ , \bullet , or \triangle (the latter representing maximal infinite branches of Sk);
- (3) distinct \triangle -vertices are adjacent to distinct trivalent vertices.

Under this correspondence $H \leftrightarrow \text{Sk} \leftrightarrow \text{Sk}^c$ one has (anti-)isomorphisms $N(H)/H = \text{Aut } \text{Sk} = \text{Aut } \text{Sk}^c$ and $H = \pi_1^{\text{orb}}(\text{Sk}) = \pi_1^{\text{orb}}(\text{Sk}^c)$, where $\pi_1^{\text{orb}}(\text{Sk}^c)$ is defined similar to $\pi_1^{\text{orb}}(\text{Sk})$, as the fundamental group of the space $(\text{Sk}^c)^\bullet$ obtained from Sk^c by replacing each monovalent \circ - or \bullet -vertex with a copy of D_2^2 or D_3^2 , respectively.

4. PSEUDO-TREES

Here, we introduce and count admissible trees and related ribbon graphs, called pseudo-trees; they are the principal source of most exponentially large examples stated in the introduction.

4.1. Admissible trees and pseudo-trees. An embedded tree $\Xi \subset S^2$ is called *admissible* if all its vertices have valency 3 (*nodes*) or 1 (*leaves*). Two such trees are called *isomorphic* if they are related by an orientation preserving auto-homeomorphism of S^2 . Each admissible tree Ξ gives rise to its *associated 3-skeleton* Sk_Ξ : one attaches a small loop to each leaf of Ξ , see Figure 4, left. A 3-skeleton obtained in this way is called a *pseudo-tree*. Clearly, each pseudo-tree is a skeleton of genus 0; two pseudo-trees $\text{Sk}_{\Xi'}$ and $\text{Sk}_{\Xi''}$ are isomorphic as ribbon graphs if and only if the trees Ξ' and Ξ'' are isomorphic.




FIGURE 4. An admissible tree Ξ (black) and associated 3-skeleton Sk_Ξ (left); the related binary tree (right)

An admissible tree has a certain number $k \geq 0$ of nodes and $(k + 2)$ leaves. The number of isomorphism classes of admissible trees with k nodes is denoted by $T(k)$; it equals to the number of isomorphism classes of pseudo-trees with $(2k + 2)$ vertices.

4.1.1. A *marking* of an admissible tree Ξ is a choice of one of its leaves v_1 . Given a marking, one can number all leaves of Ξ consecutively, starting from v_1 and moving in the clockwise direction (see Figure 4, where the indices of the leaves are shown inside the loops). Declaring the node adjacent to v_1 the root and removing all leaves, one obtains an oriented rooted binary tree with k vertices, see Figure 4, right; conversely, an oriented rooted binary tree B gives rise to a unique marked admissible tree: one attaches a leaf v_1 at the root of B and an extra leaf instead of each missing branch of B . As a consequence, the number of isomorphism classes of marked admissible trees with k nodes is given by the Catalan number $C(k)$.

4.1.2. The *vertex distance* m_i between two consecutive leaves v_i, v_{i+1} of a marked admissible tree Ξ is the vertex length of the shortest left turn path in Ξ from v_i to v_{i+1} . For example, in Figure 4 one has $(m_1, m_2, m_3, m_4, m_5) = (5, 3, 4, 5, 3)$; for another example, see Figure 7 in Subsection 5.3. The vertex distance between two leaves $v_i, v_j, j > i$, is defined to be $\sum_{s=i}^{j-1} m_s$; it is the vertex length of the shortest left turn path connecting v_i to v_j in the associated 3-skeleton Sk_Ξ .

One can extend the sequence (m_1, \dots, m_{k+1}) by appending the vertex distance m_{k+2} from v_{k+2} to v_1 ; then one has $m_1 + \dots + m_{k+2} = 5k + 4$. Two marked trees are isomorphic if and only if their sequences (m_1, \dots, m_{k+1}) are equal. Two unmarked trees are isomorphic if and only if the corresponding extended sequences $(m_1, \dots, m_{k+1}, m_{k+2})$ differ by a cyclic permutation. Note that *not* any sequence (m_1, \dots, m_{k+1}) gives rise to a marked admissible tree, see [12] for a criterion.

4.2. Counts. As above, let $T(k)$ be the number of isomorphism classes of pseudo-trees with $(2k + 2)$ vertices. Let, further, $T_i(k), i \geq 0$, be the number of classes of pseudo-trees Sk with $|\text{Aut Sk}| = i$.

For a pseudo-tree Sk with $(2k + 2)$ vertices, denote by \mathfrak{D}_{Sk} the orbit of $\mathbb{X}\mathbb{Y}$ corresponding to the outer $(5k + 4)$ -gonal region of Sk . The number of isomorphism classes of pointed 3-skeletons (Sk, e) , where Sk is a pseudo-tree with $(2k + 2)$ vertices and $e \in \mathfrak{D}_{\text{Sk}}$, is denoted by $\tilde{T}(k)$.

4.2.1. Lemma. *For a pseudo-tree $\text{Sk} = \text{Sk}_{\Xi}$ one has $|\text{Aut Sk}| \leq 3$, i.e., $T_i(k) = 0$ for $i > 3$. The numbers $T_1(k), T_2(k), T_3(k)$ are subject to the relations*

$$\sum_{i=1}^3 \frac{T_i(k)}{i} = \frac{C(k)}{k+2},$$

$$T_2(k) = \begin{cases} C(k'), & \text{if } k = 2k', \\ 0, & \text{otherwise,} \end{cases} \quad T_3(k) = \begin{cases} C(k'), & \text{if } k = 3k' + 1, \\ 0, & \text{otherwise.} \end{cases}$$

Furthermore, the group $\text{Aut Sk} = \text{Aut } \Xi$ acts freely on the set of leaves of the original tree Ξ and on the set \mathcal{E}_{Sk} of edge ends of Sk .

Proof. Obviously, one has $\text{Aut Sk}_{\Xi} = \text{Aut } \Xi$. Any combinatorial automorphism of Ξ is represented by a piecewise linear auto-homeomorphism $\varphi: \Xi \rightarrow \Xi$. Since Ξ is contractible, φ has a fixed point p , which is necessarily isolated (assuming that $\varphi \neq \text{id}$, as an automorphism of a connected ribbon graph fixing an edge is the identity). If p is at the center of an edge of Ξ (respectively, p is a vertex of Ξ), then φ^2 (respectively, φ^3) fixes a whole edge of Ξ and thus is the identity.




FIGURE 5. An automorphism of an admissible tree

A tree Ξ with an automorphism φ is shown in Figure 5. It is clear that such a tree admits no automorphisms other than powers of φ : the fixed point q of such an automorphism would belong to one of the grey areas and the vertices of Ξ would be distributed unevenly about q . Let k' be the number of nodes of the subtree Ξ' shown in the figure. In Figure 5, left ($|\text{Aut } \Xi| = 2$), one has $k = 2k'$; in Figure 5, right ($|\text{Aut } \Xi| = 3$), one has $k = 3k' + 1$. In each case, the trees Ξ admitting such an automorphism φ can be parameterized by the marked subtrees Ξ' , distinguished being the leaf extending towards the fixed point of φ . Their number is $C(k')$, which proves the expressions for $T_2(k)$ and $T_3(k)$.

It is also clear from Figure 5 that a non-trivial automorphism does not fix a leaf of Ξ or an edge end of Sk . Then the first relation in the statement is the usual orbit count: a tree Ξ with $|\text{Aut } \Xi| = i$ admits $(k + 2)/i$ essentially distinct markings, and the total number of marked trees is $C(k)$. \square

4.2.2. Corollary. *For each integer $k \geq 0$, one has*

$$T(k) = \frac{C(k)}{k+2} + \frac{T_2(k)}{2} + \frac{2T_3(k)}{3}, \quad \tilde{T}(k) = \frac{5k+4}{k+2}C(k),$$

where $T_2(k)$ and $T_3(k)$ are given by Lemma 4.2.1.

Proof. Since $T_i(k) = 0$ for $i > 3$, the expression for $T(k) = T_1(k) + T_2(k) + T_3(k)$ follows directly from Lemma 4.2.1.

For each pseudo-tree Sk , one has $|\mathfrak{D}_{\text{Sk}}| = 5k + 4$ and Aut Sk acts freely on \mathfrak{D}_{Sk} . Hence $\tilde{T}(k) = (5k + 4) \sum_{i=1}^3 T_i(k)/i = (5k + 4)C(k)/(k + 2)$ due to the first relation in Lemma 4.2.1. \square

4.3. Proof of Theorem 1.3.2. The surfaces in question were constructed in [8]. Each surface X corresponds to a pseudo-tree Sk with $(2k + 2)$ vertices, with the type specification (see Subsection 3.4 and Remark 3.4.3) chosen so that the singular fiber of X inside each monogonal region of Sk should be of type I_1 . The type of the singular fiber inside the remaining $(5k + 4)$ -gonal region (the outer region in Figure 4, left) is then determined by the parity of k : it is of type I_{5k+4} if k is odd or I_{5k+4}^* if k is even.

The $T(k)$ distinct pseudo-trees with $(2k + 2)$ vertices give rise to $T(k)$ pairwise non-isomorphic extremal elliptic surfaces; Theorem 1.3.1 implies that they are not related by a 2-orientation preserving fiberwise homeomorphism. \square

4.4. Generalized pseudo-trees. The construction of Subsection 4.1 producing a 3-skeleton from a tree can be generalized. A function ℓ defined on the set of leaves of an admissible tree Ξ and taking values in $\{0, \circ, \bullet, \Delta\}$ is called *admissible* if no two leaves v_1, v_2 with $\ell(v_1) = \ell(v_2) = \Delta$ are adjacent to the same node. An *admissible pair* is a pair (Ξ, ℓ) , where Ξ is an admissible tree and ℓ is an admissible function on the set of leaves of Ξ . Each admissible pair (Ξ, ℓ) gives rise to an (almost contractible) $(3, 1)$ -skeleton $\text{Sk}_{(\Xi, \ell)}$, whose compact part Sk^c is obtained from Ξ by attaching a small loop to each leaf v with $\ell(v) = 0$ and replacing each other leaf v with a monovalent vertex of type $\ell(v)$, cf. Figures 8 and 9 in Section 5. Thus, one has $\text{Sk}_{\Xi} = \text{Sk}_{(\Xi, 0)}$. A generalized $(3, 1)$ -skeleton obtained in this way is called a *generalized pseudo-tree*.

Clearly, two generalized pseudo-trees $\text{Sk}_{(\Xi', \ell')}$ and $\text{Sk}_{(\Xi'', \ell'')}$ are isomorphic if and only if so are pairs (Ξ', ℓ') and (Ξ'', ℓ'') , i.e., if there exists an isomorphism $\varphi: \Xi' \rightarrow \Xi''$ such that $\ell' = \ell'' \circ \varphi$.

For a generalized pseudo-tree $\text{Sk} = \text{Sk}_{(\Xi, \ell)}$, we denote by $n_*(\text{Sk})$, $* \in \{\circ, \bullet, \Delta\}$, the number of *monovalent* $*$ -vertices of the compact part Sk^c . Thus, $n_*(\text{Sk}) = |\ell^{-1}(*)|$.

4.4.1. Proposition. *Let $H \subset \Gamma$ be a proper finitely generated subgroup. Then H is generated by $H \cap \llbracket \mathbb{X}\mathbb{Y} \rrbracket_{\Gamma}$ if and only if Γ/H is a generalized pseudo-tree without monovalent vertices (i.e., a skeleton $\text{Sk}_{(\Xi, \ell)}$ with ℓ taking values in $\{0, \Delta\}$). If this is the case, H admits a free basis consisting of elements conjugate to $\mathbb{X}\mathbb{Y}$.*

Proof. Let $\text{Sk} = \Gamma/H$. It is an almost contractible $(3, 1)$ -skeleton, see 3.6.3. Since H is proper, Sk has a well defined compact part Sk^c , which is *not* isomorphic to the skeleton $\bullet \text{---} \circ$ representing Γ itself. Hence, each monogonal region of Sk (orbit of $\mathbb{X}\mathbb{Y}$ of length one) is bounded by an edge with both ends attached to a trivalent \bullet -vertex. (The only exceptional monogonal region is the ‘outer’ region in the skeleton $\bullet \text{---} \circ$ representing Γ .) It follows that the edge bounding a monogonal region cannot belong to any subtree of Sk^c .

Let Ξ be a maximal tree in Sk^c not containing a monovalent \circ - or \bullet -vertex. Contracting Ξ establishes a homotopy equivalence of the space $(\text{Sk}^c)^{\bullet}$ computing $\pi_1^{\text{orb}}(\text{Sk}^c) = H$, see 3.6.3, to a wedge W of circles and copies of D_2^2 and D_3^2 . Each

monogonal region of Sk produces a separate circle in W , and the H -conjugacy classes of loops represented by these circles constitute the intersection $H \cap \llbracket \mathbb{X}\mathbb{Y} \rrbracket$. Thus, H is generated by $H \cap \llbracket \mathbb{X}\mathbb{Y} \rrbracket$ if and only if W has no other circles or copies of D_2^2 or D_3^2 , *i.e.*, Sk^c consists of several monogonal regions attached to the (unique) maximal subtree $\Xi \subset \text{Sk}^c$. \square

4.4.2. Remark. Proposition 4.4.1 gives a geometric characterization of the proper subgroups $H \subset \Gamma$ that can appear as the monodromy group of a simple Γ -valued BM-factorization, see Definition 5.1.3. Note that Γ itself can also appear in this way (it is generated by the images $\mathbb{X}\mathbb{Y}$ and $\mathbb{X}^2\mathbb{Y}\mathbb{X}^{-1}$ of σ_1 and σ_2 , respectively, see (5.1.1) below); it is the only monodromy group that is not free.

4.5. More examples of elliptic surfaces. Let $\text{Sk} = \text{Sk}_{(\Xi, \ell)}$ be a finite generalized pseudo-tree (thus, we assume that $n_\Delta(\text{Sk}) = 0$) obtained from an admissible tree Ξ with k nodes. Let $n_* = n_*(\text{Sk})$. For the type specification (see Subsection 3.4 and Remark 3.4.3), assign type I_1 to each monogonal region of Sk and types IV^* and III^* to the monovalent \bullet - and \circ -vertices, respectively. Then the fiber inside the remaining outer region of Sk is of type I_s if $k + n_\bullet + n_\circ$ is odd or I_s^* otherwise, where $s = 5k + 4 - n_\bullet - 2n_\circ$. (For even more examples, one could also vary the types I_1 or I_1^* of the fibers in the monogonal regions, adjusting the type of the remaining fiber accordingly.)

The skeleton Sk and the type specification described above define an extremal elliptic surface X with the combinatorial type of singular fibers

$$(k + 2 - n_\bullet - n_\circ)\text{I}_1 \oplus n_\bullet\text{IV}^* \oplus n_\circ\text{III}^* \oplus \{\text{I}_s \text{ or } \text{I}_s^*\}.$$

The surfaces corresponding to non-isomorphic pairs (Ξ, ℓ) are neither analytically isomorphic nor related by a 2-orientation preserving fiberwise homeomorphism, as they have non-conjugate reduced monodromy groups.

5. BM-FACTORIZATIONS

This section deals with BM-factorizations. We prove Theorems 1.2.1 and 1.2.2 and discuss a few sporadic examples arising from generalized pseudo-trees and from maximizing plane sextics.

5.1. Preliminaries. The *braid group* \mathbb{B}_3 is the group

$$\mathbb{B}_3 = \langle \sigma_1, \sigma_2 \mid \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \rangle = \langle u, v \mid u^3 = v^2 \rangle,$$

where $u = \sigma_2\sigma_1$ and $v = \sigma_2\sigma_1^2$. The center $Z(\mathbb{B}_3)$ is the infinite cyclic group generated by $u^3 = v^2$, and the quotient $\mathbb{B}_3/Z(\mathbb{B}_3)$ is isomorphic to Γ . In order to be consistent with Subsection 2.5, we define the epimorphism $\mathbb{B}_3 \twoheadrightarrow \tilde{\Gamma}$ (and further to Γ) *via*

$$(5.1.1) \quad \sigma_1 \mapsto \mathbb{X}\mathbb{Y}, \quad \sigma_2 \mapsto \mathbb{X}^2\mathbb{Y}\mathbb{X}^{-1}.$$

(Then $u \mapsto -\mathbb{X}^{-1}$ and $v \mapsto -\mathbb{Y}$.)

5.1.2. The abelianization $\mathbb{B}_3/[\mathbb{B}_3, \mathbb{B}_3]$ is the cyclic group \mathbb{Z} . The image of a braid $\beta \in \mathbb{B}_3$ in the abelianization $\mathbb{B}_3/[\mathbb{B}_3, \mathbb{B}_3] = \mathbb{Z}$ is called its *degree* $\deg \beta$. (By convention, $\deg \sigma_1 = 1$.) A braid $\beta \in \mathbb{B}_3$ is uniquely recovered from its image $\bar{\beta} \in \Gamma$ and its degree $\deg \beta$; the latter is determined by $\bar{\beta}$ up to a multiple of 6. (The degree of an element of Γ or $\tilde{\Gamma}$ is defined, respectively, modulo 6 or 12.)

5.1.3. Definition. A \mathbb{B}_3 - (respectively, Γ - or $\tilde{\Gamma}$ -) valued BM-factorization (\mathbf{m}_i) , $i = 1, \dots, r$, is called *simple* if each entry \mathbf{m}_i belongs to the conjugacy class $[[\sigma_1]]$ (respectively, $[[\mathbb{X}\mathbb{Y}]]_\Gamma$ or $[[\mathbb{X}\mathbb{Y}]]_{\tilde{\Gamma}}$).

5.1.4. Proposition. For each $r \geq 1$, the epimorphisms $\mathbb{B}_3 \twoheadrightarrow \tilde{\Gamma} \twoheadrightarrow \Gamma$ establish bijections between the sets of simple \mathbb{B}_3 -, $\tilde{\Gamma}$ -, and Γ -valued BM-factorizations of length r ; these bijections preserve the weak/strong equivalence classes.

Proof. Each element $x \in [[\mathbb{X}\mathbb{Y}]] \subset \Gamma$ lifts to a unique element $x' \in [[\sigma_1]] \subset \mathbb{B}_3$ and to a unique element $x'' \in [[\mathbb{X}\mathbb{Y}]] \subset \tilde{\Gamma}$ (characterized by the requirement that $\deg x' = 1$ and $\deg x'' = 1 \pmod{12}$), establishing a one-to-one correspondence between the sets of BM-factorizations. The weak and strong Hurwitz equivalences are preserved due to the fact that both $\mathbb{B}_3 \twoheadrightarrow \Gamma$ and $\tilde{\Gamma} \twoheadrightarrow \Gamma$ are central extensions. \square

5.1.5. The advantage of considering the braid group \mathbb{B}_3 rather than the modular group Γ is the fact that, in \mathbb{B}_3 , the length r of a BM-factorization of an element $\mathbf{m}_\infty \in \mathbb{B}_3$ is uniquely determined by \mathbf{m}_∞ : one has $r = \deg \mathbf{m}_\infty$. Hence, for \mathbb{B}_3 , the problem of uniqueness of a BM-factorization of a given element can be restated in the language of *factorization semigroup*, see [18] and [25].

5.1.6. Definition. The *factorization semigroup* is the semigroup \mathfrak{B}_n (with the group operation denoted by \cdot) generated by the elements $\beta \in [[\sigma_1]]_{\mathbb{B}_n}$ subject to the *Hurwitz relations* $\beta_1 \cdot \beta_2 = \beta_1^{-1} \beta_2 \beta_1 \cdot \beta_1 = \beta_2 \cdot \beta_2 \beta_1 \beta_2^{-1}$. The *evaluation anti-homomorphism* $v: \mathfrak{B}_n \rightarrow \mathbb{B}_n$ is defined via $v: \beta_1 \cdot \beta_2 \cdot \dots \cdot \beta_r \mapsto \beta_r \dots \beta_2 \beta_1$.

5.1.7. It is clear that an element $\bar{\mathbf{m}} \in \mathfrak{B}_n$ represents a strong Hurwitz equivalence class of simple \mathbb{B}_n -valued BM-factorizations (of length $\deg v(\bar{\mathbf{m}})$) and the value $v(\bar{\mathbf{m}})$ is merely the monodromy at infinity $\mathbf{m}_\infty(\bar{\mathbf{m}})$. Our Theorem 1.2.1 states that, for $n = 3$, the evaluation map v is not injective; moreover, the size of the pull-back $v^{-1}(\beta)$, $\beta \in \mathbb{B}_3$, may grow exponentially in the degree $\deg \beta$. Using the canonical inclusion $\mathbb{B}_3 \hookrightarrow \mathbb{B}_n$, one can easily conclude that the same assertion holds for any integer $n \geq 3$: the size of the pull-back $v^{-1}(\beta)$, $\beta \in \mathbb{B}_n$, may grow exponentially in the degree $\deg \beta$.

According to [25], the fact that v is not injective implies that \mathfrak{B}_n does *not* have the cancellation property, *i.e.*, an equality $\alpha_1 \cdot \beta = \alpha_2 \cdot \beta$ or $\beta \cdot \alpha_1 = \beta \cdot \alpha_2$ in \mathfrak{B}_n does not necessarily imply that $\alpha_1 = \alpha_2$.

5.2. Proof of Theorems 1.2.1 and 1.2.2. Consider a marked admissible tree (Ξ, v_1) with k nodes and $(k+2)$ leaves and let $\text{Sk} = \text{Sk}_\Xi$ be the associated pseudo-tree, see Subsection 4.1. Let (m_1, \dots, m_{k+1}) be the sequence of consecutive vertex distances, see 4.1.2, and consider the distances


$$(5.2.1) \quad n_i = m_i + \dots + m_{k+1}, \quad i = 1, \dots, k+1, \quad n_{k+2} = 0$$

from v_i to v_{k+2} in Sk .

Let $e \in \mathcal{E}_{\text{Sk}}$ be the edge end at v_{k+2} that belongs to the original tree, see the grey dot in Figure 6, and consider the basis $\{\gamma_1, \dots, \gamma_{k+2}\}$ for $\pi_1(\text{Sk}, e)$, where γ_i is the class represented by the loop of Sk attached at v_i which is connected to e by the shortest left turn path in Sk (the grey loop in Figure 6).

In terms of Definition 2.4.1, the loop representing a basis element γ_i is (e, w_i) , where

$$w_i = (\text{nx op})^{n_i} (\text{nx op nx}^{-1} \text{nx}^{-1}) (\text{op nx}^{-1})^{n_i}.$$

FIGURE 6. A loop γ_i (grey)

The product $\gamma_1 \dots \gamma_{k+1}$ is homotopic to the boundary of the outer $(5k+4)$ -gonal region of Sk ; after cancellation, $\gamma_1 \dots \gamma_{k+2} \sim (e, (\text{nx op})^{5k+4})$.

Define the Γ -valued BM-factorization $\bar{\mathbf{m}} = \bar{\mathbf{m}}(\text{Sk}, e) = (\mathbf{m}_1, \dots, \mathbf{m}_{k+2})$ via

$$(5.2.2) \quad \mathbf{m}_i = (\text{val } \gamma_i)^{-1} = (\mathbb{X}\mathbb{Y})^{n_i} (\mathbb{X}^2 \mathbb{Y} \mathbb{X}^{-1}) (\mathbb{X}\mathbb{Y})^{-n_i}.$$

By construction, one has $\mathbf{m}_\infty(\bar{\mathbf{m}}) = (\mathbb{X}\mathbb{Y})^{-5k-4}$, see Lemma 2.4.4, and $\text{Im}(\bar{\mathbf{m}}) = \pi_1(\text{Sk}, e) = \text{Stab } e$, see Theorem 2.5.3. Regarding each \mathbf{m}_i in (5.2.2) as an element of $\tilde{\Gamma}$ and adjusting degree modulo 12, one obtains $\mathbf{m}_\infty(\bar{\mathbf{m}}) = -(-\mathbb{X}\mathbb{Y})^{-5k-4} \in \tilde{\Gamma}$.

5.2.3. Remark. Note that the particular choice of a basis $\{\gamma_i\}$ used above is not very important: by Artin's theorem [3], any other basis $\{\gamma'_i\}$ with the property that each γ'_i is conjugate to some γ_j and $\gamma'_1 \dots \gamma'_{k+2} = \gamma_1 \dots \gamma_{k+2}$ is obtained from $\{\gamma_i\}$ by a sequence of Hurwitz moves; hence the resulting BM-factorization $\bar{\mathbf{m}}'$ would be strongly equivalent to $\bar{\mathbf{m}}$.

Now, observe that e belongs to the orbit \mathfrak{D}_{Sk} introduced in 4.2. Let $e' \in \mathfrak{D}_{\text{Sk}}$ be another element of this orbit, $e' = (\mathbb{X}\mathbb{Y})^s e$, and consider the BM-factorization $\bar{\mathbf{m}}' = \bar{\mathbf{m}}(\text{Sk}, e') := (\mathbb{X}\mathbb{Y})^s \bar{\mathbf{m}}(\text{Sk}, e) (\mathbb{X}\mathbb{Y})^{-s}$. Clearly, one has $\mathbf{m}_\infty(\bar{\mathbf{m}}') = (\mathbb{X}\mathbb{Y})^{-5k-4}$ and $\text{Im}(\bar{\mathbf{m}}') = \pi_1(\text{Sk}, e') = \text{Stab } e'$. As above, the strong equivalence class of $\bar{\mathbf{m}}(\text{Sk}, e')$ does not depend on the particular choice of a basis for $\pi_1(\text{Sk}, e')$; for this reason, we omit the reference to the marking of the original tree Ξ in the notation.

Considering all $\tilde{T}(k)$ pairwise non-isomorphic pairs (Sk, e) , $e \in \mathfrak{D}_{\text{Sk}}$, see Subsection 4.2 and Corollary 4.2.2, one obtains $\tilde{T}(k)$ distinct BM-factorizations $\bar{\mathbf{m}}(\text{Sk}, e)$; they differ by the monodromy groups $\text{Im}(\bar{\mathbf{m}}(\text{Sk}, e)) = \text{Stab } e$, see Theorem 2.3.4. Disregarding the base points e , one arrives at $T(k)$ weak equivalence classes, which differ by the conjugacy class $[[\text{Im}(\bar{\mathbf{m}}(\text{Sk}, e))] = [[\text{Stab } \text{Sk}]]$, see Corollary 2.3.5.

The transcendental lattices and fundamental groups of the BM-factorizations constructed above are computed in [12]; for the former, see Example 7.2.3. \square

5.2.4. Remark. The BM-factorizations (5.2.2) represent the reduced homological invariants of the extremal elliptic surfaces constructed in Subsection 4.3.

5.3. Examples. Thus, the $T(k)$ weak equivalence classes of BM-factorizations given by Theorem 1.2.2 are numbered by the isomorphism classes of admissible trees with k nodes. They are given by (5.2.2), where the sequence (n_1, \dots, n_{k+2}) is obtained from the vertex distances (m_1, \dots, m_{k+1}) of the tree, see 4.1.2. The lifts to simple \mathbb{B}_3 -valued BM-factorizations are

$$(5.3.1) \quad \mathbf{m}_i = \sigma_1^{n_i} \sigma_2 \sigma_1^{-n_i}, \quad i = 1, \dots, k+2, \quad \mathbf{m}_\infty = (\sigma_1 \sigma_2)^{3(k+1)} \sigma_1^{-5k-4}.$$

(For \mathbf{m}_∞ , we multiply σ_1^{-5k-4} by a power of the central element $(\sigma_1 \sigma_2)^3$ in order to match the degree.)

5.3.2. Example. The simplest example of non-equivalent BM-factorizations given by Theorem 1.2.2 is obtained when $k = 4$. The four admissible trees with four nodes and their vertex distances are shown in Figure 7. The fact that the resulting BM-factorizations are not equivalent can be proved directly, using GAP [15]. Let \bar{m} be one of the BM-factorizations, let $H = \text{Im}(\bar{m})$ be its monodromy group, and let N be the normalizer of H in Γ . Then, as Corollary 2.3.6 predicts, the index $[N : H]$ equals 1, 2, and 3 for the trees in Figure 7, left, middle, and right, respectively. In particular, the four groups belong to at least three distinct conjugacy classes. The two groups corresponding to the two trees in the middle (which are related by an orientation reversing diffeomorphism of the sphere) are conjugate in $PGL(2, \mathbb{Z})$ but not in Γ .




FIGURE 7. Admissible trees with four nodes

5.3.3. Example. The simplest example of weakly but not strongly equivalent BM-factorizations with the same monodromy at infinity is given by Theorem 1.2.1 with $k = 0$. The only admissible tree without nodes (two leaves connected by an edge) gives rise to two BM-factorizations:

$$\bar{m}' = (\sigma_1^2 \sigma_2 \sigma_1^{-2}, \sigma_2), \quad \bar{m}'' = (\sigma_1 \sigma_2 \sigma_1^{-1}, \sigma_1^{-1} \sigma_2 \sigma_1).$$

Let $H', H'' \subset \Gamma$ be their monodromy groups (reduced to Γ). Using GAP [15], one can see that $[\Gamma : H'] = [\Gamma : H''] = 6$ whereas $[\Gamma : H' \cap H''] = 24$. Hence $H' \neq H''$.

5.4. Non-equivalent BM-factorizations of length two. Consider the almost contractible generalized pseudo-trees represented by the two ribbon graphs shown in Figure 8. (Recall that each Δ -vertex is to be extended to a maximal infinite branch, which is a ‘half’ of the Farey three, see Subsection 3.1.) They are obviously not isomorphic; hence their stabilizers are not conjugate.



FIGURE 8. Almost contractible pseudo-trees with two loops

In each skeleton S_k , let $e \in \mathcal{E}_{S_k}$ be the edge end represented by a grey dot in the figure, and pick a basis $\{\gamma_1, \gamma_2\}$ for $\pi_1(S_k, e)$ so that each γ_i , $i = 1, 2$, is conjugate to the boundary of a monogonal region of S_k and $\gamma_1 \gamma_2$ is homotopic to a circle

encompassing the compact part Sk^c of Sk . (The particular choice of bases is not important, see Remark 5.2.3.) Let $\bar{\mathbf{m}}(\text{Sk}) = ((\text{val } \gamma_1)^{-1}, (\text{val } \gamma_2)^{-1})$. For example, the bases can be chosen so that

$$\begin{aligned}\bar{\mathbf{m}}(\text{Sk}_{\text{left}}) &= ((\mathbb{X}\mathbb{Y})(\mathbb{X}^2\mathbb{Y}\mathbb{X}^{-1})(\mathbb{X}\mathbb{Y})^{-1}, (\mathbb{Y}\mathbb{X}\mathbb{Y})(\mathbb{X}^2\mathbb{Y}\mathbb{X}^{-1})(\mathbb{Y}\mathbb{X}\mathbb{Y})^{-1}), \\ \bar{\mathbf{m}}(\text{Sk}_{\text{right}}) &= (\mathbb{X}^2\mathbb{Y}\mathbb{X}^{-1}, (\mathbb{Y}\mathbb{X}\mathbb{Y}\mathbb{X}^2\mathbb{Y})(\mathbb{X}^2\mathbb{Y}\mathbb{X}^{-1})(\mathbb{Y}\mathbb{X}\mathbb{Y}\mathbb{X}^2\mathbb{Y})^{-1}).\end{aligned}$$

The \mathbb{B}_3 -valued simple lifts of the two factorizations are

$$\bar{\mathbf{m}}(\text{Sk}_{\text{left}}) = (\sigma_1\sigma_2\sigma_1^{-1}, \sigma_2\sigma_1^3\sigma_2\sigma_1^{-3}\sigma_2^{-1}), \quad \bar{\mathbf{m}}(\text{Sk}_{\text{right}}) = (\sigma_2, \beta\sigma_2\beta^{-1}),$$

where $\beta = \sigma_2\sigma_1^2\sigma_2^{-1}\sigma_1$. One has

$$\mathbf{m}_\infty(\bar{\mathbf{m}}(\text{Sk}_{\text{left}})) = \mathbf{m}_\infty(\bar{\mathbf{m}}(\text{Sk}_{\text{right}})) = \mathbb{Y}\mathbb{X}(\mathbb{X}\mathbb{Y})^{-3}\mathbb{Y}\mathbb{X}(\mathbb{X}\mathbb{Y})^{-3}$$

(or, respectively, $\mathbf{m}_\infty = (\sigma_2\sigma_1^3\sigma_2\sigma_1^{-1})^2 \in \mathbb{B}_3$). On the other hand, the monodromy groups $[[\text{Im}(\bar{\mathbf{m}}(\text{Sk}))]] = [[\text{Stab } \text{Sk}]]$ are not conjugate in Γ (although they are conjugate in $PGL(2, \mathbb{Z})$); hence the two BM-factorizations are not weakly equivalent.

5.4.1. Remark. The two pseudo-trees differ by an orientation reversing auto-homeomorphism of the sphere. This fact implies that the corresponding Hurwitz curves and Lefschetz fibrations are anti-isomorphic. Hence, the two BM-factorizations have isomorphic fundamental groups and transcendental lattices, see 1.1.3.

5.5. Non-simple BM-factorizations. Let $\text{Sk} = \text{Sk}_{(\Xi, \ell)}$ be a generalized pseudo-tree obtained from an admissible tree Ξ with k nodes, see Subsection 4.4. Denote $n_* = n_*(\text{Sk})$ for $*$ in $\{\bullet, \circ, \Delta\}$.

Consider an embedding $\text{Sk}^c \subset S^2$, patch each monogonal region of Sk^c with a disk, and let B be a regular neighborhood of the result. Denote by B^\sharp the punctured disk obtained from B by removing a point inside each monogonal region of Sk and all monovalent \bullet - and \circ -vertices of Sk , see the shaded area in Figure 9. There is an epimorphism $\rho: \pi_1(B^\sharp) \twoheadrightarrow \pi_1^{\text{orb}}(\text{Sk})$, cf. Theorem 3.3.1.




FIGURE 9. A generalized pseudo-tree Sk^c and punctured disk B^\sharp

Fix a point $b \in \partial B$ and pick a geometric basis $\{\gamma_1, \dots, \gamma_s\}$ for $\pi_1(B, b)$ such that $\gamma_1 \dots \gamma_s = [\partial B]$. (The precise choice is not important as different bases would produce weakly equivalent BM-factorizations, cf. Remark 5.2.3.) Define the BM-factorization $\bar{\mathbf{m}}(\text{Sk}) = (\mathbf{m}_1, \dots, \mathbf{m}_s)$ of length $s = k + 2 - n_\Delta$ via $\mathbf{m}_i = (\text{val } \rho(\gamma_i))^{-1}$, $i = 1, \dots, s$. It has n_\bullet elements in $[[\mathbb{X}]]$, n_\circ elements in $[[\mathbb{Y}]]$, and $k + 2 - n_\bullet - n_\circ - n_\Delta$ elements in $[[\mathbb{X}\mathbb{Y}]]$. Thus, $\bar{\mathbf{m}}$ is simple if and only if $n_\bullet = n_\circ = 0$.

If $n_\Delta = 0$, the conjugacy class of the monodromy at infinity $\mathbf{m}_\infty(\bar{\mathbf{m}}(\text{Sk}))$ equals $[[\mathbb{X}\mathbb{Y}]^{-n}]$, where $n = 5k + 4 - n_\bullet - 2n_\circ$, and $\bar{\mathbf{m}}(\text{Sk})$ represents the reduced homological invariant of an extremal elliptic surface constructed in Subsection 4.5. In

general, the monodromy at infinity can be found as follows. Let $(m_1, \dots, m_{n_\Delta})$ be the sequence of vertex distances in Sk^c between consecutive Δ -vertices, each distance being the length of the shortest left turn path connecting two Δ -vertices, with only \bullet -vertices counted. (For example, for the graph shown in Figure 9, starting from the upper left corner, one has $(m_1, m_2, m_3) = (6, 9, 4)$; in Figure 8, for both graphs one has $(m_1, m_2) = (5, 5)$.) Then, the conjugacy class of the monodromy at infinity $\mathfrak{m}_\infty(\bar{\mathfrak{m}}(\text{Sk}))$ is represented by the *right to left* product

$$(5.5.1) \quad \prod_{i=1}^{n_\Delta} (\mathbb{X}\mathbb{Y})^{m_i-1} \mathbb{X} = \dots (\mathbb{X}\mathbb{Y})^{m_2-1} \mathbb{X} (\mathbb{X}\mathbb{Y})^{m_1-1} \mathbb{X}.$$

Note that $\sum_{i=1}^{n_\Delta} m_i = 5k + 4 - n_\bullet - 2n_\circ - 2n_\Delta$.

5.5.2. Lemma. *Given two generalized pseudo-trees Sk' , Sk'' , the monodromies at infinity $\mathfrak{m}_\infty(\bar{\mathfrak{m}}(\text{Sk}'))$ and $\mathfrak{m}_\infty(\bar{\mathfrak{m}}(\text{Sk}''))$ are conjugate in Γ if and only if the corresponding sequences (m'_i) and (m''_j) differ by a cyclic permutation.*

Proof. The ‘if’ part is obvious. For the converse, observe that the admissibility condition in Subsection 4.4 implies that each entry m'_i, m''_j is at least 2. Then the cyclic word w given by (5.5.1) admits no cancellations and the distances m_i can be recovered from the distances in w between consecutive occurrences of \mathbb{X}^2 . \square

5.6. Maximizing plane sextics. We conclude this section with a few examples arising from maximizing plane sextics.

Consider a plane sextic $C \subset \mathbb{P}^2$ with simple singularities only and with a distinguished type \mathbf{E} singular point P . Let L_∞ be the (only) tangent to C at P . Assume that L_∞ is not a component of C and let $C^a \subset \mathbb{C}^2 = \mathbb{P}^2 \setminus L_\infty$ be the affine part of C . It is a *horizontal curve* in the sense of [2] (or *Hurwitz curve* in the sense of [20]) of degree 3 with respect to the pencil $\mathcal{P} = \{L_t\}$, $t \in \mathbb{C}^1$, of lines through P ; in other words, the projection $C^a \rightarrow \mathbb{C}^1$ defined by \mathcal{P} is a proper map. Hence, using \mathcal{P} and an appropriately chosen section of the projection, one can define the braid monodromy $\mu_C: \pi_1(B^\sharp) \rightarrow \mathbb{B}_3$, where B^\sharp is the base \mathbb{C}^1 of the pencil with the singular fibers removed. Then, choosing a geometric basis for $\pi_1(B^\sharp)$, one can represent μ_C by a BM-factorization $\bar{\mathfrak{m}}_C$, which is well defined up to weak Hurwitz equivalence.

The minimal resolution of singularities X of the double plane ramified at a sextic C as above is a $K3$ -surface, and the pencil \mathcal{P} lifts to an elliptic pencil $X \rightarrow \mathbb{P}^1$ with a distinguished section. One can easily show (see, *e.g.*, [9]) that X is extremal if and only if C is *maximizing*, *i.e.*, if its total Milnor number takes its maximal possible value 19. When this is the case, the combinatorial type of singular fibers of X is determined by the combinatorial type of singularities of C as follows:

- the distinguished singular point P of type \mathbf{E}_6 , \mathbf{E}_7 , or \mathbf{E}_8 gives rise to a singular fiber of type \mathbf{I}_6 , \mathbf{I}_2^* , or \mathbf{III}^* , respectively,
- each other singular point gives rise to a singular fiber of the following type: $\mathbf{A}_p \mapsto \mathbf{I}_{p+1}$, $p \geq 1$, $\mathbf{D}_q \mapsto \mathbf{I}_{q-4}^*$, $q \geq 4$, $\mathbf{E}_6 \mapsto \mathbf{IV}^*$, $\mathbf{E}_7 \mapsto \mathbf{III}^*$, $\mathbf{E}_8 \mapsto \mathbf{II}^*$,
- a number of type \mathbf{I}_1 fibers are added to make the total multiplicity 24.

Furthermore, the $\tilde{\Gamma}$ -valued reduction of the braid monodromy μ_C is the homological invariant $\tilde{\mathfrak{h}}_X$.

In [2], the authors construct a pair of reducible maximizing sextics C_1, C_2 with the set of singularities $\mathbf{E}_6 \oplus \mathbf{A}_7 \oplus \mathbf{A}_3 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$ and, using the fact that both curves

and all their singular fibers can be chosen real, compute their BM-factorizations \bar{m}_1 , \bar{m}_2 . Then, reducing \bar{m}_1 and \bar{m}_2 to the finite group $SL(2, \mathbb{Z}_{32})$ and using GAP [15], they compute their Hurwitz orbits and show that they are disjoint, concluding that \bar{m}_1 and \bar{m}_2 are not weakly equivalent and thus distinguishing the curves. (Both orbits are of length 15360.) In [11], the same pair of sextics is constructed using trigonal curves or, equivalently, extremal elliptic $K3$ -surfaces; their skeletons are as shown in Figure 10, with the distinguished fiber L_∞ corresponding to the outer region. Since the skeletons are obviously not isomorphic, Theorem 2.5.3 implies that $[[\text{Im}(\bar{m}_1)]] \neq [[\text{Im}(\bar{m}_2)]]$.




FIGURE 10. The set of singularities $\mathbf{E}_6 \oplus \mathbf{A}_7 \oplus \mathbf{A}_3 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$

5.6.1. Remark. Strictly speaking, constructed in [11] is merely a pair of not deformation equivalent sextics with the set of singularities $\mathbf{E}_6 \oplus \mathbf{A}_7 \oplus \mathbf{A}_3 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$. However, it follows from [27] that this set of singularities is realized by exactly two equisingular deformation families. Hence, the pairs found in [2] and [11] coincide.

A number of other examples is found in [10] and [11]. Listed in Table 2 are all sets of singularities realized by a pair C_1, C_2 of *irreducible* maximizing plane sextics with a distinguished type \mathbf{E} singular point and with essentially different skeletons. (More precisely, we ignore pairs of anti-isomorphic curves.) For each such pair, Theorem 2.5.3 implies that the corresponding BM-factorizations \bar{m}_1, \bar{m}_2 are not weakly equivalent, as their monodromy groups are not conjugate. For the sets of singularities marked with a *, the corresponding BM-factorizations differ by their transcendental lattices, see Example 7.2.2 below.

TABLE 2. Irreducible maximizing sextics with a type \mathbf{E} singular point

* $\mathbf{E}_8 \oplus \mathbf{A}_{10} \oplus \mathbf{A}_1$	$\mathbf{E}_6 \oplus \mathbf{D}_5 \oplus \mathbf{A}_8$	$(\mathbf{E}_6 \oplus \mathbf{A}_8 \oplus \mathbf{A}_2) \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$
$\mathbf{E}_8 \oplus \mathbf{A}_8 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	* $\mathbf{E}_6 \oplus \mathbf{D}_5 \oplus \mathbf{A}_6 \oplus \mathbf{A}_2$	* $\mathbf{E}_6 \oplus \mathbf{A}_7 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2$
* $\mathbf{E}_8 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_1$	* $\mathbf{E}_6 \oplus \mathbf{A}_{10} \oplus \mathbf{A}_3$	* $\mathbf{E}_6 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$
$\mathbf{E}_8 \oplus \mathbf{A}_5 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2$	* $\mathbf{E}_6 \oplus \mathbf{A}_{10} \oplus \mathbf{A}_2 \oplus \mathbf{A}_1$	$\mathbf{E}_6 \oplus \mathbf{A}_5 \oplus 2\mathbf{A}_4$
$(2\mathbf{E}_6 \oplus \mathbf{A}_5) \oplus \mathbf{A}_2$	$\mathbf{E}_6 \oplus \mathbf{A}_9 \oplus \mathbf{A}_4$	$(\mathbf{E}_6 \oplus \mathbf{A}_5 \oplus 2\mathbf{A}_2) \oplus \mathbf{A}_4$
$2\mathbf{E}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_3$	$\mathbf{E}_6 \oplus \mathbf{A}_8 \oplus \mathbf{A}_4 \oplus \mathbf{A}_1$	

5.6.2. Remark. It is worth mentioning that there also are three pairs C_1, C_2 of irreducible maximizing sextics, those with the sets of singularities

$$\mathbf{E}_7 \oplus \mathbf{E}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2, \quad \mathbf{E}_7 \oplus \mathbf{A}_{10} \oplus \mathbf{A}_2, \quad \mathbf{E}_7 \oplus \mathbf{A}_6 \oplus \mathbf{A}_4 \oplus \mathbf{A}_2$$

(the distinguished point P being that of type \mathbf{E}_7), such that, within each pair, the curves are not deformation equivalent but are represented by isomorphic skeletons, hence have equivalent BM-factorizations. It follows that the affine parts C_1^a, C_2^a

are isotopic in the class of Hurwitz curves, see [18]. In fact, the curves constituting each pair are related by a quadratic birational transformation biholomorphic in the affine part $\mathbb{P}^2 \setminus L_\infty$.

6. REAL TRIGONAL CURVES

Here, we give a brief introduction to theory of real trigonal curves (see [13] for more details), prove Theorem 1.4.1, and consider a few generalizations.

6.1. Dessins. Recall that a *real structure* on a complex analytic variety X is an anti-holomorphic involution $\text{conj}: X \rightarrow X$. A map, subvariety, *etc.* is called *real* if it commutes with/is preserved by conj .

For each Hirzebruch surface $\Sigma_k \rightarrow B \cong \mathbb{P}^1$, $k \geq 1$, fix a (unique up to automorphism) real structure $\text{conj}: \Sigma_k \rightarrow \Sigma_k$ with nonempty real part. Recall that the ruling of Σ_k restricts to an S^1 -fibration $(\Sigma_k)_\mathbb{R} \rightarrow B_\mathbb{R} \cong \mathbb{P}_\mathbb{R}^1 \cong S^1$ of the real parts, which is orientable if and only if k is even. The real part $E_\mathbb{R}$ of the exceptional section $E \subset \Sigma_k$ is a section of this fibration.

In what follows, we fix an orientation of $B_\mathbb{R}$ and denote by B_+ the closure of the connected component of $B \setminus B_\mathbb{R}$ whose complex orientation agrees with the chosen orientation of the boundary $\partial B_+ = B_\mathbb{R}$.

6.1.1. Given a trigonal curve $C \subset \Sigma_k$, one can define the *j -invariant* $j_C: B \rightarrow \mathbb{P}^1$ by sending a nonsingular fiber \bar{F} to the j -invariant of the elliptic curve F covering \bar{F} and ramified at $\bar{F} \cap (C \cup E)$. (Here, the target is the standard Riemann sphere $\mathbb{C} \cup \{\infty\}$.) Following [24] (see also [13] for more details), define the *dessin* of C as the graph $j_C^{-1}(\mathbb{P}_\mathbb{R}^1) \subset B$ with the following extra decoration:

- the pull-backs of 0, 1, and ∞ are \bullet -, \circ -, and \times -vertices, respectively;
- the pull-backs of $[0, 1]$, $[1, \infty]$, and $[-\infty, 0]$ are bold, dotted, and solid edges, respectively.

(Thus, the skeleton introduced in 2.2.4 is obtained from the dessin by removing all \times -vertices and solid and dotted edges.) The dessin of a real curve is invariant under the complex conjugation in B ; for this reason, we only draw the part contained in the closed disk B_+ . Vertices and edges of the dessin that belong to the boundary ∂B_+ are called *real*.

6.1.2. Remark. Note that the j -invariant of a real curve may have real critical values other than 0, 1, or ∞ not removable by a small equivariant deformation. For this reason, a generic symmetric dessin may have non-removable *monochrome* vertices in the boundary ∂B_+ , *cf.* Figure 11.

According to [24] and [13], a dessin in the *topological* disk B_+ determines a real trigonal curve C , which is well defined up to equivariant fiberwise deformation. (The converse is not true: a deformation of C may result in a non-trivial modification of its dessin, see [13] for details. We do not use this fact here.)

6.1.3. From now on, we assume all curves nonsingular and generic, *i.e.*, we assume that all singular fibers are of Kodaira type I_1 .

The real part $C_\mathbb{R} = \text{Fix conj}|_C$ of a real trigonal curve $C \subset \Sigma_k$ consists of a *long component* L isotopic to $E_\mathbb{R}$ and a number of *ovals*, *i.e.*, components contractible in $(\Sigma_k)_\mathbb{R}$. The critical values of the restriction $p: C_\mathbb{R} \rightarrow B_\mathbb{R}$ of the ruling are the real \times -vertices of the dessin of C . Pairs of such vertices bound *maximal dotted segments*

in ∂B_+ , each segment containing a number of monochrome vertices and, possibly, a number of real \circ -vertices. The projection p is three-to-one over the interior of each dotted segment, and it is one-to-one outside the dotted segments. A maximal dotted segment containing an even number of \circ -vertices is the projection of an oval, cf. Figure 11(a) and (b); a segment containing an odd number of \circ -vertices is the projection of a *zigzag* in L , cf. Figure 11(c).

The real \circ -vertices of the dessin are the points where $C_{\mathbb{R}}$ crosses the zero section of Σ_k . It follows that, if k is even, two ovals of $C_{\mathbb{R}}$ belong to the same connected component of the complement $(\Sigma_k)_{\mathbb{R}} \setminus (L \cup E_{\mathbb{R}})$ if and only if they are separated by an even number of real \circ -vertices.

6.2. Proof of Theorem 1.4.1. To construct a curve C_i as in the statement, consider one of the $T(k)$ pseudo-trees Sk_i with k nodes, see Subsection 4.1, and extend it to a dessin as shown in Figure 11(a) and (b). More precisely, embed Sk_i to the sphere S^2 (which is *not* the base of the elliptic pencil being constructed), patch each loop of Sk_i with the disk bounded by this loop, and take for B_+ a regular neighborhood of the result in S^2 . Then, place a \circ -vertex at the center of each edge and a \times -vertex at the center of each disk bounded by a loop, connect all \bullet - and \circ -vertices by appropriate edges to the boundary ∂B_+ in the radial manner, and connect the resulting monochrome vertices in ∂B_+ through \times -vertices.




FIGURE 11. Extending a skeleton Sk (bold) to a dessin

Each loop of Sk_i gives rise to an oval in ∂B_+ , see Figure 11(a), and each edge of the original tree Ξ_i gives rise to two ovals, see Figure 11(b). Thus, we obtain the dessin of a real trigonal curve in Σ_{2k+2} with $(5k+4)$ ovals. All curves obtained are topologically distinct: they differ by the monodromy group $[[\text{Stab Sk}_i]]$ of the monodromy $\pi_1(B_+^{\sharp}) \rightarrow \Gamma$, where B_+^{\sharp} is the interior of B_+ with the inner \times -vertices removed. \square

6.3. A generalization: ribbon curves. The real trigonal curves constructed in Subsection 6.2 are *ribbon curves* in the sense of [13]. This construction can be generalized. Let $\text{Sk} = \text{Sk}_{(\Xi, \ell)}$ be the generalized pseudo-tree obtained from an admissible tree Ξ with k nodes and a function ℓ taking values in $\{0, \bullet\}$, see Subsection 4.4. Let $z = n_{\bullet}(\text{Sk})$. Extend Sk to a dessin as shown in Figure 11. The new element here is Figure 11(c): the edge adjacent to a monovalent \bullet -vertex v is extended towards ∂B_+ and v is replaced with a \circ -vertex (which is bivalent in the complete dessin in B), giving rise to a zigzag rather than an oval. The result is the dessin of a real trigonal curve $C \subset \Sigma := \Sigma_{2k+2-z}$ with $(5k+4-z)$ ovals and z zigzags.

6.3.1. To distinguish the curves topologically, consider the region B_+^{\sharp} obtained from the interior of B_+ by adding small regular neighborhoods of the zigzags and removing the zigzags themselves and all inner \times -vertices, see Figure 12. Since




FIGURE 12. The region B_+^\sharp

zigzags are clearly distinguishable topologically, the monodromy $\pi_1(B_+^\sharp) \rightarrow \Gamma$ is a topological invariant of the curve. On the other hand, at least topologically, a pair of \times -vertices constituting a zigzag can collapse to a single type II singular fiber; hence the Γ -valued monodromy about a whole zigzag equals that about a monovalent \bullet -vertex. Thus, the image of the monodromy $\pi_1(B_+^\sharp) \rightarrow \Gamma$ equals $[[\text{Stab Sk}]]$, and distinct skeletons produce non-isotopic curves.

6.3.2. Let (m_1, \dots, m_z) be the sequence of vertex distances between consecutive monovalent \bullet -vertices of Sk (cf. Subsection 5.5; the monovalent \bullet -vertices themselves are also included into the count, so that each $m_i \geq 3$). Then the topology of pair $(\Sigma_{\mathbb{R}}, C_{\mathbb{R}})$ is uniquely determined by the following two properties:

- $C_{\mathbb{R}}$ does not intersect the zero section except once inside each zigzag;
- the pair of zigzags of $C_{\mathbb{R}}$ corresponding to a pair of consecutive monovalent \bullet -vertices at a distance m is separated by $(m - 3)$ ovals.

Similar to Lemma 5.5.2, one can easily see that the curves C' , C'' obtained from two skeletons Sk' , Sk'' as above have fiberwise isotopic real parts if and only if the corresponding sequences (m'_i) , (m''_j) differ by a cyclic permutation.

6.3.3. If $z = n_{\bullet}(\text{Sk})$ is even, the double covering X of Σ ramified at C and E is a generic Jacobian real elliptic surface. The surfaces obtained from distinct skeletons Sk or distinct (not related by an automorphism of Sk) lifts of the real structure are neither deformation equivalent nor isomorphic in the class of directed real Lefschetz fibrations, as they differ by the homological invariants, cf. 6.3.1. The necklace diagram of X , see [26], can be recovered from the sequence (m_1, \dots, m_z) introduced in 6.3.2: reading from m_z down to m_1 , each pair m_{2i}, m_{2i-1} gives rise to a copy of \rightarrow , followed by $(m_{2i} - 3)$ copies of $-\circ-$, a copy of \leftarrow , and $(m_{2i-1} - 3)$ copies of $-\square-$. Two sequences produce isomorphic necklace diagrams if and only if they differ by an *even* cyclic permutation. (Thus, the lift of the real structure is encoded in the choice of a marked monovalent \bullet -vertex of Sk .)

6.3.4. Remark. In the terminology of [13], the curves constructed in this section are *ribbon curves* with all blocks of type I_1 or II_3 . Conversely, any such curve C over the rational base is obtained by the above construction, and the ribbon curve structure of C is encoded by the original skeleton Sk . It follows that both the fiberwise deformation type and the fiberwise isotopy type of C determine its ribbon curve structure. In [13], a similar assertion is stated for ribbon curves with all blocks of type I_2 or II_3 .

6.3.5. Remark. It is worth emphasizing that the analytic and topological classifications of the curves constructed above coincide. This fact substantiates the conjecture that real trigonal curves are *quasi-simple*, i.e., the fiberwise equisingular deformation type of such a curve $C \subset \Sigma_k$ is determined by the topological type of the quadruple $(\Sigma_k, C; \text{pr}, \text{conj})$, where $\text{pr}: \Sigma_k \rightarrow \mathbb{P}^1$ is the ruling.

7. THE TRANSCENDENTAL LATTICE

In this section, we give a formal definition of a new invariant of BM-factorizations, which we call the *transcendental lattice*, and discuss a few open questions.

7.1. The construction. Fix a commutative ring R , two R -modules \mathcal{L} , \mathcal{V} , and a skew-symmetric bilinear form $\bigwedge^2 \mathcal{L} \rightarrow \mathcal{V}$, $x \wedge y \mapsto x \cdot y$. (In case \mathcal{V} has a 2-torsion, we assume, in addition, that $x \cdot x = 0$ for all $x \in \mathcal{L}$.) Fix, further, a symplectic (with respect to the chosen form) representation $G \rightarrow \mathrm{Sp} \mathcal{L}$.

7.1.1. Definition. Given a G -valued BM-factorization $\bar{\mathfrak{m}} = (\mathfrak{m}_1, \dots, \mathfrak{m}_r)$, define the following objects:

- (1) the R -module $\mathcal{L} \otimes \bar{\mathfrak{m}} := \bigoplus_{i=1}^r \mathcal{L}$;
- (2) the R -linear map $\chi: \mathcal{L} \otimes \bar{\mathfrak{m}} \rightarrow \mathcal{L}$, $\bigoplus_i x_i \mapsto \sum_i (\mathfrak{m}_i - 1)x_i$;
- (3) the R -quadratic map $q: \mathcal{L} \otimes \bar{\mathfrak{m}} \rightarrow \mathcal{V}$,

$$\bigoplus x_i \mapsto - \sum_{i=1}^r x_i \cdot \mathfrak{m}_i x_i + \sum_{1 \leq i < j \leq r} (\mathfrak{m}_i - 1)x_i \cdot (\mathfrak{m}_j - 1)x_j.$$

(Here, q is R -quadratic in the sense that $q(rx) = r^2 q(x)$ for all $x \in \mathcal{L} \otimes \bar{\mathfrak{m}}$, $r \in R$ and $(x, y) \mapsto q(x+y) - q(x) - q(y) \in \mathcal{V}$ is a \mathcal{V} -valued bilinear form.)

Let $\mathcal{L}_{\bar{\mathfrak{m}}} = \mathrm{Ker} \chi$, and define $\mathcal{L}_{\bar{\mathfrak{m}}}^{\perp} = \{x \in \mathcal{L}_{\bar{\mathfrak{m}}} \mid q(y+x) = q(y) \text{ for all } y \in \mathcal{L}_{\bar{\mathfrak{m}}}\}$. Then, $\mathcal{L}_{\bar{\mathfrak{m}}}^{\perp} \subset \mathcal{L}_{\bar{\mathfrak{m}}}$ is an R -submodule and the quotient $\mathcal{T}(\bar{\mathfrak{m}}) := \mathcal{L}_{\bar{\mathfrak{m}}}/\mathcal{L}_{\bar{\mathfrak{m}}}^{\perp}$ inherits a quadratic map $q: \mathcal{T}(\bar{\mathfrak{m}}) \rightarrow \mathcal{V}$. It is called the *transcendental lattice* of $\bar{\mathfrak{m}}$ (defined by the representation $G \rightarrow \mathrm{Sp} \mathcal{L}$).

7.1.2. Lemma. *One has $q(x+y) - q(x) - q(y) = \chi(x) \cdot \chi(y) \pmod{2\mathcal{V}}$ for any pair $x, y \in \mathcal{L} \otimes \bar{\mathfrak{m}}$.*

Proof. The proof is a simple computation taking into account the fact that each \mathfrak{m}_i is an isometry, so that $\mathfrak{m}_i x_i \cdot \mathfrak{m}_i y_i + x_i \cdot y_i = 2(x_i \cdot y_i) = 0 \pmod{2\mathcal{V}}$. \square

7.1.3. Corollary. *If \mathcal{V} is free of 2-torsion, the quadratic form $q: \mathcal{L}_{\bar{\mathfrak{m}}} \rightarrow \mathcal{V}$ extends to a symmetric bilinear form $\mathcal{L}_{\bar{\mathfrak{m}}} \otimes \mathcal{L}_{\bar{\mathfrak{m}}} \rightarrow \mathcal{V}$. \square*

The symmetric bilinear extension of q is also denoted by q . Its kernel equals the submodule $\mathcal{L}_{\bar{\mathfrak{m}}}^{\perp}$ defined above, and q factors to a nondegenerate symmetric bilinear form $q: \mathcal{T}(\bar{\mathfrak{m}}) \otimes \mathcal{T}(\bar{\mathfrak{m}}) \rightarrow \mathcal{V}$. The pair $(\mathcal{T}(\bar{\mathfrak{m}}), q)$ is still called the *transcendental lattice* of $\bar{\mathfrak{m}}$.

7.1.4. Remark. Assume that $\mathcal{L} = H_1(F)$ for a punctured oriented surface F and that the map $G \rightarrow \mathrm{Sp} \mathcal{L}$ is induced by a certain representation of G in the mapping class group of F . In these settings, a weak Hurwitz equivalence class of a G -valued BM-factorization $\bar{\mathfrak{m}}$ of length r represents an F -bundle $X \rightarrow B^{\sharp}$ over a disk B^{\sharp} with r punctures, see 1.1.2, one has $\mathcal{L}_{\bar{\mathfrak{m}}} = H_2(X)$, and the symmetric bilinear form $q: \mathcal{L}_{\bar{\mathfrak{m}}} \otimes \mathcal{L}_{\bar{\mathfrak{m}}} \rightarrow \mathbb{Z}$ is given by the intersection index, $q: x \otimes y \mapsto x \circ y$. (Definition 7.1.1 is merely a generalization of a simple algorithm computing $H_2(X)$ and the self-intersections of 2-cycles.) The group $\mathcal{L} \otimes \bar{\mathfrak{m}}$ can be interpreted as $H_2(X, F_b)$, where F_b is the fiber over a point $b \in \partial B^{\sharp}$, but the form $q: \mathcal{L} \otimes \bar{\mathfrak{m}} \rightarrow \mathbb{Z}$ does not seem to have a geometric meaning. Examples show that the associated bilinear form does not need to be divisible by 2, see [12] or Example 7.2.6 below.

7.1.5. Definition. A (*weak*) *isomorphism* between two triples $(\mathcal{M}_1; \chi_1, q_1)$ and $(\mathcal{M}_2; \chi_2, q_2)$, where \mathcal{M}_i is an R -module, $\chi_i: \mathcal{M}_i \rightarrow \mathcal{L}$ is an R -linear map, and $q_i: \mathcal{M}_i \rightarrow \mathcal{V}$ is an R -quadratic map, is an R -isomorphism $\varphi: \mathcal{M}_1 \rightarrow \mathcal{M}_2$ such that $q_1 = q_2 \circ \varphi$ and $\chi_1 = \chi_2 \circ \varphi$ (respectively, $\chi_1 = g \circ \chi_2 \circ \varphi$ for some $g \in G$).

7.1.6. Proposition. *The triples $(\mathcal{L} \otimes \bar{\mathfrak{m}}; \chi, q)$ and $(\mathcal{L} \otimes \bar{\mathfrak{m}}'; \chi', q')$ corresponding to two strongly (respectively, weakly) equivalent BM-factorizations $\bar{\mathfrak{m}}$ and $\bar{\mathfrak{m}}'$ are isomorphic (respectively, weakly isomorphic). In particular, the transcendental lattice $q: \mathcal{T}(\bar{\mathfrak{m}}) \rightarrow \mathcal{V}$ is a weak equivalence invariant of $\bar{\mathfrak{m}}$.*

Proof. If $\bar{\mathfrak{m}}'$ is obtained from $\bar{\mathfrak{m}}$ by a global conjugation, $\mathfrak{m}'_i = g^{-1}\mathfrak{m}_i g$, $g \in G$, the weak isomorphism $\mathcal{L} \otimes \bar{\mathfrak{m}}' \rightarrow \mathcal{L} \otimes \bar{\mathfrak{m}}$ is $\varphi: \bigoplus x'_i \mapsto \bigoplus g x'_i$; then $\chi' = g^{-1} \circ \chi \circ \varphi$.

Assume that $\bar{\mathfrak{m}}'$ is obtained from $\bar{\mathfrak{m}}$ by one inverse Hurwitz move,

$$\mathfrak{m}'_i = \mathfrak{m}_{i+1}, \quad \mathfrak{m}'_{i+1} = \mathfrak{m}_{i+1}\mathfrak{m}_i\mathfrak{m}_{i+1}^{-1}, \quad \mathfrak{m}'_j = \mathfrak{m}_j, \quad j \neq i, i+1.$$

Then the isomorphism $\varphi: \bigoplus x'_i \mapsto \bigoplus x_i$ is given by

$$x_i = \mathfrak{m}_{i+1}^{-1}x'_{i+1}, \quad x_{i+1} = x'_i + (\mathfrak{m}_i - 1)\mathfrak{m}_{i+1}^{-1}x'_{i+1}, \quad x_j = x'_j, \quad j \neq i, i+1.$$

It is straightforward that

$$(7.1.7) \quad (\mathfrak{m}_i - 1)x_i + (\mathfrak{m}_{i+1} - 1)x_{i+1} = (\mathfrak{m}'_i - 1)x'_i + (\mathfrak{m}'_{i+1} - 1)x'_{i+1};$$

hence $\chi' = \chi \circ \varphi$. Furthermore, due to (7.1.7), the essentially different terms in the expressions for q and q' are

$$-x_i \cdot \mathfrak{m}_i x_i - x_{i+1} \cdot \mathfrak{m}_{i+1} x_{i+1} + (\mathfrak{m}_i - 1)x_i \cdot (\mathfrak{m}_{i+1} - 1)x_{i+1}$$

(and the corresponding primed terms). Rewrite the latter sum in the form

$$-x_i \cdot \mathfrak{m}_i x_i + [(\mathfrak{m}_i - 1)x_i - x_{i+1}] \cdot (\mathfrak{m}_{i+1} - 1)x_{i+1}$$

(using $x_{i+1} \cdot x_{i+1} = 0$) and observe that $(\mathfrak{m}_i - 1)x_i - x_{i+1} = -x'_i$ and

$$(\mathfrak{m}_{i+1} - 1)x_{i+1} = (\mathfrak{m}'_i - 1)x'_i + (\mathfrak{m}'_{i+1} - 1)x'_{i+1} - \mathfrak{m}_{i+1}^{-1}(\mathfrak{m}'_{i+1} - 1)x'_{i+1}.$$

Multiplying out and using the fact that \cdot is skew-symmetric and $\mathfrak{m}_{i+1} = \mathfrak{m}'_i$ is an isometry, one obtains $q' = q \circ \varphi$. \square

7.2. Examples and open questions. The transcendental lattice $q: \mathcal{T}(\bar{\mathfrak{m}}) \rightarrow \mathcal{V}$ is a relatively new invariant (regarded as an invariant of a BM-factorization) and I do not know how powerful it is. In particular, I do not know if it can be expressed in terms of other known invariants.

7.2.1. Problem. Is there a relation between $\mathcal{T}(\bar{\mathfrak{m}})$ and other known invariants, for example $[\text{Im}(\bar{\mathfrak{m}})]$ and $[\mathfrak{m}_\infty(\bar{\mathfrak{m}})]$?

Most known examples of computation of $\mathcal{T}(\bar{\mathfrak{m}})$ use the identity representation $\tilde{\Gamma} = \text{Sp } \mathcal{H}$, see 2.1, and deal with a BM-factorization representing the homological invariant of an extremal elliptic surfaces X . In this case, \mathcal{T} is indeed the transcendental lattice of X , *i.e.*, the orthogonal complement $\text{NS}(X)^\perp \subset H_2(X)$, with the form induced by the intersection index; this relation explains the terminology, and it is the computation in [12] that inspired Definition 7.1.1.

7.2.2. Example. The $\tilde{\Gamma}$ -valued reductions of the (non-simple) BM-factorizations arising from the pairs of plane sextics with the sets of singularities marked with a * in Table 2, see Subsection 5.6, differ by their transcendental lattices. An easy way to prove this fact is to compare the geometric classification of curves found in [10], [11] and their arithmetic classification found in [27]. The same argument shows that the other pairs in Table 2 have isomorphic transcendental lattices.

7.2.3. Example. For each $k \geq 0$, the simple $\tilde{\Gamma}$ -valued BM-factorizations given by Theorem 1.2.1 have isomorphic transcendental lattices, see [12]. If k is even, one has $\mathcal{T} \cong \mathbf{D}_k$ (with the usual convention $\mathbf{D}_0 = 0$ and $\mathbf{D}_2 = 2\mathbf{A}_1$); if $k = 2s - 1$ is odd, then \mathcal{T} is the orthogonal complement $(3\mathbf{v}_1 + \dots + 3\mathbf{v}_s + \mathbf{v}_{s+1} + \dots + \mathbf{v}_{2s-1})^\perp$ in the orthogonal direct sum $\bigoplus_{i=1}^{2s-1} \mathbb{Z}\mathbf{v}_i$, $\mathbf{v}_1^2 = 1$.

7.2.4. A *coloring* of length r is a function $\ell: \{1, \dots, r\} \rightarrow \{\pm 1\}$. Given a simple Γ -valued BM-factorization $\tilde{\mathbf{m}} = (\mathbf{m}_1, \dots, \mathbf{m}_r)$ and a coloring ℓ of length r , define the lattice $\mathcal{T}(\tilde{\mathbf{m}}, \ell)$ as the transcendental lattice of the $\tilde{\Gamma}$ -valued lift of $\tilde{\mathbf{m}}$ obtained as follows: an entry $\mathbf{m}_i = g_i^{-1} \mathbb{X} \mathbb{Y} g_i$, $i = 1, \dots, r$, $g_i \in \Gamma$, lifts to $g_i^{-1} \ell(i) \mathbb{X} \mathbb{Y} g_i \subset \tilde{\Gamma}$. Alternatively, this lift can be described as the one with the eigenvalues of sign $\ell(i)$; in this form, the concept can be extended to a wider class of BM-factorizations, for example, to those with unipotent entries, which arise from elliptic surfaces/trigonal curves over the rational base and without exceptional singular fibers. The following statement is immediate.

7.2.5. Proposition. *Assume that two simple Γ -valued BM-factorizations $\tilde{\mathbf{m}}'$, $\tilde{\mathbf{m}}''$ of length r are weakly equivalent. Then there is a permutation $\sigma \in \mathbb{S}_r$ such that, for any coloring ℓ of length r , one has $\mathcal{T}(\tilde{\mathbf{m}}', \ell) \cong \mathcal{T}(\tilde{\mathbf{m}}'', \ell \circ \sigma)$. \square*

7.2.6. Example. In [12], the lattices $\mathcal{T}(\tilde{\mathbf{m}}, \ell)$ are computed for all Γ -valued BM-factorizations given by Theorem 1.2.1, see (5.2.2), and all colorings ℓ taking exactly one value -1 . It turns out that the isomorphism class of $\mathcal{T}(\tilde{\mathbf{m}}, \ell)$ depends on k only. The corresponding quadratic forms $q: \mathcal{H} \otimes \tilde{\mathbf{m}} \rightarrow \mathbb{Z}$ are also computed; in general, they do *not* extend to integral symmetric bilinear forms.

7.2.7. Problem. Does Proposition 7.2.5 distinguish the weak equivalence classes given by Theorem 1.2.2?

7.2.8. Example. We conclude with the only known to me example of a direct computation of the transcendental lattice using a representation other than $\tilde{\Gamma} = \mathrm{Sp} \mathcal{H}$. In [1], the authors give an explicit construction of a pair of reducible sextics (each splitting into an irreducible quintic Q and a line L) with the set of singularities $\mathbf{A}_{10} \oplus \mathbf{A}_9$ and compute their \mathbb{B}_5 -valued braid monodromies with respect to the pencil of lines through a generic point in L . Then, following more or less the lines of Definition 7.1.1 and using the obvious representation $\mathbb{B}_5 \rightarrow \mathrm{Sp} H_1(F)$, where F is a punctured surface of genus 2, they compute the transcendental lattices and show that they are distinct (the latter fact being predicted beforehand using theory of K3-surfaces). It is worth mentioning that the two sextics are conjugate over $\mathbb{Q}(\sqrt{5})$; thus, \mathcal{T} is a topological, but not algebraic, invariant.

REFERENCES

1. K. Arima, I. Shimada, *Zariski–van Kampen method and transcendental lattices of certain singular K3 surfaces*, [arXiv:0806.3311](https://arxiv.org/abs/0806.3311) (to appear).

2. E. Artal, J. Carmona, and J. I. Cogolludo, *Effective invariants of braid monodromy*, Trans. Amer. Math. Soc. **359** (2007), no. 1, 165–183.
3. E. Artin, *Theory of braids*, Abh. Math. Sem. Hamburgischen Univ. **4** (1926), 47–72.
4. B. Birch, *Noncongruence subgroups, covers and drawings*, The Grothendieck theory of dessins d'enfants (Luminy, 1993), London Math. Soc. Lecture Note Ser., vol. 200, Cambridge Univ. Press, Cambridge, 1994, pp. 25–46.
5. F. Bogomolov, Yu. Tschinkel, *Monodromy of elliptic surfaces*, Galois Groups and Fundamental Groups (L. Schneps, ed.), MSRI Publications, vol. 41, Cambridge Univ. Press, 2003, pp. 167–181.
6. O. Chisini, *Una suggestiva rappresentazione reale per le curve algebriche piane*, Rend. Ist. Lombardo II **66** (1933), 1141–1155.
7. O. Chisini, *Courbes de diramation des plans multiples et tresses algébriques*, Deuxième Colloque de Géométrie Algébrique, Masson, Paris 1952, pp. 11–27.
8. A. Degtyarev, *Zariski k -plets via dessins d'enfants*, Comment. Math. Helv. **84** (2009), no. 3, 639–671.
9. A. Degtyarev, *Plane sextics via dessins d'enfants*, Geom. Topol. **14** (2010), no. 1, 393–433.
10. A. Degtyarev, *Plane sextics with a type E_8 singular point*, arXiv:0902.2281.
11. A. Degtyarev, *Plane sextics with a type E_6 singular point*, arXiv:0907.4714.
12. A. Degtyarev, *Transcendental lattice of an extremal elliptic surface*, arXiv:0907.1809.
13. A. Degtyarev, I. Itenberg, V. Kharlamov, *On deformation types of real elliptic surfaces*, Amer. J. Math. **130** (2008), no. 6, 1561–1627.
14. R. Friedman, J. W. Morgan, *Smooth four-manifolds and complex surfaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3), Springer-Verlag, Berlin-New York, 1994.
15. The GAP Group, *GAP—Groups, Algorithms, and Programming*, Version 4.4.10, 2007, (<http://www.gap-system.org>).
16. A. Hurwitz, *Über Riemannsche Flächen mit gegebenen Verzweigungspuncten*, Math. Ann. **38** (1891), 1–61.
17. E. R. van Kampen, *On the fundamental group of an algebraic curve*, Amer. J. Math. **55** (1933), 255–260.
18. V. M. Kharlamov, Vik. S. Kulikov, *On braid monodromy factorizations*, Izv. Ross. Akad. Nauk Ser. Mat. **67** (2003), no. 3, 79–118 (Russian); translation in: Izv. Math. **67** (2003), no. 3, 499–534.
19. K. Kodaira, *On compact analytic surfaces, II–III*, Annals of Math. vol 77–78 (1963), 563–626, 1–40.
20. Vik. S. Kulikov, *Hurwitz curves*, Uspekhi Mat. Nauk **62** (2007), no. 6(378), 3–86 (Russian); English transl. in Russian Math. Surveys **62** (2007), no. 6, 1043–1119.
21. B. Moishezon, *Complex surfaces and connected sums of complex projective planes*, Lecture Notes in Math., vol. 603, Springer-Verlag, Berlin-New York, 1977.
22. B. Moishezon, *Stable branch curves and braid monodromies*, Algebraic geometry (Chicago, Ill., 1980), Lecture Notes in Math., vol. 862, Springer, Berlin-New York, 1981, pp. 107–192.
23. M. Nori, *On certain elliptic surfaces with maximal Picard number*, Topology **24** (1985), no. 2, 175–186.
24. S. Orevkov, *Riemann existence theorem and construction of real algebraic curves*, Ann. Fac. Sci. Toulouse Math. (6) **12** (2003), no. 4, 517–531.
25. S. Orevkov, *On braid monodromy monoid*, Talk at MSRI, 2004, <http://www.msri.org/publications/ln/msri/2004/topgeom/orevkov/1/index.html>.
26. N. Salepci, *Real Lefschetz fibrations*, Ph. D. thesis, IRMA, Strasbourg, 2007, pp. 136.
27. I. Shimada, *On the connected components of the moduli of polarized $K3$ surfaces* (to appear).
28. T. Shioda, *On elliptic modular surfaces*, J. Math. Soc. Japan **24** (1972), 20–59.
29. O. Zariski, *On the problem of existence of algebraic functions of two variables possessing a given branch curve*, Amer. J. Math. **51** (1929), 305–328.

DEPARTMENT OF MATHEMATICS, BILKENT UNIVERSITY, 06800 ANKARA, TURKEY

E-mail address: degt@fen.bilkent.edu.tr